



5.455.

# JOURNAL

OF

# NATURAL PHILOSOPHY,

CHEMISTRY,

AND

THE ARTS.

VOL. III.

Illustrated with Engravings.

BY WILLIAM NICHOLSON.

#### LONDON:

PRINTED BY W. STRATFORD, CROWN-COURT, TEMPLE-BAR; FOR
THE AUTHOR, No. 10, SOHO-SQUARE
AND SOLD BY

G. AND J. ROBINSONS, PATERNOSTER-ROW.

1802.

# TAUTO



# PREFACE.

THE third Volume of this Miscellany being now completed, I have again the opportunity of making my acknowledgements to its friends and supporters, and presenting the respectable list of its Authors.

The authors of original Papers are John Bostock, M. D. Mr. Rob. Jameson; the Rev. James Wilson, A.M.; John Gough, Esq.; Mr. Ez. Walker; Rev. Joseph Priestley, L.L.D. F.R.S.; Rev. W. Pearson, P. R. I.; Dr. Thomas Young, F. R. S. and Professor at the Royal Institution; Mr. Peter Keir; Mr. John Dalton; Mr. Wm. Wilson; R. Chenevix, Esq. F. R. S.; Mr. John Cuthbertson; Mr. John Murray; J. B.; W. N.; and Professor Aldini.—Of foreign works; Hapel Lachenaie; Brongniart; J. B. Berard; Guyton; Gay Lussac; Ekeberg; B. Prevost.—And of English Memoirs abridged or extracted, Dr. Young; Mr. R. Phillips; Mr. J. Besant; Edw. Howard, Esq.; F. R.S.; Sir H. C. Englesield, Bart. F. R. S.; Mr. J. Dalton; Mr. H. Davy, Professor at the Royal Institution; T. Wedgwood, Esq.; R. Chenevix, Esq. F. R. S.

Of the Engravings the Subjects are, 1. A powerful Blowpipe by Alcohol. 2. Mr. Wilson's two new Barometers. 3. An improved Water Wheel, by Mr. Besant. 4. Mr. R. Phillips's Tubes for driving Copper Bolts into Ships. 5. The Hydrometer of Atkins, for exhibiting Specific Gravities and Strengths

#### PREFACE.

Strengths of Spirits without Computation. 6. Mr. Read's very simple and cheap Pneumatic Apparatus. 7. The new moveable Crane erected by Mr. Keir at Ramsgate. 8. Haas's Blow-pipe Apparatus. 9. Figures by Mr. Jameson, to illustrate the Formation of Basalt. 10. Elements of the Galvanic Pile, by Dr. Bostock. 11. Distinguishers of Galvanism and Electricity. 12. Gay Lussac's Apparatus for measuring the Expansions of the Gases. 13. The Indian Hand Mill. 14. A Secret Lock, which can be opened only by the Proprietor. 15. Improvement of Woulfe's Apparatus, by Mr. Murray. 16. Dr. Young's Apparatus for exhibiting the colours of thin plates. 17. Figures by Prevost, for explaining the Phenomena of Dew.

Soho Square, Dec. 26, 1802.

## TABLE OF CONTENTS

#### TO THIS THIRD VOLUME.

#### SEFTEMBER 1802.

- FINGRAVINGS of the following Objects: 1. An Apparatus in which the Flame of a Lamp is very powerfully urged by the Vapour of Alcohol; 2. Two new Barometers, by the Rev. James Wilson, A. M.; 3. Mr. Besant's improved Water Wheel; 4. Mr. R. Phillips's Tubes for driving Copper Bolts into Ships; 5. An Hydrometer by Mr. Atkins, by which Specific Gravities are exhibited directly without Computation; 6. Very simple and cheap Pneumatic Apparatus, by Mr. Read.
- I. Description of a cheap and simple Apparatus or Blow-pipe, in which the Flame of Oil or Tallow is impelled by the Vapour of Alcohol. From a Correspondent.

  Page 1
- II. Outline of the History of Galvanism: with a Theory of the Action of the Galvanic Apparatus. By John Bostock, M. D. From the Author. 3
- III. On the supposed Existence of Mechanical Deposits and Petrifactions, in the Primitive Mountains, and an Account of Petrifactions which have been discovered in the newest Flotz Trapp Formation. By Mr. Robert Jameson.

  Communicated by the Author.
- IV. A Method of increasing the Sensibility of the Barometer, ad libitum.

  By the Rev. James Wilson, A. M. Communicated by the Author. 21
- V. A Summary of the most useful Parts of Hydraulics, chiefly extracted and abridged from Eytelwein's Handbuch der Mechanik und der Hydraulik. Berlin, 1801. By Thomas Young, M. D. F. R. S. 25
- VI. Experimental Proof, that Corrections deduced from the Arcs of Vibrations of a Pendulum in Vacuo, are practically useful. By Mr. Ezekiel Walker.
- VII. Description of an Apparatus of Tubes for facilitating the driving of Copper Bolts into Ships. By Mr. Richard Phillips. 36
- VIII. Reply to Dr. Young's Letter on the Theory of Compound Sounds. In a Letter from Mr. John Gough.
- IX. A new Process for Claying Sugars, proposed by Cit. Hapel Lachenaie,
  Chief Apothecary of the Military Hospitals of Guadaloupe to the Agents of the
  Consuls of the French Republic in the Windward Islands.

  41
- X. Description of a Water Wheel. By Mr. J. Besant. In a Letter to Mr. Charles Taylor, Secretary to the Society for the Encouragement of Arts. From their Transactions.
- XI. Description of an Hydrometer which gives the Specific Gravity by Inspection, constructed by Mr. Atkins. W. N. 50
- XII. On the Air from finery Cinder and Charcoal, with other Remarks on the Experiments and Observations of Mr. Cruickshank. In a Letter from Joseph Priestley, L. L. D. F. R. S. &c. &c. 52
- XIII. Description of Mr. Read's Pneumatic Apparatus. By a Correspondent.
- Scientific News. Experiments of the Invifible Girl—Aerostatic experiments of Garnerin—Commercial College at Hull—A System of Chemistry in four Volumes—Death of Thomas Garnet, M. D. late Professor of Chemistry and Natural Philosophy at the Royal Institution; &c. &c. 56 to 63

Vol. III,-1802. a OCTOBER

#### OCTOBER 1802.

- Engravings of the following Objects: 1. View and Plan of a new moveable Crane, erected for Ramigate Harbour, by Mr. Keir; 2. Haas's Blow-pipe Apparatus; 3. Figures to illustrate the Formation of Basalt, &c. by Mr. Jameson; 4. Elements of the Galvanic Pile, by Dr. Bostock; and, 5. Distinguishers of Galvanish and Electricity.
- I. Observations upon some remarkable Wells near the Sea Coast at Brighthelmstone, and other Places contiguous. By the Rev. W. Pearson, P.R.I. - - Page 65
- II. On the Theory of Galvanism. By John Bostock, M. D. From the Author.
- 111. A Summary of the most useful Parts of Hydraulics, chiefly extracted and abridged from Eytelwein's Handbuch der Mechanik und der Hydraulik. Berlin, 1801. By Thomas Young, M. D. F. R. S.
- IV. Experiments and Observations on certain stony and metalline Substances, which at different Times are said to have fallen on the Earth; also on various Kinds of native Iron. By Edward Howard, Esq. F. R. S. From the Philosophical Transactions, 1802.
- V. On the Colours obtained from Metallic Oxides, and fixed by Means of Fufiou on different Vitreous Bodies. By Alex. Brongniart, Director of the National Manufactory of Porcelain at Sévres, Engineer of Mines, &c. 101
- VI. Examination of the supposed igneous Origin of the Rocks of the Trappformation. By Mr. Robert Jameson. From the Author.
- VII. Description of the Blow-Pipe Apparatus of Haas. From a Correspondent.
- VIII. Note respecting the Instruments by which the two Kinds of Electricity are distinguished, or its Direction ascertained. W. N. 121
- IX. Section and Plan of a moveable Crane, capable of heaving four Tons with four Men. Defigned and executed for the Pier at Ramfgate, by Order of the Trustees for the Management of the Harbour at that Place. By Mr. Peter Keir, of St. Pancras. Communicated by Mr. Peter Keir. 124
- X. Experiments on the Separation of Light and Heat by Refraction. In a Letter from Sir H. C. Englefield, Bart. F. R. S. to Thomas Young, M. D. F. R. S. From the Journal of the Royal Institution, p. 202.
- XI. On the Expansion of the Elastic Fluids by Heat. By Mr. John Dalton. - - - 130
- XII. Account of fome Experiments made in the Laboratory of the Royal Institution, relating to the Agencies of Galvanic Electricity, in producing Heat, and in effecting Changes in different fluid Substances. By H. Davy, Prof. Chem.
- Scientific News. National Inftitute of France—Prize proposed at the Public Sitting of the 15 Germinal, in the 10th Year. (April 5, 1802.)—Subject of the Prize of Chemistry—Prize Questions from the Batavian Society at Haarlem—Sixth Descent of Garnerin with his-Parachute. 139 to 143

#### NOVEMBER 1802.

- Engravings of the following Objects: 1. Cit. Gay Lussac's Apparatus for determining the Expansions of the Gases by Heat; 2. The Indian Hand Mill; 3. and 4. The Parts and Structure of a Secret Lock, which can be opened only by the Proprietor. By Cit. J. B. Berard.
- I. Letter from Thomas Young, M. D. F. R. S. In Reply to Mr. Gough's Letter, at Page 36 of the present Volume. On the Phenomena of Sound.

  Page 145
- II. On the Galvanic Effect of very minute Particles of Zinc and Copper in Water. In a Letter from Mr. William Wilson.
- III. On the Colours obtained from Metallic Oxides, and fixed by Means of Fusion on different Vitreous Bodies. By Alex. Brongniart, Director of the National Manufactory of Porcelain at Sévres, Engineer of Mines, &c. 148
- IV. Experiments and Observations on certain Stony and Metalline Substances, which at different Times are said to have fallen on the Earth; also on various Kinds of Native Iron. By Edward Howard, Esq. F. R.S. From the Philosophical Transactions, 1802.
- V. Experiments and Observations on the Heat and Cold produced by the Mechanical Condensation and Rarefaction of Air. By John Dalton 160
- VI. An Account of a Method of copying Paintings upon Glass, and of making Profiles by the Agency of Light upon Nitrate of Silver. Invented by T. Wedgwood, Esq. With Observations by H. Davy. From the Journals of the Royal Institution, I, 173.
- VII. Observations and Experiments upon Oxigenized and Hyperoxigenized Muriatic Acid; and upon some Combinations of the Muriatic Acid in its Three States. By Richard Chenevix, Esq. F. R. S. M. R. I. A. From the Philosophical Transactions for 1802.
- VIII. Description of the Indian Hand Mill. From a Correspondent. 186
- IX. An Effect hitherto unpublished, by which the two Electricities are distinguished. In a Letter from Mr. John Cuthbertson. 183
- X. Palpable Mathematics; or Methods of writing and calculating for the Use of the Blind. By J. B. Berard.
- XI. Observations on the real Nature of the Precipitates formed by the Prusfiates in Acid Solutions of Barites, and the Affinities of the Prussic Acid. By Citizen Guyton. - - 193
- XII. A Review of some Experiments, which have been supposed to disprove the Materiality of Heat. By William Henry. From the Memoirs of the Literary and Philosophical Society of Manchester, Vol. V. Part II.
- XIII. Enquiries concerning the Dilatation of the Gases and Vapors. Read to the National Institute of France. By Cit. Gay Lussac. 207
- XIV. Description of a new Secret Lock. By J. B. Berard. 216
- Scientific News. Cheap Method of producing Light.—On the Reflection of obfcure Heat.—On certain Facts commonly urged against the Doctrine of two Electric Fluids.

   222, 223

#### DECEMBER 1802.

- Engravings of the following Objects: 1. Mr. Murray's Improvement of Woulfe's Apparatus; 2. Dr. Young's Apparatus for exhibiting the Colours of thin Plates; 3. Prevost on the Phenomena of Dew.
- I. Description of an Improvement of Woulse's Apparatus, by which the great Inconveniences of that Method of connecting the Vessels are obviated. In a Letter from Mr. John Murray.

   Page 225
- II. Observations and Experiments upon oxigenized and hyperoxigenized Muriatic Acid; and upon some Combinations of the Muriatic Acid in its three States. By Richard Chenevix, Esq. F. R. S. M. R. I. A. From the Philosophical Transactions for 1802.
- III. On the present State of the British Museum. By a Correspondent. 246
- 1V. Of the Properties of the Earth Yttria, compared with those of Glucine; of Fossils, in which the first of these earths is contained; and of the Discovery of a new Substance of a metallic nature (Tantalium.) By A. G. Ekeberg.
- V. On the Probability that the Eruptions of Lunar Volcanos may reach the Earth. By J. B. - 255
- VI. Enquiries concerning the Dilatation of the Gases and Vapors. Read to the National Institute of France. By Cit. Gay Lussac. 257
- VII. New Theory of the Constitution of mixed Gases elucidated. In a Letter from Mr. J. Dalton. - 267
- VIII. A Method of increasing the Quantity of Light afforded by Candles, and to obviate the Necessity of snussing them. In a Letter from Mr. Ezekiel Walker.
- IX. Description of a new Secret Lock. By J. B. Berard. Concluded from page 222.
- X. Description of Dr. Young's Apparatus for exhibiting the Colours of thin Plates, by means of the Solar Microscope. From the Journals of the Royal Institution, I. 241.
- XI. Extract of a Letter from R. Chenevix, Esq. on the Magnetic Property of Nickel. - 286
- XII. On the unexpected Production of Sulphate of Magnefia. By J. Bostock,
  M. D. - - 288
- XIII. Abstract of a Memoir on the Dew. By Cit. Benedict Prevost. 290
- Scientific News, &c. Abstract of the late Experiments of Professor Aldini on Galvanism.—Examen, &c. Chemical Examination of a New Gas.—Notice concerning certain peculiar Habits of the Shark and Pilot fish.—Notice of a new Metal.

  298 to 304

# JOURNAL

0 F

## NATURAL PHILOSOPHY, CHEMISTRY,

AND

#### THE ARTS.

SEPTEMBER, 1802

#### ARTICLE I.

Description of a cheap and simple Apparatus or Blow-pipe, in which the Flame of Oil or Tallow is impelled by the Vapour of Alcohol-From a Correspondent.

#### To Mr. NICHOLSON.

SIR.

London, July 9, 1802.

SEFUL as the blow-pipe is for the professed mineralogist Inconveniences and chemist, as well as for the amateur of the science, it is, of the blow-pipe, and the substihowever, liable to the inconvenience of requiring one hand of tutes commonly the operator to be employed; besides, that persons of weak used. lungs, or not well accustomed to its use, find it laborious, even when they attempt to manage it according to the directions given by Bergman and Engstrom; more especially, when a continued stream of air is required for some time. For this reafon, fome have recourse to a bladder filled with air, and force it out by pressing the bladder between the arm and their body; others use a small pair of double bellows, to be fixed upon the table: others again employ a receiver, or rather gazometer, in the form of Mr. Watt's hydraulic bellows. But all these means are not always ready at hand, or not eafily portable to accom-Vol. III .- SEPTEMBER, 1802. pany

rany the travelling mineralogist, and in every case they require fome preparation. And even the ingenious contrivance of Mr. Haas, though it permits the free use of the operator's hands, is expensive, and still requires the blowing through the mouth.

Very afteful lamp exhibited by Professor Pictet.

When Professor Pictet of Geneva visited London last year, he brought along with him and exhibited to his friends an apparatus, which at the same time that it is simple and very portable, is the most advantageous substitute for the blow-pipe, and removes all its inconveniences. An imitation of this apparatus was exhibited in the lecture-room of the Royal Institution, Albemarle-street.

Description of two lamps for blowing the flame of oil by alcohol. The drawing of this inftrument which I subjoin, was copied from that communicated by a friend. A, plate 1, is a tin-box filled with common oil, serving to feed the flame at F, thence to be directed on the substance under examination, and at the other flame, to heat the spirit of wine contained in the copper or tin lamp B. The boiling spirit being converted into steam by the heat, rushes through the bended tube C upon the flame of the wick at F. Lastly, D is a slider with loops and screws to raise or lower the spirit lamp. The whole being very simple and clear, any farther explanation would be needless.

Long ago ufed by Nollet. I do not know whether Professor Pictet pretends to be the inventor of this apparatus; but certain it is, that Abbe Nollet more than sifty years ago employed such a contrivance, as may be seen in his VArt de faire Experiences. Another application of the spirit lamp for this purpose I have read in the Annales de Chimie, but do not recollect in which volume.

There is no doubt but the communication of the lamp above described will be acceptable to many of the readers of your excellent Journal; whence, if you agree with me in this opinion, be so kind as to insert it.

I am, Sir,

Your humble fervant.

N. N.

#### ANNOTATION. W. N.

Powerful effect of a fmall colipile. The application of the elastic and combustible vapour of alcohol to this useful puppose is so obvious, that it can scarcely be appropriated as an invention. The glass-blowers have long sold a little implement, consisting of an egg-shaped ball, having

I frem or handle, and a nozzle or tube. It is an eolipile, and when half filled with water or spirit, and heated over a candle, a ftrong current either of vapour or of the dense fluid will issue out accordingly as it is held. With the vapour of spirit thus blown An halfpenny through the large flame of a tallow lamp, I fused a halfpenny, glass stopper perand softened the stopper of a bottle so as to run a wire through forated. it; though I must confess not without apprehension of the thin brittle veffel burfling, and covering the furrounding bodies with flame. The lamp B, which is the effential part, may be made The lamp appacheap; and indeed the whole would cost but a trifle. I took ratus is very a small lamp of that figure,  $2\frac{1}{2}$  inches wide and  $1\frac{1}{2}$  inch high (of the fort called agitable, from the cap for the wick being made to screw in and defend the oil), and in the place of the wick-pipe, I foldered a piece of a common blow-pipe. The lamp cost eighteen pence, and the blow-pipe fix-pence, and the

#### II.

combination answers perfectly well.

Outline of the History of Galvanism: with a Theory of the Action of the Galvanic Apparatus. By JOHN BOSTOCK, M. D. From the Author:

### (Concluded from Page 304.)

COL. HALDANE found that the effects of the apparatus Col. Haldane were suspended when it was immersed under water, and that immersed the this was also the case when it was placed under the vacuum of He discovered an air-pump. He conceived that the effects of the pile were that its action is not increased by its being connected with the conductor of an vacuo. electrical machine, and he thought that the charging of the Leyden phial was retarded when the apparatus was placed between its outfide and infide coatings. Upon the whole he Thinks it not concludes that its effects cannot be referred to electricity. electric. Upon these experiments Mr. Nicholson remarks, that in the Observations to case where the pile was connected with the conductor, the the contrary. electric stream of the conductor might have been in a contrary direction to the electric stream of the pile. In the experiment where the pile was connected with the Leyden phial, it is certain that the phial could not be charged higher than the pile B 2 itfelf.

itself, and from the effects of galvanism in general, it may be conjectured that it refembles a large furface of an electric charged to a low intenfity.

Cruickshank's for keeping the gases distinct.

Mr. Cruickshank, in a fecond communication, gives an acfimple apparatus count of some experiments which he performed with an interrupted wire of gold. By means of a bent glass tube he contrived to keep the gafes obtained from each wire perfectly distinct. That from the filver end of the pile he found to be chiefly hidrogen, and that from the zinc end nearly pure oxigen. Both the gold wire and one which he used of platina were evidently tarnished on the zinc end, after being exposed His general con- for fome time to the influence. He draws the following geneclusions. 1. Sil-ral conclusions from his experiments. 1st, From the filver end,

ver end of the pile gives chiefly if it be immerfed in water, whatever be the nature of the wire, tion.

hidrogen. 2. Re- there is disengaged a quantity of hidrogen gas mixed with a duction of me-tallic folution at small quantity of oxigen gas and ammonia. 2d, Where a methis end. 3. Zinc tallic folution is used instead of water, the metal is reduced and end gives chiefly is deposited at the end of the wire. 3d, When the wire at the oxigen. 4. Or produces oxida- zinc end confifts of gold or platina, oxigen gas is difengaged mixed with a little azote and nitrous acid; the quantity of gas is only about one-third of what is difengaged at the filver end, fo that the gases exist in the proportion requisite to produce water. 4th, When the zinc wire is formed of an imperfect metal, it is partly oxidated, partly diffolved, and a little oxigen only is difengaged. Fluids which contain no oxigen Mr. Cruickshank thinks are incapable of transmitting the influence, whilst all those which contain it appear capable of conducting His Invention of the influence. Mr. Cruickfhank then proceeds to give an account of a new method of placing the metals horizontally in a

kind of trough, which in some respects appears to be more con-

venient than the form of the apparatus originally invented by

a new galvanic arrangement by a trough.

Volta. Mr. Davy proposed this curious subject of enquiry, whether tains the oxigen the oxigen and hidrogen gases evolved by the interrupted wire of water, and the when immerfed in water, can be procured from two portions of water kept distinct from each other? For this purpose he caused the wires to terminate in two glasses of water, and by means of his fingers established a communication between the two glasses. The gases he found were produced as in the for-When the water used was very carefully boiled, he mer cafe. found that the gafes evolved were nearly pure, and in the pro-

portions

Mr. Davy obfrom one portion hidrogen from another unconnected portion.

portions requifite to form water. Mr. Davy then wished to Reverse effect ascertain whether it was essential for the wires to be in contact on a single inter-mediate wire. with the metallic plates of the pile; the ends of the pile were therefore made to communicate with two glasses of water by pieces of mufcular fibre, and the glasses were connected by a filter wire. The effects of the pile were now reversed. Hidrogen was difengaged by the wire at the zinc end, while the wire at the filver end became oxidated.

Col. Haldane, in a fecond memoir, details a feries of experi- Col. Haldane ments made to determine the power of different combinations flews that the pile acts more of metals. He found the pile acted more powerfully when im-frongly in oxmerfed in a given quantity of oxigen gas, than when confined in igen, and not at the same bulk of atmospherical air; its action was entirely suf-vacuo. pended in azote or in a vacuum. On this account he conjectures that the effects of galvanism depend upon a chemical operation, and that oxigen is attracted by the apparatus from the atmosphere.

Mr. Davy discovered that the influence of the pile was Davy's applicacapable of being transmitted through charcoal, when an appa-tion of charcoal ratus was conftructed of this fubftance and filver. It decomposed water in the usual manner, the filver end giving out hidrogen with a little carbon in folution, while the zinc end evolved but little gas of any kind; this was probably owing to the abforption of the carbonic acid which would be formed at this end of the wire. He found that when the water between the His discoveries pairs of metallic plates was perfectly pure, i. e. when it that pure water is ineffectual in contained neither acid, falt, nor gas of any kind, the pile the pile, and that ceased to act. He conjectures that the energy of the pile is its effects are proportioned to nearly in proportion to the rapidity with which the zinc be-the oxidation. comes oxidated; and confequently the effects were found to be the most powerful when nitric acid was interposed between the metals. Mr. Davy appears in this inftance to have made He applies the the first step towards the true theory of the action of Volta's acids, pile: as might be inferred from this notion, he found that concentrated fulphuric acid interposed between the metals, acted less powerfully than diluted sulphuric acid, because this last has more power in oxidating zinc than when in its pure state. He and causes the also discovered that the pile can act in vacuo, provided a little pile to act in acid be interposed between the plates. A pile composed of zine and charcoal was found to possess great energy.

All these valuable discoveries and experiments are taken from original papers communicated to the 4th vol. of Nicholfon's Journal.

Anonym. that galvanism than men.

In the 5th vol. of this work we meet with fome curious obwomen are more fervations detailed by an anonymous correspondent at Edinburgh. He found that water which had been for some time fubjected to the influence of the pile, continued for feveral days to emit gas, and to deposit a mucous matter. He has observed that when both men and women form the galvanic circuit, the women appear to receive a larger share of the shock than the Dr. Rutherford of Edinburgh conceives that vegetables are non-conductors of galvanism, and therefore in this particular that it differs from electricity.

Mr. Davy's pile with one metal only.

Mr. Davy afterwards discovered that a pile may be conftructed with only one metal, provided proper fluids be applied to its furfaces. He employed the following feries: metal, nitrous acid, water, fulphuret of potash, and then metal. This curious discovery, as Mr. Nicholson judiciously observes, completely overthrows the hypothesis of Volta.

Cruickshank charges a jar with galvanism.

Mr. Nicholfon about this time was informed by Mr. Cruickfhank, that he had at length succeeded, by using proper precautions, in charging the Leyden phial by means of the galvanic pile.

Hitherto nearly all the wonderful discoveries that had been made by means of Volta's apparatus were confined to England, and were principally due to Mr. Nicholfon, Mr. Cruickshank, Mr. William Henry, Mr. Davy, and Col. Haldane. fubject began now to attract the attention of foreigners, and we have an account of some curious experiments performed by Professor Tromsdorff. He constructed a pile of zinc and copper, and fixed a piece of gold leaf to the zinc end of the wire;

Tromfdorff burns gold leaf and other metals.

he found that it was eafily confumed, and that the fame thing Remarkable dif-happened to other metallic leaves. This experiment was recovery of Four-peated by Fourcroy, and he adds this remarkable circumstance, croy, that the shock is greatest that fix large zinc and filver plates gave only a slight shock, but when the repeti-confumed the wire rapidly; whereas the same surface of metal tion of plates is most numerous; arranged in the form of a greater number of smaller plates, afthe combustion forded a more confiderable shock, but would not burn the greatest the metal.

larger the fur-

About this time Mr. Cruickshank observed, that when the Cruickshank ob-wire of his trough apparatus terminated at the surface of the Serves two difwater,

water, the filver end emitted from the water a brush of fire, at ferent kinds of the same time that the zinc end produced only a dense spark. This circumstance induced him to suppose that the filver was in the plus and the zinc in the minus state of electricity, which indeed agrees with Bennet's original experiments, though it differs from the account given by Mr. Nicholson in his first paper, and also from the experiments of Dr. Wollaston, Dr. Van Marum, and others, as we shall see below.

Dr. Wollaston read a judicious paper to the Royal Society, Dr. Wollaston. which appeared in the Transactions for the year 1801. He Oxidation differremarks, as Mr. Davy had previously done, that the energy of the pile feems proportionate to the disposition which one of the plates has to be oxidated or acted upon by the interposed fluid. If a piece of filver and a piece of zinc be immerfed in diluted fulphuric acid, the zinc immediately begins to decompose the water and to evolve hidrogen; the filver is not acted upon, but if we bring them into contact, the filver also begins to produce hidrogen. He therefore concludes, that in the folution of a metal by an acid, electricity is always difengaged. In from the amalorder that the electrical machine should act strongly, it is ne-gam of the el. ceffary that the amalgam upon the cushion should be composed of a metal which is easily oxidated; the Doctor also found that which will not an electrical machine plunged into carbonic acid gas is incapable acid. of being excited. He concludes by the opinion which was fuggested by Mr. Nicholson, that electricity and galvanism are the same principle, but that in the latter it is generally met with in large quantity, but in a state of low intensity. Dr. Wollaston He first procoated two filver wires, leaving the ends only of them exposed; duced the galvathese wires were connected with the two conductors of an elec- water by comtrical machine, and the spark was taken from one to the other mon electricity, through a folution of copper: he found that the end of the wire connected with the negative conductor was covered with reduced copper; no change had taken place in the other end. The negative conductor appears therefore, from these experiments, to correspond with the filver end of the pile.

Mr. Davy's very curious discovery, that a galvanic pile Mr. Davy's pile might be constructed by one metal alone, provided its different with one single surfaces were exposed to the action of different sluids, has been already announced: In addition to these facts the same intelligent experimenter informs us, in the first vol. of Nicholson's Journal, N. S. that he has constructed a galvanic apparatus which

and of charcoal with no metal.

which contains no metal, and which confifts folely of pieces of charcoal, which have their different fides exposed to the action of different fluids.

Volta's account of his own theory.

In the same volume there is an account of a paper written by Volta himself; but from the opinions which are advanced in it, we may conclude that he was unacquainted with many of the most interesting circumstances which had been observed in this country, respecting the operations of the apparatus originally discovered by himself.

Dr. Van Marum charges a large battety by galvanism.

In the 40th volume of the Annales de Chimie, is a letter from Dr. Van Marum to Sig. Volta, containing an account of experiments performed in order to compare the effects of the galvanic pile with those of the great Teylerian machine at Haarlem. Both fingle jars and batteries were charged by means of the galvanic pile, and in all cases they were charged to the same degree of intensity with that which the pile itself indicated to an electrometer. When the zinc was at the top of the pile, and communicated with the infide of the jar, the electricity of the infide was found to be positive, and when the pile was reverfed, it was negative. It was found that the shocks given by the battery charged from the electrical machine, were not perceptibly different from those given by the battery when charged to an equal intenfity by the pile. constructing a pile of large plates of zinc and copper, he was enabled to fuse thick iron wires; he even succeeded in sufing a wire of platina. He found that piles which confift of the fame number of plates, but of different diameters, give equal intenfities and equal shocks; yet those with the larger plates he found to be confiderably more powerful in fufing metals.

Dr. Priestley's experiments.

In the first volume of Nicholson's Journal, N. S. is a letter from Dr. Priestley on the pile of Volta. It contains a number of interesting experiments performed by this venerable philosopher, principally with a view of establishing his favourite hypothesis of phlogiston, to which he still adheres. Some of the facts which he notices had been previously observed in England, though from his remote situation he had not the opportunity of becoming acquainted with them. He interposed four glasses of water between the two ends of the pile; the glasses were connected by means of silver wires, and the usual operations went on at the legs of the wires; but it was found that when one of the portions of water had its surface covered with

Galvanism said not to act on water under oil; oil, the action was suspended in the whole of the glasses. This not confirmed. experiment has not, however, been found to answer in England.

#### Theory of the Action of the Galvanic Apparatus.

THE phenomena of the galvanic pile, which at first view Elementary poappear almost incapable of being arranged into any systematic fitions of galvanform, may, I think, be all eafily explained by admitting the truth of the three following postulates. 1st, That the electric 1. Electricity is fluid is always liberated or generated when a metal or any ox-oxidation. 2. idable fubstance is united to oxigen; 2dly, That the electric Electricity fluid has a strong attraction for hidrogen; and, 3dly, That hidrogen. 3. when the electric fluid, in passing along a chain of conductors, Electricity passes leaves an oxidable fubstance to be conveyed through water, it (invisibly) through water, unites itself to hidrogen, from which it is again disengaged in combination when it returns to the oxidable conductor.

The first of these propositions may be considered as almost Nature of the proved by the experiments of Fabroni, Davy, and Wollaston: proofs. the fecond and third have not been directly established by experiment, but will not appear improbable when it is feen in what a fimple and eafy manner they account for all the phenomena hitherto observed. In endeavouring to explain the action of the pile, two distinct subjects of inquiry present themselves: 1st, What are the operations carried on at the ends of the wires in the interrupted circuit, as discovered by Mr. Nicholson? and, 2dly, What is the operation carried on in the body of the pile it felf?

As the current of the electric fluid appears to pass from the Theory of what zinc or plus end of the apparatus to the filver end, we must choison's tube. first endeavour to ascertain the action which takes place at the El. passes from zinc end of the wire. This in fact appears to be the difen-the z. end and decomposes the gagement of oxigen in a concentrated flate, by which the wire water: the oxitself, when oxidable, is corroded, but which, when the wire is igen either apformed of a perfect metal, is disengaged in the form of oxigen the metal: gas. This oxigen appears to be derived from the decomposition of the water in which the wire terminates, in confequence of the attraction which the electrical fluid possesses for hidrogen, and its incapacity of passing through water without being united to this fubstance, according to the second and third postulates.

The elect. and hidrogen país invisibly to the other point, where the first is absorbed and the latter becomes gas; or if the fluid reduces the metal and again forms water.

The electric fluid thus united to hidrogen, is carried to the other point of the wire, where, upon entering the oxidable conductor, it is difengaged in the form of hidrogenous gas, if water be the medium of communication; if a folution of a metallic oxide be employed; it unites with the oxide, and reduces The decomposition of water is therefore effected at the contain oxide, it zinc point alone, though the different gafes which compose it are difengaged at each of the points; and this process will continue even when the points terminate in two different portions of water, as was discovered by Mr. Davy, provided that the glasses are united by a conductor which is not oxidable.

Theory of the

This may be confidered as a general explanation of the first apparatus or pile. fubject of inquiry; we must next proceed to investigate the nature of the operation carried on in the body of the apparatus itself. In the construction of the pile there are two points which are essential to its action; 1st, That the electric fluid be disengaged; and, 2dly, That it be confined and carried forward in one direction, fo as to be concentrated in the end of the appa-

the zinc developes elect. which decomand is attracted

The oxidation of ratus. The first object is evidently attained by the oxidation of the zinc or other oxidable body employed. If both fides of the zinc were oxidated, the electric fluid would indeed be liposes the water, berated, but it would be immediately dispersed, and its effects by the hidrogen; could not be observed. As foon, however, as the electric matter is evolved, it is immediately attracted by the hidrogen, which is at the same time necessarily generated in the fluid

and therefore passes to the filver through the fluid, and not the reverse. Its passage to the next oxidating furface adds to the quantity of el. and increases fluid being in some degree accumulated in the metallic plate, the oxidation;

which oxidates the metal, and it is by this means conveyed across the water to the filver plate, when two metals are used; or, in other cases, simply to the opposite surface of the oxidating substance. The electric fluid then enters the filver plate, and instantly passing on to the contiguous zinc plate, arrives at a fecond oxidating furface. The fame feries of events which have been described is here repeated, except that the electric

and this effect will increase with the number of pairs.

is difengaged by the fecond oxidating furface in larger quantity, and in a more concentrated flate, than before. By purfuing the fame train of operations, it is eafy to fee how the electric matter will continue to be accumulated in each fuccef-

five pair of plates, until, by fufficient repetition, it may be made to exist in the zinc end of the pile in any assigned degree of force.

Having

Having thus exhibited a general view of the hypothesis Remarks and which has suggested itself to my mind, as affording a simple inferences from this theory. explanation of the phenomena, without offering violence to any of the generally received opinions respecting chemistry or electricity, I shall proceed to illustrate its truth, by applying it to the explanation of some of the most remarkable properties of the pile. It is fufficiently obvious that upon these principles the action of the pile will continue as long as the furface of the zinc continues to be oxidated, and that, the fize of the pile remaining the fame, its energy will be proportionate to the rapidity of this operation. The real difference between the two ends of the apparatus will appear to be not fo much a difference in the absolute quantity of electrical matter contained in them, as a difference in the direction of its current; because exactly the same quantity of fluid which passes out of the zinc end must be received at the filver end, when a communication is established between them. It may be farther remarked, that when the pile is constructed as directed by Volta, beginning with zinc, then filver, card and zinc, &c. though the zinc forms the basis of the pile, yet as no change takes place in the electric fluid during its passage from the filver to the zinc, this might with equal propriety have been called the filver end: in order, however, to avoid confusion, the usual language has been adopted in this paper; the terms discharging and receiving ends would be more appropriate.

It has excited a good deal of surprize, that, in the experi-Considerations ments of Mr. Davy and Mr. Cruickshank, apparatuses, con-tuses of Davy structed of various shapes and of such very different materials, and Cruickshould be capable of producing the same operations. We may shank, in a few words enumerate the circumstances which are effential to its formation. 1st. An oxidable substance, one end or fide of which only is to be oxidated; 2d, A fubstance from which the electric fluid generated during the oxidation can abstract hidrogen: these two substances must be alternately arranged, either perpendicularly or horizontally; the first fets the fluid at liberty, the fecond confines it and conducts it in the proper direction. When we admit the necessity of the oxidation of the zinc, we immediately fee the reason why the pile ceases to act when perfectly pure water is interposed between the plates, why the action is suspended in vacuo and increased by oxigen gas. The hypothesis advanced above shews us how

and Davy's extrication of the gases in separate veffels.

an apparatus may be conftructed of one metal and charcoal, of one metal alone, or even of charcoal alone, provided a proper fluid be interpoled to each furface. The experiment related by Mr. Davy, where, upon uniting the ends of the pile to two glasses of water by means of muscular fibres, while the glasses themselves were connected by a filver wire, the effects of the ends of the wire were reverfed, admits of a fimple folution. The fluid must have been conducted along the fibres folely in confequence of the moisture attached to them; when it left the plus end of the pile it would therefore decompose part of this water, and passalong in union with hidrogen to the filver wire, where the hidrogen would be difengaged. The other end of the filver will then become oxidated, and the hidrogen absorbed will be evolved at the minus end of the pile. I have protracted this communication to fo unufual a length,

Fourcroy and Thenard's expe-that I shall only add one additional remark with respect to the riment that the power of giving shocks and burning wire depend on the mode metals. of building the

exhibiting the coated jars.

fact first ascertained by Fourcroy, that the power of the pile in giving shocks is not proportionate to its power in consuming A few alternations of large plates are more powerful in effecting the latter operation, while the same quantity of metal employed in conftructing a higher pile of less diameter, referred to elec- affords a more powerful shock. It appears from every cirtrical phenomena cumstance, that the force of the shock depends almost entirely fame refult with upon the concentration of the electric fluid; the fame quantity which, when contained in a few jars highly charged, would be fufficient to deftroy animal life, if dispersed over 100 jars would be fearcely perceptible. On the other hand, in the combustion of metals the absolute quantity of electricity is of great importance; provided the current be rapid (which, from the experiments of Dr. Van Marum, appears to be the case with the pile of Volta in an amazing degree), and if the quantity of fluid be confiderable, its flate of intenfity is of less moment \*. The intelligent reader will at once perceive in what way the fact obferved by Fourcroy admits of explanation from this train of reafoning.

> \* Cuthbertson first observed in a letter to me (1793) the striking fact, that the combustion of equal lengths of the same wire is made by equal quantities of electricity, whether the intensity be greater or less (within certain extended limits). See Account of Experiments. Philof. Journal, quarto, II. 215. August, 1798.-N.

#### III.

On the Supposed Existence of Mechanical Deposits and Petrefactions in the Primitive Mountains, and an Account of Petrefactions which have been discovered in the newest Flotz Trapp formation. By Mr. ROBERT [AMESON. Communicated by the Author.

JUR globe, according to the Wernerian geognofia, even The primitive during the deposition of the newer primitive strata, appears to rocks precipitated from a have been covered to a great height with water, as is evinced great depth of by the want of all mechanical deposit \*. After the precipi-water which covered the globe. tation of these great rock formations, the level of the water became so low, as to allow it to act mechanically upon the subjacent rocks; this occasioned the first mechanical deposition, The transition which discovers itself in the transition rocks (Ubergangsge-rocks when the burge). Nearly at the fame time organization commenced, at which time as it is in the transition rocks we find the first traces of organic organization beremains: these are generally zoophytes and sea plants, a fact gan. which goes deep not only into geology, but natural history.

Professor Playsair, in his illustrations of the Huttonian the-Instances of mein primaryfrata:

ory, mentions feveral inftances of mechanical depofits and chanical depofits petrefactions which have been discovered among the primary offered by Prostrata, and from these he concludes, that no such series of strata fessor Playsair. as the transition exist. I shall now examine the statements he has given; and first respecting the occurrence of petrefactions

in primary mountains.

At page 164 he observes, "Another spot, affording in-Shells in primiftances of shells in primitive limestone, is in Devonshire, on tive limestone, the sea shore, on the east side of Plymouth dock, opposite to of Plymouth, Stonehouse, I found a specimen of shistose micaceous lime-&c. stone, containing a shell of the bivalve kind; it was struck off from the folid rock, and cannot possibly be considered as an adventitious fossil. Now, no rocks can be more decidedly primary than those about Plymouth; they confist of calcareous strata, in the form either of marble or micaceous limestone, alternating with shiftus of the same kind, which prevails through Cornwall to the west, and extends eastward into Dartmoor, and on the sea coast as far as Berry Head. These all inter-

\* The exception to this in the sienite formation I explained in a former paper.

But it is not proved that those strata are primitive.

fect the horizontal plane in a line from east to west nearly a they are very erect, those at Plymouth being elevated to the north." That petrefactions exist in the limestone at Plymouth is evident; but that these strata are primitive, still remain to be proved. The character given of the limestone does not exclude it from the transition strata; but of the shiftus we cannot judge, as neither its oryclognoffical, or geognoffical characters are given. The other inftances which are alluded to, are liable to the same objection. I cannot therefore agree with Professor Playsair in believing "Though, therefore, the reconclusion of the mains of marine animals are not frequent among the primary rocks, they are not excluded from them; and hence the exiftence of shell sish and zoophytes, is clearly proved to be anterior to the formation even of those parts of the present land which are justly accounted the most ancient."

Whence the Profesior is not agreed to.

The polition that vegetable

matters are

tound in pri-

mitive ftrata

Professor Playfair agrees with Dr. Hutton in assirming, that vegetable matters occur in the primitive ftrata: I do not find it necessary to enter into an examination of what they have faid upon this fubject, as they have evidently confounded a need not be exgeognofical with an oryctognofical investigation.

amined. Prof. Playfair's proofs of mechanical depofits in primitive ftrata examined.

I shall now examine the proofs which Professor Playfair has brought to establish the existence of mechanical deposits in primitive strata.

Mount St. Gothard a central primitive mountain, has arenaceous strata in its vicinity.

The first we meet with is from Saussure. Professor Playfair remarks, "St. Gothard is a central point, in one of the greateft tracts of primary mountains on the face of the earth, yet arenaceous firata are found in its vicinity. Between Ayrolo and the Hospice of St. Gothard, Saussure found a rock, composed of an arenaceous or granular paste, including in it hornblende and garnets. He is fomewhat unwilling to give the name gres to this ftone, which Mr. Besson has done; but he nevertheless describes it as having a granulated structure."

Observation. The rock here mentioned was encifs or mica flate.

The rock of Ayrolo is primitive, and is either gneis or mica flate. Garnets are feldom found in gneifs, but are characteriftic for mica flate; the geognoti, therefore, would not hefitate to confider the rock here mentioned as belonging to mica flate.

Instances by P. Playfair of indurated arenacous granite.

Professor Playsair continues, "Among the most indurated rocks that compose the mountains of this island, many are arenaceous. Thus, on the western coast of Scotland, the great body of high and rugged mountains on the shores of Arasaig,

&c. from Ardnamurchan to Glenelg, confifts, in a great mea" fure, of a granitic fandstone, in vertical beds. This stone fometimes occupies great tracts; at other times, it is alternated with the micaceous, and other varieties of primary shiftus; it occurs, likewife, in feveral of the islands, and is a fossil which we hardly find described or named by writers on mineralogy."

The granitic fandstone of Glen-elg is most certainly gneifs \*, Obs. This graand a variety which is not uncommon; and I may venture to nitic fandstone is fay, that the strata in Arasaig, &c. are of the same nature.

Professor Playfair concludes with stating the following exam-Other examples ples, as a further confirmation of his opinion—" Much also of of Pr. Playfair. Granulated a highly indurated, but granulated quartz, is found in feveral quartz in beds places in Scotland, in beds of strata, alternated with the com-alternating with non shiftus of the mountain. Remarkable instances of this may be feen on the north fide of the ferry of Balachulish, and again on the fea shore at Cullen. At the latter, the strata are remarkably regular, alternating with different species of shiftus. At the former, the quartz is fo pure, that the frone has been mistaken for marble.

"These examples are perhaps sufficient; but I must add, that in the micaceous and talcofe schissi themselves, thin layers of fand are often found interpoled between the layers of mica or talc. I have feen a specimen from the summit of one of the highest of the Grampian mountains, where the thin plates, of a talcy or afbeftine fubftance, are feparated by layers of a very fine quartzy fand, not much confolidated. The mountain from which it was brought, confifts of vertical firata, much interfected by quartz veins. It is impossible to doubt, Whence he in this instance, that the thin plates of the one substance, and concludes that they were depo-the small grains of the other, were deposited together at the sited (inechanibottom of the fea, and that they were alike produced from cally) at the the degradation of rocks more ancient than any which now fame time. exift."

I am furprized Professor Playsair should adduce granular Obs. Granular quartz as a proof of mechanical deposition, as it has no more quartz is not a claim to fuch a character than granite. The following obser-a chemical depovations will render this evident; granular quartz differs from fition. mica flate, in the absence of the flaty fracture, and mica; we from mica flate

\* Mineralogy of the Scottish isles, vol. ii. p. 160.

mechanical, but For this differs in the absence of flaty fracture; have, but the feries

between each are gradual. Mica flate differs from gneifs in but here also the transition is And gneifs differs from granite ture, which it as gradually lofes. The gradation being perfect, the one is no more a mechanical deposit than the other. Loofe texture is no proof of mechanical deposition; and layers of fand are found in veins which in the volcanic theory are faid to bear every mark of complete fusion. General observations.

have, however, a feries from the most complete granular quartz to the most perfect mica slate. Again, mica slate, which differs from gneis in wanting selspar, is to be observed in all the intermediate stages until it passes into complete gneis; the gneis, which is principally distinguished from granite by its slaty fracture, gradually loses this fracture, and at length is not to be distinguished from granite. Thus we have a complete gradually loses.

It therefore follows, that if granular quartz is a mechanical deposit, so is granite, a position which I believe the Huttonian system would not allow.

That the granular quartz should occur sometimes of a very loose texture; nay, even as Prosessor Playsair remarks sandy, is not surprizing, for in granite and basalt we have similar appearances. Even in veins, which to use Prosessor Playsair's own words, bear all the marks of complete sustion, layers of granular quartz, from the most compact to the looseness of sand have been observed. The great sand veins in the Hartz afford remarkable instances of this; also, as I have more lately discovered, the lead veins at Wanlock-head in Lancashire.

Account of Organic remains which have been discovered in the newest Flötz Trapp formation.

Many different accounts had been given of the geognoftic relations of the rocks of this formation, before they engaged the attention of Werner, the great founder of true geognofia. After having made the remarkable discovery upon the hill of Scheibenberg, he extended his inquiries to all the basalt hills in Germany, and found in every quarter corresponding appearances. This confirmed more completely the conclusions he had then drawn, and intirely overthrew the volcanic system. He however did not stop here; his after observations disclosed a connection among these appearances, which at first he was not probably aware of; they placed the Neptunian system beyond the reach of attack, and completely annihilated a host of hypotheses. He proved,

Refults. 1. That this is the neweft of the great rocks. 2. That all the now unconnec-

Refults. 1. That 1. That this is the newest of all the great rock formations, this is the new- of which the crust of the earth is composed.

2. That all the apparently unconnected hills and maffes of this formation, have formerly flood in connection with each other.

3. That

3. That it exists in all quarters of the globe.

These facts lead him to the great discovery, that this for- this formation mation at one time, extended as a cover around the whole nected. earth. From these observations it is evident, that we may 3. That it exexpect to meet with vestiges of many of the organic and un- 4. That it fororganic matter which at that time existed upon the crust of the merly covered earth, in the rocks of this formation. Accordingly, the in- and must conveffigations of geognosts have discovered organic remains in tain organic reevery rock of this formation; the following are given as in-mains. flances:

ted maffes of ifts every where.

Organic remains in Greenstone, Basalt, Wacke, and Trapp Buccin.

#### QUADRUPEDS.

1. Werner in his geognostic lectures, informs us that the Organic remains wacken of Kalten-nordheim is sometimes sound to contain in greenstone, basalt, &c. deers horns.

found; viz. of

2. The Abbe Fortis discovered the head of an unknown quadrupeds. animal in a foft wacken, in the valley of Ronca in the Veronese. Vide Beschrebbung des Thales Ronca, s. 96.

3. Sauffure observed bones of quadrupeds in the wacken of the catacombs of Rome.

Lettre a Mr. le Chev. Hamilton, J. de Physique. Tom. VII.

#### SHELLS AND ZOOPHYTES.

1. Dr. Richardson found shells in rocks of the Trapp for- Of shells and mation at Ballycastle in Ireland.—Kirwan's Geol. Essays.

2. Mr. Von Buch informs us, that in the county of Landeck, he observed a bed of wacken, which contained besides pebbles of chalcedony, turbinites in a state of complete prefervation. Versuch einer mineralogisten Veschreibung von Landeck, von Leopold von Buch. f. 35.

3. Abbé Fortis observed numerous petrefactions of shells in Wacken, and Trapp Breccia, also a few in Basalt, in the formation of the valley of Ronca.

4. Berolding found a cornu ammonis, which still retained its mother of pearl lustre, in the basalt of Torez. In the bafalt of Thurgau, near the Boden lake, he observed gryphites, ammonites, and glossopetræ, Chem. Annal. 1794, p. 103.

5. In the Wernerian collection of petrefactions I faw specimens of greenstone, containing petrefactions of shells.

Vol. III .- SEPTEMBER, 1802.

Of vegetables.

#### VEGETABLE REMAINS.

- 1. Werner observed great trees, with branches, leaves, and fruit, in the wacken, at Joachimsthal.
- 2. Friesleben describes the impression of a plant in the Kawsower Berg, near to Podsedlitz.
- 3. In the islands of Banna and Skye I observed pieces of wood in Trapp Breccia, and Mineralogy of the Scottish isles, vol. ii. p. 58—75.

Organic remains in flaty clay, limestone, and sandstone, of this formation.

Organic remains found in flaty clay, limestone, and fandstone, which belongs to the Flötz Trapp formation.

#### SHELLS.

Shells.

- 1. Abbé Fortis, in his account of the valley of Ronca, informs us, that limestone and slaty clay often alternate with basalt, wacken, and trapp breccia. The limestone and slaty clay contains numerous petrefactions of shells, which are of the same kinds with those found in the basalt, wacken, and trapp breccia.
- 2. In the island of Eigg, where there appears to be a fimilar formation with that of Ronca, I observed that the limestone, slaty clay, and fandstone, contained numerous petrefactions of shells. Mineralogy of the Scottish isles, vol. ii. P.

#### Of vegetables.

#### VEGETABLE REMAINS.

- 1. At the northern extremity of the island of Skye, where basalt alternates with limestone and slaty clay, I observed pieces of carbonated wood in the limestone. Mineralogy of the Scottish isles, vol. ii. p. 80 \*.
- 2. In the flinty fandstone, which usually accompanies this formation, I have observed branches of shrubs; vegetable matters that occur inveloped in rocks are generally, either carbonated or bituminated, here however they are not altered.

R. JAMESON,

Sheriff Bra, Leith.

\* Mr. Kirwan, in the first volume of his System of Mineralogy, has given us an excellent account of the different opinions respecting the formation of basalt; and has adduced many arguments that shew the fallacy of the Volcanic and Plutonic hypothesis.

POSTSCRIPT.

#### POSTSCRIPT.

Since writing the inclosed, I have examined one of those appearances, which are confidered by Dr. Hutton and Professor Playsair, as demonstrating the existence of petrefactions in primitive mountains. Dr. Hutton, at p. 334 of his Theory of the Earth, remarks, " I have already observed, that one Dr. Hutton's fingle example of a shell, or of its print, in a schissus, or in a fastement of sacts to shew the ftone stratified among those vertical or erected masses, suffices mechanical orito prove the origin of these bodies to have been, what I had gin of primitive maintained them to be, water formed firata created from the the existence of bottom of the fea, like every other confolidated stratum of the organic remains. earth. But now, I think, I may affirm that there is not, or rarely, any confiderable extent of country of the primary kind, in which fome mark of this origin will not be found, upon careful examination; and now I will give my reason for this I have been examining the fouth alpine country of Scotland occasionally, for forty years back, and I could not find any mark of an organized body in the schistus of those mountains. It is true, that I knew of only one place where limestone is found among the strata: this is upon Tweedside near the Crook. This quarry I had carefully examined long ago, but could find no mark of any organized body in it. suppose they are now working some other of the vertical strata near to those which I had examined: for, in the summer of 1792, I received a letter from Sir James Hall, which I shall now transcribe. It is dated Mosfat, June 2, 1792.

"As I was riding yesterday between Noble House and the Sir James Hall's Crook, on the road to this place, I sell in with a quarry of account of orgaalpine limestone; it consists of sour or sive strata, about three limestone stated feet thick, one of them single, and the rest contiguous; they to be primitive. all stand between the strata of slate and schist, that are at that place nearly vertical. In the neighbourhood, a slate quarry is worked of pure blue slate; several of the strata of slate near the limestone, are silled with fragments of limestone scattered about like the fragments of schist in the sandstone, in the neighbourhood of the junction on our coast. Among the masses of limestone lately broken off for use, and having the fracture fresh, I sound the forms of cockles quite distinct, and in great abundance. I send you three pieces of this kind,"

It

It may perhaps be alledged, that those mountains of Cumberland and Tweedale are not the primary mountains, but composed of the secondary schistus, which is every where known to contain these objects belonging to a former earth. Naturalists who have not an opportunity of convincing themfelves by their proper examination, must judge with regard to that geological fact by the description of others. Now it is most fortunate for natural history, that it has been in this range of mountains that we have discovered those marks of a marine origin; for, I shall afterwards have occasion to give the cleareft light into this subject, from observations made in other parts of those fame mountains of schift, by which it will be proved that they are primary strata; and thus no manner of doubt will remain in the minds of naturalists, who might otherwise sufpect that we were deceiving ourselves, by mistaking the secondary for the primitive schistus. Dr. Hutton's account of the mountains in the fouth of Scot-

Remark that the limestone is not primitive,

land is confused and unscientifical, and hardly comes within the pale of true geognofia. It is not my intention, at present, to enter into an examination of his observations; the object of this postscript is to shew, that the limestone between Noble House and the Crook Inn does not belong to the primitive The beds of limestone mentioned by Sir James mountains. Hall, I observed lying between strata of transition slate, and alternating with this flate alternating with strata of grey wacke; consequently the whole belongs to the transition class of rocks.

transition slate; strata of grey wacke. Description of

for it lies between strata of

The limestone has a blueish grey colour, fracture is foliated, the feveral rocks, the distinct concretions are from coarse to fine grained, and it is hardly transfucid on the edges. It is often traverfed by veins of calcareous spar, and fometimes it contains thin beds of flinty flate (Kiesel Schiefer of Werner). The transition flate has a blueish or fmoke grey colour, has generally lefs lustre than the primitive slate, and contains much interspersed mica. I observed it in all the stages from nearly pure slate to grey wacke. The grey wacke is composed of fragments of transition flate, flinty flate, and quartz, connected by a basis of transition slate. It is frequently traversed by veins of quartz, and is to be observed where the fragments are hardly distinguishable from their fize; it has much the appearance of a breccia. I shall take another opportunity of fending you drawings of the different kinds of petrefactions that occur in the limeftone.

#### IV.

A Method of increasing the Sensibility of the Barometer, ad libitum. By the Rev. JAMES WILSON, A. M. Communicated by the Author.

HOUGH there have been many contrivances for making Contrivances to barometers with indefinite scales, they all seem to labour under enlarge the scale of the basome difficulty either in their construction or use.

The evaporation of water in that of Des Cartes, the fluggish by water; by motion of the mercury in the horizontal and diagonal ones, the clined columns, friction in Dr. Hooke's, and the unfteadiness of floating bodies by mechanism in Rownings, have rendered these ingenious contrivances have not proved scarce preferable to the common vertical one.

horizontal or in-

None of these defects. I believe, will attach to that which I Description of a propose. It is as follows: (See Fig. 1. Plate II.) A B, a tube new barometer. differing in no respect from that of the common barometer, floats in the but that it is wider and longer, so that a cylindrical rod, qr, mercury of a fyphon baromemay float freely on the mercury, and that the lower end is ter. connected to a small bent tube BCD, instead of being immerfed in a ciftern. To the lower end of the floating rod is fastened a hair, or fine iron wire, (what I used was a piece of the Indian weed used by anglers) which is brought through the mercury, and out at D, fo that the rod may be either drawn down, or fuffered to ascend at pleasure.

Upon the tube CD is a mark N, to which by moving the rod we can at any time bring the surface of the mercury. For by drawing it down we force up mercury into each tube, and on the other hand, by fuffering it to afcend, we fuffer the mercury to descend in each.

The furface of the mercury then being brought to this mark, and is drawn and the rod being fastened, by tying the hair to the pin E, if down into the mercury till the on inspecting it afterwards, we find that the surface of the internal rise mercury has moved from N, we are fure there has been fome causes the lower furface to stand variation in the weight of the atmosphere, viz. if it has fallen, at a fixed mark. we know that this weight has increased, and vice versa.

The variations either at M or N are in a given ratio to the The variations variations in the length M N of the fustained column. For at the inner or outer fursace are the rife or fall at M, is to the fimultaneous fall or rife at N, in proportioned to the inverse ratio of the bases, or horizontal sections of the those of the whole column.

mercury at M and N; namely, as the area of the horizonta fection of the tube D C is to the difference of the areas of the horizontal fections of the tube A B, and of the rod. That is, if D, d, and r be the diameter of the tubes A B, D C, and rod, as  $d^2: \overline{D^2-r^2}$ , which is a given ratio, and confequently each quantity has to the fum of both, which is the variation in the length M N a given ratio, e. gr.  $d^2$  (or  $D^2-r^2$ ):  $D^2+d^2-r^2$ .

The variations either at M or N would therefore, if we could measure them with sufficient accuracy, express truly the variation in the weight of the atmosphere.

and so are those in the rife or fall of the rod; if it be in every case adjusted to the mark.

And this meafure may be enlarged at will.

Proof.

But if instead of attempting this, we move the rod by means of the hair until the mercury return to N, by the length of rod moved either into the mercury, or out of it, we have another measure, and one as long as we please, of the variations in the weight of the atmosphere.

For, supposing the mercury to have fallen at N, any space Nx, and risen at M, a space My, for the adjustment we must draw down the rod until it has displaced as much mercury as will fill the space Nx, and also a space yx in the other tube of precisely the same bulk, through which it must rise at the same time. The length of rod drawn down for each of these purposes will have a given ratio to the space Nx, and consequently their sum added to My will have a given ratio to it: and thence, as Nx is a measure of the variation of the atmosphere, this portion of rod will also be a measure: and manifestly it may be increased at pleasure by lessening the thickness of the rod.

This appears as follows. The proportion of the rife in one tube to the fimultaneous fall in the other upon any change of the weight of the atmosphere, was above determined to be as  $\frac{1}{D^2 - r^2} : \frac{1}{d^2}$ . Let q be the quantity of mercury transferred from one tube to the other, the heights My and Nx will be  $= \frac{1}{D^2 - r^2}$  and  $\frac{q}{d^2}$  and the sum of these is the variation which

The variation  $My = \frac{q}{D^2 - r^2}$  is feen on the rod without moving it; upon moving it there is added to this, first,  $\frac{q}{r^2}$ 

would take place in De Luc's barometer.

to occasion the ascent N x (the bulk of rod drawn down for that purpose being = to the bulk of q, and thence its altitude  $=\frac{q}{r^2}$ ) and secondly,  $\frac{q D^2 - q r^2}{d^2 r^2}$  to occasion the equal simultaneous ascent yx (since the bulk of rod drawn for that purpose is greater than the bulk q in the ratio  $D^2 - r^2$ :  $d^2$ , and therefore equal to  $\frac{q D^2 - q r^2}{d^2}$  and its altitude equal to  $\frac{q D^2 - q r^2}{d^2 r^2}$ ).

The fum of these three  $\frac{q}{D^2 r^2} + \frac{q}{r^2} + \frac{qD^2 - qr^2}{d^2 r^2}$  is the

whole variation feen on the rod. This is to  $\frac{q}{D^2-r^2} + \frac{q}{d^2}$ , the variation on De Luc's fcale, (by bringing them to a common denominator) as  $D^4-2D^2r^2+D^2d^2+r^4:D^2r^2+d^2r^2-r^4$ , which is a given ratio; and therefore the rod affords an accurate measure.

Let s and 1 be the rod scale, and the common scale  $s = \frac{D^4 - 2 D^2 r^2 + D^2 d^2 + r^4}{D^2 r^2 + d^2 r^2 - r^4}$ , which by performing the divi-

fion gives the infinite feries  $\frac{D^2}{r^2} - 1 + \frac{d^2}{D^2} - \frac{d^4}{D^4} + \&c.$   $\frac{D^2}{r^2}$  is the principal term of this quote; for the fum of the infi-

nite feries commencing with -1 is  $\frac{-D^2 + r^2}{D^2 + d^2 - r^2}$  which is necessarily less than a negative unit; and if d be supposed to vary, decreases still as it increases.

The scale therefore being increased in so trisling a degree by the increase of d, may be said to be as  $\frac{D^2}{r^2}$ .

The narrower the tube DC is I should suppose the better; since though the scale is diminished in a small degree, yet the variation Nx is rendered more sensible, which is a point of great importance.

Injury from agitation is thus also in some degree prevented.

Though we can thus from D, d, and r calculate the scale, The scale of this yet as we can scarcely measure them with sufficient accuracy, barometer best obtained from the observation.

the best way is to determine it by observation, viz. by finding how much of the rod expresses some larger variation in the common barometer, and then taking out the rod and graduating, &c. or rather by taking the mean of a great number of observations.

Should this be found on a fufficient trial to answer the purpose, I can point out a method of making it serve also as a thermometer.

This barometer

Before I conclude, I shall just mention the idea which led was suggested by me to this construction, which though the greatest increase of other construction fcale it gives is 14:1, may possibly in the end be found to answer better.

Water poured on the exposed fur. face of a syphon mercurial barometer gives a barometer which thew variations 14 times as long change.

It struck me that if we were to pour in water into a long tube DF, connected with the tube of the common barometer as in Fig. 2, or to take out water fo as at any time to bring the mercury to the fixed point N, the altitude of water either may be made to poured in or taken out would give us a measure for the variations of the barometer fourteen times larger than in the comas the mercurial mon one, fince water is fourteen times lighter than mercury. This is evident from the common hydrostatical principles, as the altitude of water poured in is to be a counterpoise to the fum of the fall of mercury in AB, and rife in CD, and e contra, the altitude of water taken out is to be a counterpoise to the fum of the rife of mercury in A B, and fall in C D.

If the tube DF therefore were graduated, it would give a fcale 42 inches long, and the rife of water would indicate diminution in the weight of the atmosphere and v. v. thence the scale to be inverted. To facilitate the pouring out and in of water, let there be a bucket H moveable up and down by rackwork, or otherwise, from D to F; to this let there be attached the fyphon S.

By raifing the bucket until the level of the water in it be higher than the level of that in the tube, we pour in water at pleasure, and e contra, by bringing down the bucket we take out water at pleafure.

This instrument will have all the advantages of a water barometer without its unmanageable length; nor do I see any objection to it, except perhaps that its appendages are too bulky \*.

V. A Summary

\* The principal objection to the ingenious contrivances in the present communication are, that the results originally depend on the

A Summary of the most useful Parts of Hydraulics, chiefly extracted and abridged from Eytelwein's Handbuch der Mechanik und der Hydraulik. Berlin, 1801. By THOMAS YOUNG, M. D. F. R. S. \*

THE theory of hydraulics has never been carried to a very State of hydrauhigh degree of perfection upon mathematical foundations alone; lie theory, and difficulties of apnor has it hitherto, even with the affiftance of experiment, plying it to pracbeen rendered of much practical utility. Newton began the tice. investigation of the motions of fluids: Daniel Bernoulli added to Newton's propositions much valuable matter, both from calculation and from experiment; D'Alembert and many later authors have exercifed their analytical talents in inquiries of a But another and a more practicable mode of fimilar nature. attaining hydraulic knowledge has been attempted by a distinct class of investigators, at the head of whom stands the Chevalier de Buat. These have begun from experiment alone, and have laboriously deduced from very ample observations of the actual refults of various particular cases, the general laws by which the phenomena appear to be regulated, or at least the formulas by which the effect of new combinations may be predicted. But it must be confessed that these formulas, however accurate. are too intricate to be retained in the memory, or to be very eafily applied to calculations from particular data.

Mr. Eytelwein, a gentleman already known to the public Mr. Eytelwein's by his translation of Buat's work into German, with important valuable comadditions of his own, and honoured with feveral employments and titles relative to the public architecture of the Pruffian dominions, has collected into this compendium of mechanics and hydraulics, the principal facts that have been afcertained, as well by his own experiments, as by those of former authors,

the accuracy of adjustment at N, in which the errors are precisely the fame as in the common barometer. The double barometer affords an instance of a lighter sluid floating on a larger; but with a somewhat shorter scale than that of fig. 2 .- N.

\* This excellent memoir is copied from the Journals of the Royal Institution.

especially

especially such as are the most capable of practical application; and he appears to have done this in so judicious a manner, as to make his book a most valuable abstract of every thing that can be deduced from theory respecting natural and artificial hydraulics. This elegant conciseness deserves so much the more praise, as his countrymen in general appear too often to make a merit of prolixity; and we shall have occasion to remark, that besides the convenience of simplicity, he has sometimes been fortunate enough to unite with it the advantage of superior accuracy.

Part 1. Mechanics. The first part of the work is but short; it relates to proper mechanics, and has little that is remarkably new or interesting. In treating of pendulums, the author informs us, with reference to another work of his own, that the Rhinland or Brandenburg foot contains 139.13 French lines. Hence it appears that 100 Rhinland feet are exactly 103 English; and in this paper, the measures will be reduced accordingly.

Part 2. Hy-draulics.

The fecond part, relative to hydraulics, contains besides a short introduction, twenty-four chapters, almost every one of which presents to us something of importance.

Water issuing from refervoirs.

Chapter 1. Of the motion of water flowing out of refervoirs, and of the contraction of the stream.

Law of its velocity: As the square root of the height. § 89. The velocity of water flowing out of a horizontal aperture, is as the square root of the height of the head of water.

That is, the pressure, and consequently the height, is as the square of the velocity: for the quantity slowing out in any short time, is as the velocity; and the force required to produce a velocity in a certain quantity of matter in a given time, is also as that velocity; therefore the force must be as the square of the velocity. The proposition is fully confirmed by Bossur's experiments; the proportional velocities, with a pressure of 1, 4, and 9 feet, being 2722, 5436, and 8135, instead of 2722, 5444, and 8166; a very inconsiderable difference.

Another developement. There is another mode of confidering this proposition, not mentioned by Eytelwein, which is a very good approximation. Supposing a very small cylindrical plate of water immediately over the orifice, to be put in motion at each instant by means of the pressure of the whole cylinder standing on it, and supposing all the gravitation of the column to be employed in generating the velocity of the small cylindrical plate, neglecting

its own motion, this plate would be urged by a force as much greater than its own weight as the column is higher than itself; and this, through a space shorter in the same proportion than the height of the column. But where the forces are inversely as the spaces described, the final velocities are equal. (Young's Syllabus, 35.) Therefore the velocity of the water flowing out must be equal to that of a heavy body falling from the height of the head of water, which is found very nearly by multiplying the square root of that height in seet by 8, for the number of feet described in a second. Thus a head of 1 foot gives 8; a head of 9 feet, 24.

The well-known circumstance of the contraction of the The contraction stream or vein of water running out of a simple orifice in a thin of the stream. plate, reduces the area of its section at the distance of about half its diameter from the orifice, from 1 to .660 or .666 according to Bossut, to .631 according to Venturi, and to .64 or \frac{16}{25} according to the author's own experiments: hence the diameter is reduced to \frac{4}{25}.

The quantity of water discharged is very nearly, but not Practical rule for quite, sufficient to fill this section with the velocity due, or cor- a simple orisice. Multiply the responding to the height: for finding more accurately the square root of quantity discharged, the orisice must be supposed to be dimitive the height by 5 missed to .619, or nearly \frac{5}{8}. Hence we may multiply the square city in 1 second. root of the height by 5 instead of 8, for this mean velocity in a simple orisice.

If we apply the shortest pipe that will cause the stream to Additional pipe adhere every where to its sides, which will require its length to the diameter of be twice its diameter, the discharge will be about  $\frac{13}{16}$  of the full the hole requires quantity, and the velocity may be found by taking  $6\frac{1}{2}$  for a  $6\frac{1}{2}$  multiplier of multiplier.

The greatest diminution is produced by inserting a pipe so Disadvantage of as to project within the reservoir, probably because of the greater an interior pipe; interference of the motions of the particles approaching its orifice in all directions: in this case the discharge is reduced nearly to a half.

A conical tube, approaching to the figure of the contraction conical, &c. of the stream, procured a discharge af .92; and when its edges produced a discharge of o.98, were rounded off, of .98, calculating on its least section.

Venturi \* has afferted that the discharge of a cylindrical pipe Venturi's coni-

\* Venturi's treatife is given entire in our Journal, 4to, Vol. II. and III.

may be increased by the addition of a conical tube nearly in the ratio of 5 to 2; but Mr. Eytelwein finds this affertion somewhat too strong, and observes, that when the pipe is already very long, scarcely any effect is produced by the addition of fuch a tube. He proceeds to describe a number of experiments made with different pipes, where the standard of comparison is the time of filling a given vessel out of a large refervoir, which was not kept always full, as it was difficult to avoid agitation in replenishing it, and this circumstance was perfectly increases the dif-indifferent to the results of the experiments. They confirm the

charge from a fimple aperture

affertion that a compound conical pipe may increase the diftwice and a half. charge to twice and a half as much as through a simple orifice, or to more than half as much more as would fill the whole fection with the velocity due to the height: but where a confiderable length of pipe intervenes, the additional orifice appears to have little or no effect.

> The chapter concludes with a general table of the coefficients for finding the mean velocity of the water discharged by the pressure of a given head under different circumstances.

Multipliers for velocity under different circumstances.

For the whole velocity due to the height, the coefficient, by determining the which its square root is to be multiplied, is 8.0458.

For an orifice of the form of the contracted stream, 7.8.

For wide openings, of which the bottom is on a level with that of the refervoir; for fluices with walls in a line with the orifice; for bridges with pointed piers, 7.7.

For narrow openings, of which the bottom is on a level with that of the refervoir, for fmaller openings in a fluice with fide walls, for abrupt projections and square piers of bridges, 6.9.

For short pipes from two to four times as long as their diameter, 6.6.

For openings in fluices without fide walls, 5.1.

For orifices in a thin plate, 5.

Chapter 2. Of the discharge of water by horizontal and by fmall lateral orifices, in a veffel continuing full.

Particular cases.

The principles detailed in the first chapter are here applied to particular cases.

Discharge in the dam,

Chapter 3. Of the discharge by rectangular orifices in the through a notch fide of a refervoir, extending to the furface.

> The velocity varying nearly as the square root of the height, may here be represented by the ordinates of a parabola, and the

quantity

quantity of water discharged by the area of the parabola, or two thirds of that of the circumscribing rectangle. So that the has nearly two-quantity of water discharged may be found by taking two-locity due to the thirds of the velocity due to the mean height, and allowing for mean height. the contraction according to the form of the opening, as explained in the first chapter.

The author has found this mode of calculation sufficiently near to the results of Buat's experiments, and to some accurate observations of his own.

He proposes for example a lake in which a rectangular Particular exemopening is made without any oblique lateral walls, 3 feet wide, plification. and extending 2 feet below the surface of the water. Here the coefficient of the velocity, corrected for contraction, is 5.1, and the corrected mean velocity  $\frac{2}{3}\sqrt{2\times5.1}=4.8$ ; therefore the area being 6, the discharge of water in a second is 28.8 cubic feet, or nearly four hogsheads.

The fame coefficient ferves for determining the discharge over a were of considerable breadth; and hence it is easy to deduce the depth or breadth requisite for the discharge of a given quantity of water. For example, a lake has a were 3 feet in breadth, and the surface of the water stands at the height of 5 feet above it: it is required how much the were must be widened in order that the water may be a foot lower. Here the velocity is  $\frac{2}{3}\sqrt{5}\times5.1$ , and the quantity of water  $\frac{2}{3}\sqrt{5}\times5.1\times3\times5$ ; but the velocity must be reduced to  $\frac{2}{3}\sqrt{4}\times5.1$ , and then the section will be  $\frac{2}{3}\sqrt{4}\times5.1$ 

 $\frac{\sqrt{5 \times 3 \times 5}}{\sqrt{4}}$ =7.5. $\sqrt{5}$ ; and the height being 4, the breadth must be  $\frac{7.5}{4}\sqrt{5}$ =4.19 feet.

Chapter 4. Of the discharge from reservoirs with lateral By lateral oriorifices of considerable magnitude, with a constant head of water. able magnitude,

This may be found by determining the difference in the difcharge by two open orifices of different heights: but in most cases the problem may be solved with nearly equal accuracy, the velocity due by considering the velocity due to the distance of the centre of to the center of gravity of the orifice below the surface.

On the control of the centre of the centre of the center of the center

Chapter 5. Of the discharge from reservoirs receiving no taken. supply of water.

Refervoirs not supplied. For prismatic vessels, all the particulars of the discharge may be calculated from the general law that twice as much would be discharged from the same orifice if the vessel were kept full during the time which is required for its emptying itself. (Young's Syllabus, 245). Where the form is less simple, the calculations become intricate, and are of little importance.

Divided refer-

Chapter 6. Of the discharge from compound or divided reservoirs.

The author observes from Buat, that the discharge through an orifice between two refervoirs, below the furface, is the fame as if the water ran into the open air. Hence he calculates the discharge when the water has to pass through several orifices in the fides of as many refervoirs open above. In fuch cases, where the orifices are small, the velocity in each may be confidered as generated by the difference of the heights in the two contiguous refervoirs, and the square root of the difference will therefore reprefent the velocity; which must be in the feveral orifices, inverfely as their respective areas; so that we may calculate from hence the heights in the different refervoirs when the orifices are given. Mr. Eytelwein then confiders the case of a lock which is filled from a canal of an invariable height, and determines the time required, by comparing it with that of a vessel emptying itself by the pressure of the water that it contains, observing that the motion is retarded in both cases in a similar manner; and he finds the calculation agree fufficiently well with experiments made on a large scale. The motion of water through different compartments of a closed cavity is also determined.

A lock filled from a canal.

Velocity of a

Chapter 7. Of the motion of water in rivers.

The fimple theorem by which the velocity of a river is determined, appears to be the most valuable of M. Eytelwein's improvements; although the reasoning from which it is deduced is somewhat exceptionable. The friction is nearly as the square of the velocity; not because a number of particles proportional to the velocity are torn asunder in a time proportionally short, for according to the analogy of solid bodies, no more force is destroyed by friction when the motion is rapid than when slow, but because when a body is moving in lines of a given curvature, the destecting forces are as the squares of the velocities (Young's Syllabus, 40, 35, 46), and the particles of water in contact

contact with the fides and bottom must be destected, in confequence of the minute irregularities of the furfaces on which they flide, nearly into the same curvilinear path, whatever their velocity may be.

At any rate we may fafely fet out with the hypothesis, that Elementary positive principal part of the friction is as the square of the velocity. tion, and inductions.

And the friction is nearly the same at all depths: for Professor Robifon found that the time of the ofcillation of the fluid in a bent tube was not increased by increasing the pressure against the fides, being nearly the same when the principal part of the tube was fituated horizontally as when vertically. The friction will however vary according to the furface of the fluid which is in contact with the folid, in proportion to the whole quantity of fluid: that is, the friction for any given quantity of water will be as the furface of the bottom and fides of a river directly, and as the whole quantity of water in the river inverfely: or supposing the whole quantity of water to be spread on a horizontal furface, equal to the bottom and fides, the friction is inverfely as the height at which the river would then fland, which is called the hydraulic mean depth.

Now when a river flows with an uniform motion, and is neither accelerated nor retarded by the action of gravitation, it is obvious that the whole weight of the water must be employed in overcoming this friction; and if the inclination vary, the relative weight or the force that urges the particles along the inclined plane, will vary as the height of the plane when the length is given, or as the fall in a given distance (Young's Syllabus, 54); consequently the friction, which is equal to the relative weight, must vary as the fall, and the velocity which is as the square root of the friction, must be as the square root of the fall; and supposing the hydraulic mean depth to be increased or diminished, the inclination remaining the same, the friction would be diminished or increased in the same ratio; and therefore in order to preserve its equality with the relative weight, it must be proportionally increased or diminished by increasing the square of the velocity in the ratio of the hydraulic mean depth, or the velocity in the ratio of its square root.

We may therefore expect that the velocities will be conjointly The velocities

as the square root of the hydraulic mean depth and of the fall will be as a mean proportional bein a given distance, or as a mean proportional between these tween the hytwo lines. Taking two English miles for a given length, we draulic mean depth and the must fall;

or 11-10ths of the velocity per fecond. must find a mean proportional between the hydraulic mean depth and the fall in two miles, and inquire what relation this bears to the velocity in a particular case, and thence we may expect to determine it in any other. According to Mr. Eytelwein's formula, this mean proportional is  $\frac{11}{10}$  of the velocity in a second.

Confirmation by a practical inflance.

In order to examine the accuracy of this rule, we may take an example which could not have been known to Mr. Eytelwein. Mr. Watt observed, as Professor Robison informs us in the article River of the Encyclopædia Britannica, that in a canal 18 feet wide above, and 7 below, and 4 feet deep, having a fall of 4 inches in a mile, the velocity was 17 inches in a fecond at the surface, 14 in the middle, and 10 at the bottom: so that the mean velocity may be called 14 inches, or somewhat less, in a second. Now to find the hydraulic mean depth, we must divide the area of the section, 2. (18+7)=50 by the breadth of the bottom and length of the sloping sides added toge-

ther, whence we have  $\frac{50}{20.6}$ , or 29.13 inches: and the fall in two miles being 8 inches, we have  $\sqrt{(8 \times 29.13)} = 15.26$  for the mean proportional, of which  $\frac{10}{11}$  is 13.9; agreeing exactly with Mr. Watt's observation. Professor Robison has deduced from Buat's elaborate theorems 12.568 inches for the velocity, which is considerably less accurate.

Another in-

For another example we may take the Po, which falls 1 foot in two miles, where its mean depth is 29 feet; and its velocity is observed to be about 55 inches in a second. Our rule gives 58, which is perhaps as near as the degree of accuracy of the data will allow.

On the whole, we have ample reason to be satisfied with the unexpected coincidence of so simple a theorem with observation: and in order to find the velocity of a river from its sall, or the sall from its velocity, we have only to recollect that the velocity in a second is  $\frac{10}{11}$  of a mean proportional between the hydraulic mean depth and the sall in two English miles. This is however only true of a straight river slowing through an equable channel.

True only in a strait river, &c.

Slope of the banks.

For the flope of the banks of a river or canal, Mr. Eytelwein recommends that the breadth at the bottom should be  $\frac{2}{3}$  of the depth, and at the furface  $\frac{10}{3}$ : the banks will then be in general capable of retaining their form. The area of such a section

is twice the square of the depth, and the hydraulic mean depth 3 of the actual depth. He then investigates the discharge of a canal of which the bottom is horizontal. The velocity appears in this case to be somewhat greater than in a similar canal, of which the bottom is parallel to the furface.

The author remarks that the velocity is greater near the con- Velocities when cave than the convex fide of a flexure: a circumftance probably the channel is occasioned by the centrifugal force accumulating the water on that fide. No general rule can be given for the decrease of the velocity in going downwards: but fometimes the maximum appears to be a little below the furface. In the Arno the velo- at different cities are at 2 feet below the surface,  $39\frac{1}{2}$  inches; at 4,  $38\frac{1}{2}$ ; depths. at 8, 37; at 16, 331; at 17, 31. In the Rhine at 1 foot, 58 inches; at 5, 56; at 10, 52; at 15, 43. As an approximation to the mean velocity, the author directs us to deduct from the superficial velocity  $\frac{1}{130}$  for every foot of the whole depth. For instance; if the depth were 13 feet, and the superficial velocity 5 feet, to take  $4\frac{1}{2}$  as the average velocity of the whole river. This can however only be true in large rivers: for in the canal, measured by Mr. Watt, the superficial velocity must be diminished nearly \( \frac{1}{5} \) for a depth of only 4 feet. And we may in general come quite as near to the mean velocity by taking of the superficial velocity; although this may still differ materially from the true medium. But comparing this with the former theorem for the velocity, which gives a refult oftener above than below the truth, we may bring them both into a form eafily recollected, thus;

The superficial velocity of a river is nearly a mean propor- Concile deductional between the hydraulic mean depth and the fall in two tion of the hydraulic mean miles; and the mean velocity of the whole water is, still more depth. nearly, nine-tenths of this mean proportional.

We may find a double confirmation of these principles in Major Rennel's account of the Ganges (Phil. Tranf. 1781, p. 87).

He informs us that "at 500 miles from the fea, the channel Instance in the is 30 feet deep when the river is at its lowest; and it conti-Ganges. nues at least this depth to the sea," that " a section of the ground, parallel to one of its branches, in length 60 miles, was taken by order of Mr. Hastings;" that "the windings of the river were fo great as to reduce the declivity on which the water ran, to less than 4 inches per mile;" that "the medium Vol. III .- SEPTEMBER, 1802.

rate of motion of the Ganges is less than three miles an hour in the dry months;" that is, its superficial velocity. Now allowing a little for the banks or shelving sides, we may take exactly 30 feet as the hydraulic mean depth; then if the fall in two miles were precisely  $\frac{2}{3}$ , we should have  $\frac{2}{3} \times 30 = 20$ ; and  $\sqrt{20} = 4.47$  for the velocity in a second, or 3.05 miles in the hour; which is a little greater than the observed velocity, because the fall was assumed somewhat too great.

Again (p. 110), "the river when full, has thrice the volume of water in it, and its motion is also accelerated in the proportion of 5 to 3. We may assume, that the hydraulic mean depth is doubled at the time of the inundation, whence the velocity will be increased in the ratio of 7 to 5: but the inclination of the surface is probably somewhat increased at the same time, which may easily be supposed to increase the velocity still surther, from 1.4 to 1.7.

The effects of weres, falls, and contractions in rivers.

Chapter 8. Of the discharge and the swell in the case of weres, falls, and contractions, in rivers and canals.

The methods employed in the third chapter require here fome modification, fince the water arrives at the place of defcent with a confiderable velocity; and it is evident from mechanical, as well as from hydraulic confiderations, that the ultimate velocity will exceed that which is due to the depth of the stream at the place of its descent, and that it will correfpond to a height equal to the fum of the heights capable of producing these velocities. Hence we may calculate the effect of a bar in elevating the furface of a river; how broad a were must be, in order to produce a certain elevation, and how much water will run over a given were according to collateral circumstances. When a bar is below the level of the lower water. we must consider the difference of the two levels as constituting the fall; the whole of the stream below the level of the lower water deriving its additional velocity from this difference only.

The extent of the fwell produced by a given elevation of the furface of a river in confequence of the effect of a were or bar, may be determined by calculating from the rules for finding the velocity of rivers, the inclination necessary for producing a given discharge; the depth being greater, the inclination immediately above the bar will be less, but the effect of the swell does not terminate at the point where the new surface,

if straight, would have met the original surface; for on account of the rounding off of this angle, it extends nearly twice as far. The effect of reducing the breadth of a river may be determined in a manner nearly fimilar. The author remarks, that a confiderable diminution of breadth produces but a fmall elevation, a refult which appears to be conformable to experience; but that where depth is required for navigation, it may often be obtained by a projection built out from the bank, which may be sufficient to increase the river's velocity, and to cause it to excavate its bed.

(To be concluded.)

#### VI.

Experimental Proof, that Corrections deduced from the Arcs of Vibration of a Pendulum in Vacuo, are practically ufeful. Mr. EZEKIEL WALKER.

#### To Mr. NICHOLSON.

SIR.

Lynn, Aug. 20, 1802.

YOUR annotation on my last paper \* convinces me, that I Introduction. ought to have given you my reasons, derived either from theory or experience, for troubling you on the fubject which it contains.

Though theory may be too uncertain a guide to be depend- Arguments ed on, in a matter so complicated as the vibration of a pendu- might be offered in favour of aplum in a variable refisting medium, yet I think that some philo-plying the theory fophical arguments might be advanced in its favour; but these of pendulums to I shall not at present enter into, as I mean to confine myself to mere matters of experience.

The clock which I used formerly was made by the late Mr. but at present John Arnold, in his best manner. It had a dead beat, with a the author concompound pendulum that vibrated in the arc of a circle. This fines himself to pendulum frequently varied in its arc; and as the short vibra- fact. tions were performed in less time than the long ones, I was led to suppose that it might be governed by the same law, which

\* See Vol. III. p. 273.

Clock of Arnold's make with dead beat and compound pendulum, found to vary accord-

obtains in a pendulum vibrating in a non-refifting medium by the force of gravity: and, on making use of corrections deduced from that principle, I found them agree fo well with the going of the clock, as left me no reason to doubt of the truth of my ing to the theory, conjecture. The table which I fent you was therefore constantly used to correct the clock's rate, when any alteration was observed in the arc of vibration; but the rate itself was ascertained by transits of the sun and fixed stars over the meridian.

It is thought that the corrections would not be applicable in binations.

How far this table may be applicable to other clocks I know not, as this is the only instance in which I have had any expe-It is evident, however, that this theory cannot be fome other com- used to advantage where a pendulum is acted upon either by cycloidal cheeks, a faddle-piece, or pallats of a particular conffruction.

I am, SIR,

With much respect, Your humble fervant, E. WALKER.

#### VII.

Description of an Apparatus of Tubes for facilitating to driving of Copper Bolts into Ships. By Mr. RICHARD PHILLIPS \*.

Mr. Phillips's method of driving copper bolts,

MR. RICHARD PHILLIPS, of Bristol, in several letters fent to the Society, states, that he had invented a method of driving copper bolts into ships, without splitting the heads or bending them; and that by means of tubes contrived by him for the purpose, this could be effected without difficulty, and had been fatisfactorily executed in the presence of several of the principal ship-builders of Bristol.

A certificate accompanied these letters, from Mr. William James, and Mr. Samuel Haft, flip-builders, and also from Mr. George Winter, of Briftol, teffifying that they had tried

\* From the Transactions of the Society for the Encouragement of Arts, &c. for 1801. The Society adjudged a reward of forty guineas to the inventor. Models are in their repository.

the experiment of driving copper bolts through the jointed Evidence of its cylinder invented by Mr. Phillips; and that they so far approve utility. of it, that they mean to adopt the general use of them, for driving bolts in all directions, particularly on the outside of ships, whether iron or copper; as this method not only prevents the bolts from bending, but keeps the heads from splitting, and enables the bolts to be driven much tighter, than by any other means with which they were acquainted. They surther add, that by the application of Mr. Phillips's cylinder and punch, a copper bolt which had been crippled at the edge of the hole, and which could not be started by a mall, went up with ease in a perpendicular direction in the slat of a ship's bottom, not four feet from the ground.

This certificate was witneffed by Mr. William Holden.

The fame facts are also certified by Mr. Thomas Walker, and Mr. James M. Hillhouse, of Bristol, who add their opinion, that the adoption of this invention in the different dock-yards of the kingdom, will prove very advantageous.

Since Mr. Phillips's first application to the Society for a premium, he has made a confiderable improvement in the construction of his tubes. The description and engraving hereunto annexed are of the improved kind: models of both are, however, preserved in the Society's repository, for public inspection.

The inftrument employed for driving the bolts, confifts of a Description. hollow tube formed from separate pieces of cast iron, which are placed upon the heads of each other, and firmly held thereto by iron circles or rings over the joints of the tubes. The lowest ring is pointed, to keep the tube steady upon the wood. The bolt being entered into the end of the hole bored in the wood of the ship, and completely covered by the iron tube, is driven forward within the cylinder by an iron or steel punch, placed against the head of the bolt, which punch is struck by a mall: and as the bolt goes further into the wood, parts of the tubes are unscrewed and taken off, till the bolt is driven home into its place up to the head.

The tubes are about five inches in circumference, and will admit a bolt of feven-eighths of an inch in diameter.

Reference to the Engraving of Mr. R. Phillips's Method of driving Bolts into Ships. Plate III. Fig. 3, 4, 5, 6.

### Fig. 3.

A. The copper bolt, with one end entered into the wood, previous to fixing the tube.

B. A piece of timber, or ship's side, into which the bolt is intended to be driven.

### Fig. 4.

CCCC. The parts of the iron tube fastened together, ready to be put on the bolt A.

DDDDD. Iron or brass rings with thumb-screws, placed over the joints of the tube, to hold them firm together.

EEEEE. The thumb-screws, which keep the rings and tubes firm in their proper places.

F. Two points formed on the lower ring: they are to flick into the timber, and to enable the tube to be held firm in its place.

## Fig. 5.

Shows the separation of the parts of the tube, which is effected by slackening the thumb-screws and rings.

To put them together, you slide the rings over the joints, placed as close as possible; then, by tightening the thumb-screws, you will have them firm together, and may continue the tubes to any length, from one foot to whatever number is required.

### Fig. 6.

GH. Two steel punches or drifts, to be placed on the head of the copper bolt within the tube, whilst driving. The blow given upon the punch drives sorward the bolt. The shortest of them should be used first, and, when driven nearly to its head, should be taken out of the tube, and the longer punch applied in its place.

#### VIII.

Reply to Dr. Young's Letter on the Theory of Compound Sounds.

In a Letter from Mr. John Gough.

#### To Mr. NICHOLSON.

SIR,

You have published Dr. Young's answer to my remarks on On the proper the theory of compound founds; and I doubt not but you will, spirit of controwith equal impartiality, find a place for the following reply in your Journal. When a controverly is conducted with good. humour, it affords amusement; and when it furnishes fresh facts, or places the fubject of dispute in a new point of view, it proves instructive. Dr. Young's answer possesses both these qualities; and it is my duty to imitate him, at least in the obfervance of the former. Having now completed what may be called the introduction of the prefent letter, I will, with the permission of Mr. Nicholson, make a few remarks upon the principal objections contained in the answer under considera-The Doctor observes, that my theory of vibrations in an Explanatory reelastic fluid differs as widely from Dr. Smith's hypothesis as marks. from his own conceptions relating to the same operations: but he will probably recollect, that I profess to maintain compound founds to be mixtures of elementary founds, not aggregates by coalescence; in other words, I undertake to defend Dr. Smith's proposition, as a fundamental maxim both of harmonics and the general philosophy of founds. When the question is thus properly stated, the engagement I undertake to perform cannot in any fense oblige me to support the collateral arguments, with which Dr. Smith thought proper to elucidate his proposition, perhaps unfortunately. For if the positive maxim, riz. that That compound compound founds are mixtures, be established, the design of founds are mixmy effay is accomplished: and as Dr. Young does not controvert the leading conclusions of this paper, I may fairly suppose he admits them, but am at a loss to understand in what manner they favour his fide of the question; for if my inferences be just, compound founds are not formed by coalescence, which and not formed is the point in debate. I do not pretend to fay precifely what by coalescence, would be the confequence of conducting a number of founds through

Fact respecting bells, of which the founds are faid not to coalefce.

through a narrow tube, because I do not recollect a circumstance of the kind; nor do I think myfelf bound to determine the effect, because an instance, which is so particular, can have no weight in a dispute concerning a general principle. I am acquainted, however, with one circumstance which deserves attention at prefent: the notes of two or more bells are not obliged to coalesce by passing through the narrow sound-holes of the steeple of a country church: and if Dr. Young will suppose the point A, in the demonstration of my third proposition, to be placed in the mouth of his tube, he will fee that the motions of the corpufcle will not be disturbed by the edge of the pipe. The Doctor's answer blames my representation of his idea of a compound found, as amounting to a charge of ignorance in the most common occurrences. The representation appeared to be a necessary introduction to the objections which were levelled, in the fequel of my essay, at the Doctor's theory, not at his experience; and if the statement be not just, it is difficult to discover the sense of the term coalescence in that section of his paper in which the subject is examined, especially when the author infers, that the strength of the found in a concert is not in exact proportion to the number of instruments composing it. The writer of this paragraph, without doubt wished his readers to conclude, that the vibrations, which are communicated to the air by a number of founding bodies, reduce themselves to a single set of vibrations by mutual opposition: Now my objections to this doctrine must remain in force, until mankind can be convinced that unity possesses the qualities of number. The answer corrects the new theory in two essential points:

First, It gives a power of analyzing compounds to the ear; but this power cannot be admitted before the doctrine of coalefcence is established: Second, It reduces the coalesced compound to a pigmy, and joins to it the powerful reflected parts of its constituents, thereby forming a mixture. A few remarks On the mixture on the range of such a mixture shall close this letter. When a of founds differ- traveller approaches a town, he frequently hears the bells in it at the distance of four or five miles; the beating of a drum may catch his attention at one-third of that distance; when near the place, he perceives the blows of hammers, and at last a mixed noise. In his journey, then, he meets with a succession of founds at the limits of their respective ranges, but does not find

ing in their ranges.

the coalesced compound in the van of the train. If the preceding statement be natural, the range of a mixture of founds must be estimated by means of the strongest individual in the aggregate, not from the coalefeed compound, admitting its existence. This conclusion holds good in a general sense; at the fame time I know, that the ranges of certain aggregates. exceed the ranges of their strongest constituents. Of this kind Sounds of agis a general discharge of fire-arms; but though I admit the gregates heard fact, it does not appear to be an inflance of coalescence. For their loudest the report of every musket employed is nearly alike; and the constituents, musquetry. explosion of the whole number, at the same moment, agitates a large field of air. Now I know, by observation, that when an extensive surface is made to vibrate it sounds to a great diftance; for when the shear-men of Kendal beat the tenters on Shearmen beata calm morning, the strokes may be heard for two miles, or ing the tenters heard two miles. more; though the ear of a by-stander is not much affected by them, the operation being performed by striking the stretched cloth with a flick not thicker than a man's finger. Seeing then a number of founds which are nearly in unifon, forms an aggregate of more power than any one of its constituents, it follows, that when a multiplicity of instruments contains a strong combination of unifons, that combination determines the range of the concert.

JOHN GOUGH.

Middleshaw, near Kendal, August 19, 1802.

#### IX.

A new Process for Claying Sugars, proposed by Cir. HASSEL LACHENAIE, Chief Apothecary of the Military Hospitals of Guadaloupe to the Agents of the Confuls of the French Republic in the Windward Islands ..

(Concluded from Vol. II. Page 190.)

A HE third and last which I shall mention, though the least Economy of of the inconveniencies of the process of Citizens Boucherie, time. cannot be reconciled with the economy which we are obliged to use in the employment of our time; namely, the necessity of graining the fugar before it is put into the cases, &c.

After

Process of the author.

After having pointed out the principal difficulties and inconveniences which would refult in the manipulation of the claying of fugar, by the use of cases constructed after the model of those of Citizens Boucherie, nothing more remains to be done, than to flew how I have succeeded in overcoming these difficulties, by using cases fabricated according to my own principles.

Construction of the cases or receptacles for claying fugars. ference to the drawings.

My cases are without a bottom; they are constructed of four boards, united in a fquare by tenons and mortices, which are held closely together by wedges and pins, so as to form a described by re-fquare vessel, each side of which is three feet long, inside This case therefore presents a surface of nine square measure. feet. The fide h (fee Plate XII. Vol. II.) is 18 inches high at its extremity e, and  $16\frac{1}{2}$  inches at its other extremity f. The fide l, parallel to h, is cut in the fame proportions. height of the fide k corresponds with that of the extremity l; and the fide i, has the fame height as the extremity e.

I place four cases, of the construction just def ribed, upon a frame eight feet long, or plank, m, m, m, m, having a fall of three inches, and raifed two feet above the ground qq. board ferves as their bottom. In the space occupied by each case, it is perforated with twenty holes of an inch in diameter, placed in four rows: these holes, which I stop beneath with pins projecting eight inches above the bottom within, ferve for the drainage, when, after the fugar has cooled, thefe fame pins, which perforate it almost to the half of its thickness, are drawn out.

For receiving the fyrup, I place below the board feveral channels which convey it into a common trough intended to conduct it into a refervoir. Thirty-two cases disposed in this manner upon eight boards, in my fugar work, will fupply the place of 900 forms, and as many pots.

It is evident that by thus inclining the board which ferves as a bottom to the cases, it was indispensibly requisite that I should give the same inclination to the lower edges of the sides, in order that this upper part might be at the same level; and this inclination was the more necessary, as it favours the running off of the fyrup at the borders, where the cases are in contact with the board, no less than is done by the holes with which it is perforated.

After

After describing the construction and disposition of the cases which I propose, I shall now point out the manner in which I use them.

In order to avoid very fatiguing transportations, in which Manufacturing much time would be loft, I have established a cooler in my process. fugar-work, into which the boiled fugar is conveyed by a gutter from the battery. Near this cooler I have placed two backs, each of the same capacity with one of my cases, and thefe last are all of equal dimensions.

#### MANIPULATION.

I draw off my fugar by two batteries \*; when thefe are The fugar first joined together in the cooler +, I pour them into one of the cooled in a back, backs where the fugar crystallizes by cooling. This fugar is ftirred two or three times during the interval, until two other batteries have been poured into the refrigeratory: I then fill up my back, and with a paddle I mix well together the fugar thus collected; after which I leave it at rest till it acquires the confistence it is suffered to assume, when intended to be manufactured into raw fugar. In the interval other fugar is made, to be poured in the same manner into the second back.

As foon as the fugar has acquired the before-mentioned con- then put into fiftence, I use small buckets for conveying it into one of my the cases to cases, which I fill to the height of fix lines below its margin. I then leave it till coated, at which period I draw out the pegs from the board, in order that the fyrup may run out.

On the next and the following days, one of the fides of the eafily examined cases may be opened to observe how the drainage goes on; to ascertain the time for claying. but it must immediately be replaced. This facility, which has the advantage of afcertaining the most favourable moment for claying the fugar, enables us also to ascertain at pleasure the

\* The battery (batterie) is the vessel in which the boiling of the fugar is terminated, and when two boilings are united in the refrigeratory, it is customary in our manufactories to call this process drawing off the sugar by two batteries (tirer le sucre par deux batteries, l'enformer par deux batteries). These two batteries constitute the filling (emplie). The name of battery, which ferves to designate this boiler, is given to it because the sugar is beaten with a paddle, or stirrer, whenever it tends to raise itself above the sides.

+ This is a large boiler which receives the boiled fugar, and in which it is fuffered to lose its heat before it is drawn off.

effect of the claying. When, by examining the fugar in this manner, we find that there does not remain more than about two inches discoloured by the syrup, it is the proper time for claying it.

Preparation for the claying.

To prepare the fugar for receiving the clay, the very thin crust formed at its surface is first taken off, and is divided and spread out again upon this surface, which is pierced to the depth of six lines with the edge of the trowel, which is afterwards used for slattening and compressing the bottom. After this last operation is performed, it is clayed in the usual manner.

Qualities of the clay.

The fat earth or clay which is generally used here for claying of sugars, has in general the fault of retaining the water with which it is diluted too long, on which account it happens, that whilst one part of this water filtrates into the sugar, another, and that not the least considerable, evaporates into the air.

Defect from tenuity.

This earth has also another fault, which is very prejudicial to the operation of the bleaching of the fugar; namely, that of remaining for feveral days in part suspended in the water, before it is intirely precipitated from it, which renders this fluid milky. Both these saults require to be corrected.

After some trials I succeeded in doing this completely: my process is simple, easy to be executed, and requires neither expence nor loss of time; I can even affert, that the water with which my earth thus prepared is diluted, deposits nothing by filtrating into the sugar, but that it separates as limpid as the purest rain water. I send you some sugar bleached by this means.

Advantages refulting from the Use of my Cases and Manipulation for the claying of Sugar.

### FIRST ADVANTAGE.

With Regard to the Weight.

Statement of advantages in this new apparatus and process.

By cafing the fugar in the manner here described previous to subjecting it to the claying process, I prevent the formation of the fountain (la fontaine) \*, and of the thick crust with which every form presents us. Now, not having to separate the

This is described at page

fountains

fountains of the twenty-fix forms which each of my cases con-Less crust on the tains, the quantity of the sugar of these same fountains remains surface. therefore in the mass. This mass also suffers a much smaller loss in the claying operation, than it would have done if this operation had been performed upon twenty-fix separate forms.

### SECOND ADVANTAGE.

With Respect to the Regularity and Uniformity of the Grain.

When fugar is made to crystallize in our forms, the crystals The grain very that are successively precipitated from it, diminish also successively in volume in rising towards the top of the form which, in its position, then represents an inverted cone. In order to avoid this inconvenience, it is customary to stir the sugar as soon as its crust becomes folid. Nevertheless, a fountain is almost always formed, and for this reason: when the mass contained in our large forms begins to cool, the consistence which its still liquid part assumes, in proportion as it loses the caloric by which it was liquested, resisting the force of attraction which tends to unite the crystalline particles symmetrically together, in order to constitute regular crystals, these particles are precipitated, and apply themselves consustedly together, so as to produce the solid crust which constitutes this fountain.

The greater uniformity of crystallization in the mass which Cause why the my cases contain, (though I have disturbed it by agitating it grain is better-several times), compared with what happens in the forms which have not been moved, and even in those which have been, with equal care and dexterity, is obtained at the moment when I case my sugar. The consistence which it has acquired no longer permits its crystals to obey the laws of statics; but the smaller being consounded with the larger, the whole presents a perfect uniformity, not to be sound in those forms which have been moved. For in them this operation is performed at a moment when there still exists a sufficient degree of liquidity to permit the larger crystals to obey the same laws.

The reason why no syrup is collected above the sugar depo-No syrup colsited in my cases, depends on the separation of the crystals, lested at the topwhich being not at all agglutinated together, leave intervals
sufficiently large to receive it. But as this syrup still retains
much of its heat when the sugar is cased, it deposits a sufficient number of crystalline particles in cooling to agglutinate
the crystals, and to cause the whole mass to settle by its own

weight

weight in proportion as the fyrup runs off, and acquire a force of adhesion equal to that of the sugar put into forms.

#### THIRD ADVANTAGE.

As to its Whiteness or Colour.

and much less at The cased sugar bleaches as well under the clay, as that the base.

which is put into forms, and its base retains proportionally less syrup than the head of the form.

#### FOURTH ADVANTAGE.

With Respect to the Manipulation.

COMPARISON. 1. Less Labour and loss of Time.

Numerical statement of the saving of labour.

When we fabricate fugar, we employ every day a man to wash the forms, to carry them to the fugar house, and to plant them.

For putting the fugar into forms, two men carry it and divide it in the forms, which they fill. When these forms are sufficiently cool, they carry them into the resining house, in order to put them upon pots into which their syrups drain.

After fome days drainage, the hottom of each form is deeply perforated, in order to extract the fountain which is feparated from it. When this bufines is finished, the forms are again carried to other pots, destined to receive their fine syrup, after which the bottom is disposed for the claying.

In treating 416 forms in this manner, four men generally work 51,711 hours, which, multiplied by 4, amounts to 206,841 hours work for one man.

By the use of the cases, I avoid the necessity for employing this man, and the loss of this time.

For casing the fngar, it is transported, one single time, by small buckets into my cases, which are close to the backs. This operation is much less fatiguing than that of the forms, which weigh about 80 lbs. at the moment when they are thus conveyed.

When, on opening one of the fides of my cases, the sugar is found to be in the proper state for claying, its bottom is prepared as I have described, page, after which the diluted earth is poured upon it.

In this manner four men clay 16 cases, which contain as much as 416 forms, in the space of 2,840 hours, which multiplied by 4, amounts to 11,360 hours work for one man.

Thus the labour of claying 416 separate forms, with respect The new meto the time employed in doing it, is to that for the fame num-thod of claying ber of forms contained in 16 cases, as 206,844, to 11,360, or about one twenas 51,711 to 2,840. Now therefore, in order to clay the tieth part of the quantity of 416 forms of fugar in my 16 cases, (reckoning the usual timeday's work of each man at nine hours) I employ only one day and 2,360 hours, whereas the claying of 416 feparate forms requires 22 days and 8,844 hours.

I find also another faving of time and labour when the earth is to be removed and the bottom cleaned. This work, which requires a day's work of four men for the 416 forms, is performed in half a day by one man, upon the same quantity of fugar in the cases. I have made this comparison with my

watch in my hand.

### 2. Fewer Difficulties in making the Bottom.

Though our cultivators are in the habit of working the forms and making their bottom, it frequently happens that they neglect to remove the fountain completely; frequently also they do not uniformly compress the grained fugar which they had taken out of the form, in order to separate this fountain, and which they replace in it before making the bottom. In that case, the filtration of the water of the clay does not take place uniformly, but meets with obstacles towards fome of its fides; and the bottom, after this claying, exhibits distinct marks of the defective manipulation.

In my cases, as there are no The bottom is . fountains to be separated, as levelled more the crystallization is uniform throughout the whole mass, the furface of which is level before the cooling of the fugar, it is sufficient that this furface be very flightly opened before levelling the bottom: this bufiness is very easily performed, and those workmen to whom I have confided the execution for the first time, have perfectly succeeded. The uniformity of the crystallization in my cases, also favours that of the filtration of the claying water into the mass of the fugar.

### 3. Another faving of Time and Labour.

The forms being fufficiently in order to be conveyed to the

When I find the fugar fuf- The drying and drained, the fugar is taken out ficiently drained in my cases, floving is per-formed with I take them down, and leave more facility and

them effect.

stove, where the loaves are deposited upon frames, after the head, which is still wet with fyrup, has been separated from them. For, without this precaution, it would flow back into the mass, diminish its whiteness, and render the fugar more liable to attract the moisture of the atmosphere. Some days afterwards these forms are cut in pieces, in order to complete the deficcation of the fugar, and to feparate the whitest parts from the more coloured.

them for fome days exposed to the air, in order to deprive them of part of their humidity which is retained by that of the forms, when carried immediately to the stove: I afterwards break them into large pieces \*, and at the same time I feparate the whiter fugar from the reft. By this means the workmen are faved the labour of dividing the fame fugar, as is commonly done, in a hot stove which frequently must not be suffered to cool without the danger of injuring the quality of the fugar, and which, on account of the length of time they remain in it, disposes them to contract dangerous diseases.

#### FIFTH ADVANTAGE.

The last of the advantages which I shall mention, and which is not the least important, is that of economy.

It is very expenfive, and sometimes difficult to procure the forms and pots; but this apparatus is had.

It is known that it would cost an immense sum to those who, like myfelf, being without forms and pots, are under the necessity of providing cheap and easily themselves with them, fince it is almost impossible to procure them, now that the Saints and Martinique are in the power of the enemy.

With little expence, I construct and establish 32 cases. which supply the place of 802 forms and as many pots, and by this means avoid the great losses occasioned by the daily rupture of these vessels.

I think it unnecessary to recapitulate the observations contained in this memoir, and only wish that you may confider them as having fome title to your approbation.

\* Which are likewise carried into the stove, but which dry more speedily than the loaves.

X. Description

### X.

Description of a Water Wheel. By Mr. J. BESANT. In a Letter to Mr. Charles Taylor, Secretary to the Society for the Encouragement of Arts. From their Transactions.

SIR,

I BEG leave to lay before the Society some observations re- The common specting the common Undershot Water-wheel, and to point undershot wheel out the superiority of that of my invention.

First,—In common water-wheels more than half the water lets much water passes from the gate through the wheel, without giving it any pass without ef-

affiftance.

Secondly,—The floats coming out of the tail-water are re- The floats are fished with almost the whole weight of the atmosphere, at the refished at their instant they leave the surface of the water.

Thirdly,—The fame quantity of water which passed be- and they tend to tween the floats at the head, must of course pass between list the tail wathem at the tail, and consequently impede the motion of the wheel.

In the water-wheel of my invention,

First,—No water can pass but what acts, with all its force, In the new on the extremity of the wheel,

Secondly,—The floats coming out of the water in an ob- the floats are lique direction, prevent the weight of the atmosphere from oblique. taking any effect.

Thirdly,—Although the new water-wheel is heavier than It is in part that on the old construction, yet it runs lighter on its axis, the buoyant;

water having a tendency to float it.

Fourthly,—By experiments made with the models, proofs and is good for have been shown, that the new wheel has many advantages tide mills. over the common wheel, and that, when it works in deep tail water, it will carry weights in proportion of three to one, so that it will be particularly serviceable for tide-mills.

I hope on trial, before the Society, my invention will prove

fuccessful, and am,

Sir,

Your obedient Servant,

No. 26, Brompton.

J. BESANT.

To the SECRETARY of the

Society of Arts, &c.

Vol. III .- SEPTEMBER, 1802.

E

Repeated

Tried with fuceefs by the Society.

Repeated experiments of the above invention were made by the Committee; from the refult of which it appeared to possess some advantages over the common wheel, and to have a greater power of action.

Description of the late Mr. Besant's Water-wheel. Plate III. Fig. 1 & 2.

Description of the engraving.

- A. The body of the water-wheel, which is hollow in the form of a drum, and is fo constructed as to be proof against the admission of water within it.
  - B. The axis on which it turns.
- C. The float boards, placed on the periphery of the wheel. Each board is obliquely fixed firm to the rim of it, and to the body of the drum.
  - D. The refervoir, containing the water.
- E. The penstock, which regulates the quantity of water running to the wheel.
  - F. The current of water which has passed the wheel.
- Fig. 2. Is a front view of the water-wheel, shewing the oblique direction in which the float boards C are placed on the face of the wheel.

### XI.

Description of an Hydrometer which gives the Specific Gravity by Inspection, constructed by Mr. Atkins .- W. N.

Improvements in Atkins's hydrometer.

SINCE the publication of the description of Atkins's hydrometer in the last Number, that artist has communicated to me The weights of two improvements in the same. The first confists in making different figures. the four weights of different figures, namely, round, square, triangular, and pentagonal; and he stamps the figure of the weight on the fliding rule, opposite to every letter in the feries to which it belongs. By this contrivance, which indicates a confiderable degree of fagacity with regard to the practical requifites of an instrument offered for general use, he renders it impossible for the revenue officer to mistake one weight for another, or to take out his refult at a wrong part of the sliding rule.

The

The other alteration, though it will be highly acceptable to Hydrometer philosophical men, and renders the instrument of more exten-having the sp. five utility, and independent of the fliding rule, feems not so ated on its stems clearly to be an amelioration with regard to its most frequent and daily use. Instead of the letters of the alphabet, the four faces of the stem carry graduations which at once point out the fpecific gravity of the fluid according to the form now univerfally adopted; namely, by taking water as unity. Its figure is shown in Plate IV. Fig. 1. and it is provided with three weights; the inftrument itself being used without a weight at the upper range of densities.

Upon this occasion I cannot avoid noticing the strange over-Remarkable

fight of philosophers and others, who have expressed the den-oversight of philosophical chefities of acids and other fluids by certain graduations or degrees mifts in noting of instruments, such as that of Baumé and others; the com-specific gravities parative values of which were either little known to the pub- of fome particulic, or ill determined; fo that a very effential part of what we lar inftrument. read in books of chemistry is to many readers void of meaning. This is the more remarkable, fince the expression of specific gravities before mentioned has been long established, and is quite as easy to be put upon an instrument as any other notation. It is now seven years fince I urged this matter in my Long ago nochemical dictionary at the article Hydrometer, and the evil ticed by the auftill remains. If my recollection be accurate, I think Brisson and by Brisson. made the same proposal many years ago in the Memoirs of the French Academy. In the present instance, Mr. Atkins gains Advantages in the advantage of simplifying his sliding rule by leaving out the simplifying Atalphabet, fo that the operator looks for the specific gravity in rule; the first instance, and upon the opposite lines he finds the

But still more it is worthy of notice, that if he possessed the and rendering instrument only, or should in preference be desirous of refer-the instrument ring to Gilpin's tables, or any other elementary course of ex-tables, &c. periments, he can proceed immediately from the inftrument to the table without using the sliding rule to determine his specific gravity, as he must with the other.

strength and concentration.

I was at first very strongly impressed with the facility of this instrument in giving a result which ordinarily demands to be computed, and I suppose the advantages above stated will secure it the preference with many. But upon reflection, I But the alphacannot but think the alphabetic inftrument intitled to the pre-

ference adapted for dif-

rity against un-Kilfulnefs.

patch and secu- ference for commercial purposes, especially in the hands of men unused to make experiments, or to read the graduations of a rule. It is so much more easy to pitch upon a single letter than to read a graduation by stroke, as well upon the instrument as the scale, that I apprehend the literal instrument, with the variously figured weights, will in general be attended with greater certainty in the hands of all but philosophers. And for this reason the inventor is undoubtedly right in offering both to the public.

Cylindric bulb.

The cylindrical figure of the bulb affords the means of diminishing the fize of the assay jar, and consequently will demand a less quantity of the spirit to make the requisite trials.

#### XI.

On the Air from Finery Cinder and Charcoal, with other Remarks on the Experiments and Observations of Mr. Cruickshank. In a Letter from Joseph PRIESTLEY, L. L. D. F. R. S. 4c. &c.

#### To Mr. NICHOLSON.

DEAR SIR, Northumberland (America), Ap. 16, 1802.

MR. CRUICKSHANK having added a supplement to the account of his experiments on the air from finery cinder and charcoal, in answer to my objections to the new theory from the properties of that kind of air, I have given more particular attention to it, and wish you to add the following observations to those which I have already sent you on the subject.

The air from from charcoal stated to differ Inflammable air only in the proair it affords by combustion.

When I first procured this kind of air, I was far from imafinery cinder and gining that I had discovered any new species of air, essentially different from any other, so as to be entitled to a new appellafrom other dense tion, but only another variety in the heavy inflammable air, which is known to be exceedingly various in different proportion of fixed ceffes, and even in the different stages of the same process, all however agreeing in this, that when all fixed air is carefully washed out of them, more is found on the decomposition of them when they are fired together with dephlogisticated air,

And

And in this effential property the air from finery cinder and charcoal agrees with them all, differing only in the proportion of the fixed air procured in this manner. This, however, I now find must be called the gazeous oxide of carbon, while the others are called hydro-carbonates; that being faid to confift of two parts oxigen to one of carbon, and the others to be a folution of carbon in hidrogen, or the light inflammable air.

It cannot, however, be denied, that this gazeous oxide of Itis inflammable carbon is inflammable as well as the hydro-carbonates, in the dro-carbonates, composition of which a portion of hidrogen (one of the compo- and cannot with nent parts of water) is a necessary ingredient. This air, there- any confishency be called an oxfore, from finery cinder and charcoal, though called an oxide, ide, because it is must be essentially different from all the other oxides, none of combustible. which are combustible, being substances already saturated with oxigen. Thus iron is a combustible substance, ready to unite with oxigen when presented to it in a proper temperature; but when it is faturated with oxigen, and therefore called an oxide of iron, it is no longer combustible. It must, therefore, as it appears to me, be an absolute abandonment of one of the most fundamental principles of the new theory, to call the air from finery cinder and charcoal an oxide. If substances be combustible in proportion to their affinity to oxigen, and their confequent readiness to unite with it, this air, which is inflammable, must be of this class, and therefore the very reverse of the oxides, which are faturated with oxigen, and incapable of receiving more.

If this kind of air was a real oxide, it would appear to be Its oxigen has fo on the decomposition of it; when, to make the result un-never been sepaexceptionable, the oxigen it contained would either take the by transposition, form of dephlogisticated air, or become a component part of as happens with fome other substance into which oxigen was acknowledged to real oxides. enter. But this has not been done. When it is decomposed by being fired together with dephlogisticated air, the fixed air which is then formed comes, I have no doubt, from the oxigen in the dephlogisticated air, and the phlogiston in this species of inflammable air; the fame being the refult, though not quite in the same degree, of firing the heavy inflammable air from charcoal and water, from oil, &c. &c. into which it is not pretended that any oxigen enters.

#### AIR FROM FINERY CINDER AND CHARCOAL.

Mr. Cruickshank's observation of the union of the oxigen in the dephlogisticated marine acid air with inflammable air in the common temperature of the atmosphere, is extremely curious but I do not see that any thing more can be inferred from it, than from the more rapid union of the fame principles when inflammable and dephlogisticated air are fired together.

Objection to Mr. Cruickshank's theory of the re-

Mr. Cruickshank says, p. 201, that "the oxigen in the finery " cinder, uniting with the carbon in the charcoal, forms fixed duction of finery " air; and that the metal being in the same process revived, cinder and char- "decomposes this fixed air, when it becomes again to a cer-" tain degree oxigenated." But why should not the finery cinder retain a part of its oxigen, rather than first part with it, and then take it again? Besides, it is not true that after this process the iron is in any degree oxigenated, for it is completely revived, being perfect iron; and that any fixed air is either formed or decomposed in this process is altogether conjectural, and for the reasons that I have given cannot be admitted. For though it might be possible for oxigen in the finery cinder (supposing it to contain that principle) to be extracted from it by its affinity with the carbon in the charcoal, and that nothing should enter in its place, the iron thus revived could not decompose the fixed air that would be formed by their union.

> Not doubting but that Mr. Cruickshank will give a candid attention to these observations.

DEAR SIR, Your's fincerely, I. PRIESTLEY.

#### XIII.

Description of Mr. Read's Pneumatic Apparatus. By a Correspondent.

#### To Mr. NICHOLSON.

SIR,

I BEG leave to recommend to your notice a very fimple and Notice of a very cheap apparatus for displacing any elastic study out of a referfor the gases.

voir by means of water. It is described in Dr. Beddoes's Considerations on the Medicinal Use of Factitious Airs, octavo,

London, 1796; and I hope you will think with me, that it deferves a corner in your respectable publication.

I is a funnel with its pipe or tube stopped at B, and perso-Description, rated above and below with two or three small holes. Round the tube A is soldered the tube D, lest open at top. The tube G is made to circumscribe the tube D, and to be soldered to

the funnel at F. and likewise soldered at E.

When water is poured into the funnel, and the cork K K in-Effect. ferted into a bottle filled with gas, the water descends through the tube A till it arrives at the division or section B; it then slows through the small holes at C, and ascends between the tubes A and B; slows over the top of D, and descends again between the tubes D and G, till it arrives at E (where it regains the tube through an opening above E), and thence into the bottle. As the water goes in, the air escapes between the tubes A and G, through the cock H, into the mouth-piece. The lower end of G is made to perforate a cork which is properly attached to it, and which secures it air-tight into the neck of a common bottle, as seen at fig. 3.

The whole apparatus may be made for less than a guinea, It is very cheap, and may be introduced to the bed-side of the patient, if necessary, by having a cask filled instead of a bottle, and placed upon a chair. By turning the cock the patient may take an inspiration whenever he pleases, the cock being thrust into the

spiggot hole.

I am, &c.

N. N.

## SCIENTIFIC NEWS, &c.

Experiment of the Invisible Girl.

Description of the apparatus for the experiment of the Invisible Girl.

 ${
m Mons}$ . Charles, brother to the well known philosopher of that name, has for some time exhibited in Leicester-square an experiment called the Invisible Girl, which seems to include fome principle of acoustics either newly discovered, or not hitherto known to the extent there displayed. In the middle of a large lofty room in an old house, where from the appearance of the wainfcot, the position of some glass cases of natural history that occupy one fide, and from other circumstances, there does not appear to be any fituation for acoustic tubes or reflectors-is fixed a wooden railing about four feet high, and five feet wide, inclosing a square space. From the four corner posts of this railing there rise painted wires, or rods, to an eminence of ten or twelve feet, (the cieling being much loftier) and from the upper ends of these hangs, by strings resembling the lines used with curtains, a square glazed box, having a mahogany bottom nearly on a level with the top of the railing, and in the middle of this box is suspended an hemisphere (or larger portion) of tin or pewter, out of which proceed horizontally four tin trumpets at right angles to each other, and about eighteen inches long. The disk or hemisphere is covered with a portion of a quickfilvered glass globe which completes the sphere, and closes it imperfectly, so as not to prevent the inner aperture of one of the trumpets from being feen through the place of infertion. This whole apparatus is fo far detached from all furrounding objects, that it can be fwung about as far as the railing will allow; that is to fay, the box can be fwung by its four ftrings, and the globe and trumpets by the ftring that fustains them in the box. The globe may be about ten inches in diameter. In all these dimensions, which I do not suppose to be material, I speak from the recollection of some weeks ago.

Manner in which the voice iffues from the apparatus.

In the exhibition, the Invisible Girl is supposed or pretended to be included in the sphere, and the conversation is held by speaking into any one of the trumpets, and the reply comes out of them; that is to say, it is most clearly heard by applying the ear to one of their mouths. The voice is low, as if con-

The lady converses in several languages, sings, describes all that happens in the room, and displays a fund of lively wit and accomplishment, that admirably qualify her to support the character she has undertaken.

I hope to meet my reader hereafter with a little more to fay Experiment of on the nature of this experiment. At prefent I can only offer found conveyed the conjecture that the found is conveyed through the ftrings, by a thread, and to this I am led by another experiment of the celebrated Dr. Moyes. He shewed me, that if one end of a packthread, or even fewing thread, be wrapped round the finger, and that finger be put into the ear (not forcibly) while the other ear is flopped, and at the same time an affisfant wraps the other end of the string round the handle of a tuning fork, and (the string being then drawn rather tight) he causes the fork to found, too low to be heard by himfelf and the byftanders, it will nevertheless be very audible to him who holds the other end of the string. We tried it at an interval of fifty feet, and the Doctor has tried it at upwards of three hundred, without any fenfible diminution of the effect. I wrapped the firing round my wrist, and put my finger in my ear, and found the effect very nearly as ftrong. If instead of the tuning fork one observer bites hold Conversation of the string between his fore teeth, while the other applies it through a strings as before to his ear by means of his finger, the voice of the former may be transmitted; but it is more difficult to manage this experiment with fuccess than the other.

# Aerostatic Experiments of GARNERIN.

A new generation has fprung up fince the epoch at which Revival of across Air Balloons constituted the object of astonishment and specutation. In Balloons constituted the object of astonishment and specutation. It is a soft as a single period as if it were yesterday, must feel a little surprized at the vicissitudes of human assairs, when they find how many of those who are now intirely engaged in society, were then scarcely in existence; and how sew have seen a balloon. The spectacle exhibited by Mr. Garnerin, though much less impressive than the experiments of his countrymen to whom this invention is due, in all their novelty and originality have nevertheless been received with much interest. The readers Garnerin the of our Journal will recollect him as the first adventurer whose sirest operator with a paraintrepidity chute.

intrepidity and skill \* enabled him to descend through the atmosphere by means of a Parachute; and those who extend their refearches to the new relations and confequences of uncommon experiments, will probably confider thefe daring attempts as the embryon efforts towards the art of flying; which fo many eminent men have thought within the reach of human He has not, however, yet, in the uncertain climate of our island, been able to gratify us with the fight of his defcent in this manner. Three aerial voyages, perfectly refembling those of his predecessors, have constituted the whole of what he has yet been enabled to perform.

On the 28th of June he made his first ascent from Ranelagh

of the morning, the weather changed, and at five o'clock, when the whole was in readiness, there was a strong wind from the

fouth west, with occasional showers.

After the promife of a fine day by the indications

At this honr Mr.

Three voyages in England.

The first with Capt. Sowden.

Afcended two miles.

ing.

Singular narrative of Captain

Sowden.

Garnerin and Capt. Sowden afcended, and proceeded over St. James's Park, the Thames, and the bridges of Westminster and Blackfriars. After proceeding over the metropolis they threw out a quantity of ballast, and ascended to an elevation which Mr. Garnerin estimates at upwards of 10,000 feet above the furface of the earth, or nearly two miles perpendicular. At the expiration of about half an hour the valve was opened, and the balloon descended through dark clouds which had intercepted their view of the earth, upon fight of which they found themselves speedily advancing towards the sea. Dangerous land- On account of the tempeftuous weather their landing was very The wind dragged them over dangerous and unpleafant. fields and hedges, which tore their hands and cloaths, and when their anchor had engaged itself at last in a thicket near a house, the inhabitants were so far from affisting them, that they even appeared disposed to fire upon them. For want of this necessary help their cable broke, and they were dragged forward till the balloon was fufficiently lacerated to begin to collapse, and suffered them to jump out.

Capt. Sowden published an account of this excursion, in which he relates some singular observations, considerably differing from those of former aeronauts. He affirms, that his fight in looking down was fo ftrong, that he could eafily diffin-

\* Philof. Journal 4to, I. 523.

guish the minutest objects on the earth. In this vast Map, he could not only trace the different roads and interfections, but even the ruts and furrows in the field. His hearing became likewife furprifingly acute. The rattling of the carriages on the roads, the lowing of cattle, and the acclamations of the populace, were diffinctly heard at the height of fifteen thoufand feet; though at the fame time the travellers could scarcely hear themselves speak. He is persuaded that a person with a strong voice and a speaking trumpet, might be perfectly understood from the earth at that elevation in the air. Instead of finding the climate colder as he afcended, he observed the contrary.

The place of their descent was on a common four miles be- Account of their yond Colchester and fixty miles from Ranelagh, and the time landing near Colchester, after from their departure was three quarters of an hour; confe-travelling at the quently they travelled at the rate of eighty miles an hour rate of 80 miles They were very hospitably received by Mr. Kingsbery, a freeholder of the county, who imagined their vifit to be on account of the parliamentary election, in which he had determined not to interfere. By his very friendly fervices they were refreshed and proceeded to Colchester, whence they returned to London on the following day.

The fecond voyage was made on the fifth of July, from Second voyage, Lord's cricket ground, near Paddington. The very unfavour-July 5, 1802. able weather rendered his afcent much lefs firiking than it might else have been. Thousands of individuals filled the furrounding fields for some hours before the ascent, amidst rain nearly inceffant, and a continual haze. The balloon rose with Mr. Garnerin, accompanied by Mr. Locker, at ten minutes before five in the evening, and in four minutes afterwards it was loft in the mass of clouds. At the end of the fifth minute they began to descend, and landed at Chingford in Essex, after having remained in the air for one quarter of an hour. The most remarkable observation in Mr. Locker's account is, that neither himself nor Mr. Garnerin could distinguish sounds

On the 20th following a balloon was fet up from Vauxhall Pyrotechnic balgardens in the night, having an extensive system of fire works loon, July 20. attached to it, to be discharged by means of a fuze, at a great height in the air previous to the explosion of the balloon itself.

above the elevation of 3 or 4,000 feet.

This exhibition, which does not require any general philosophical explanation, was attended with the most complete success, and had a very striking effect.

Third voyage, with Mrs. Garnerin and Mr. Glasfurd.

The last voyage was made from Vauxhall Gardens by Mr. Garnerin, accompanied by Mrs. Garnerin and Mr. Glassurd, on Tuesday, the 3d of August, at a quarter past seven in the evening. The day was calm and serene, and Mr. Garnerin made the offer of descending by the parachute, which he says in his letter was rejected, but not by whom. Circumstances were highly savourable to the prospect. The travellers passed flowly over the metropolis in bright sun-shine, at an elevation which permitted them to be distinctly seen by a good perspective magnifying about 30 times, and for the most part they were much more clearly seen. A cat attached to a parachute was let fall from an height of about 600 feet. It descended swiftly at first, till the parachute was fairly expanded, and then came down slowly and with safety. They landed near Hampstead, about two miles north of the skirts of the town.

Promise of a disfertation on this object.

Notwithstanding the very ample discussion the subject of aerostation has had at the period of its invention, I apprehend that my readers will not be displeased to see an outline of what relates to it, with an inquiry into the probabilities of its becoming of any extended degree of utility; and this I purpose to do in our next.

# Commercial College at Hull.

Objects of a proposed commercial college. Dr. J. Alderson of Hull, has given out proposals for establishing a college in that town, for purposes highly interesting to that commercial intercourse which has in so many respects extended the powers and enjoyments of the civilized societies of modern times, and is so intimately connected with the improvement of the intellectual and moral habits of man. He states as the objects of the establishment, 1st, To obtain for the man of business precise information on the nature and value of every article of commerce, whether crude or manufactured; 2d, To point out the country where such articles are best to be procured; 3d, To render him acquainted with the various processes by which they are rendered marketable; and, 4th, To instruct him in the languages of the different countries to which the objects of commercial intercourse may lead him.

· The

The plan is to form a museum, and to erect a building for Plan. the accommodation of Professors, and the reception and display of specimens of raw materials and manufactured articles. One Professor to be engaged to teach the southern languages, another the northern, and a third to have the care of the museum and the office of explaining whatever relates to the specimens under his care. The estimate for the building and specimens, to be raifed by subscription, is 2,500l.; and it is proposed that the corporations should endow the Professors with 50l. per annum each.

The Doctor, who stands in the honourable situation of pro-Its great advanposer of an establishment so manifestly useful, disclaims the no-tages. tion of this being a charitable infiitution; but confiders it as the private interest of every subscriber. " As the father of a large family," he observes, "I certainly should not think myfelf at liberty to subscribe fifty pounds merely for the personal pleasure that results from the contemplation of a charitable action; but in subscribing fifty pounds to the above institution, I consider my interest and the welfare of my family. I am convinced it will add to the wealth and importance of the town; and in both of these I am deeply interested: but the greatest interest I have, is in the facility it will give to the education of my children. In a commercial country and a fea-port town, the acquisition of knowledge that evidently leads to the improvement of trade, becomes an object of the first moment in the education of youth; and the mode now proposed offers advantages which no private school can possess. To bring together under the immediate observation of youth (while yet the memory is most retentive), perfect specimens of the different articles of commerce, both of the raw material and of the manufactured, will enable him to bear in mind the precise value of any article he may be called upon to appreciate: To point out the country where every article is first procured or manufactured, will qualify him to go to the cheapest market, and by teaching him the language of fuch country, he will be at all times capable of transacting his own concerns without the intervention of interpreters, often a very ferious fource of impofition."

It will no doubt be of great advantage to the town of Hull, that this plan should meet with support, and to the community at darge, that it should be followed in other places.

A System of Chemistry in Four Volumes. By THOMAS THOMSON, M. D. Lecturer on Chemistry in Edinburgh.

Thomson's chemistry.

This work of Dr. Thomson (who is well known to the public by his excellent notes on an edition of Fourcroy's chemistry in three volumes, as well as by several important memoirs in our Journal, and other productions) is intended to exhibit a detail of the vast number of sacts which constitute the science of chemistry, blended with the history of their gradual developement and of the theories which have been founded on them, and accompanied with exact references to the original works in which the different discoveries have been registered. His plan or order of arrangement will be best seen by coumerating the contents.

After a short introduction, it is divided into parts, books, chapters, and sections. Part first relates to the principles of chemistry. Simple substances, viz. oxigen; simple combustibles, viz. sulphur, phosphorus, carbon, hidrogen, azote: Metals, 22.—Light; caloric; general observations.—Compound substances. Primary: Alkalis; earths, 10; oxides; acids. Secondary: Glass, salts, hidro-sulphurets, soaps.—Of affinity.—Examination of nature. Of the atmosphere; of water; of minerals; of vegetables; of animals.

To enter into a detail of the numerous interesting and highly valuable sacts and statements which abound under the heads above enumerated, would be impracticable in the short limits of a notice. I must therefore content myself with observing, that it is a clear, comprehensive, and accurate system, in which the learner will be orderly led from one truth to another, and the man of information will with ease find the compendium and references he may want; that it is beautifully and closely printed on sine paper, with side notes; and upon the whole must prove a desirable acquisition to our stock of means for the advancement of chemical knowledge.

Death of THOMAS GARNET, M. D. late Professor of Chemistry and Natural Philosophy at the Royal Institution, &c. &c.

Death of Dr. Garnet.

On the 28th of June last died of a typhous fever, caught in the gratuitous exercise of his profession, Dr. Thomas Garnet, a man who, for his philosophical knowledge, his assiduous exertions in diffusing information by his lectures, and his amiable

amiable and unassuming disposition in fociety, will be long regretted by the world, and by his friends in particular. He Some account of was a native of Westmoreland, and after a regular education, his life. studied physic at Edinburgh, which he afterwards practifed at Harrowgate. Subsequent to this period, he commenced a course of lectures on chemistry and experimental philosophy at Liverpool, which he repeated at Manchester with such fuccess, that when a vacancy offered in Anderson's Institution at Glasgow, he became a candidate for the appointment, and obtained it. In this fituation he acquired fo high reputation, that he was applied to by the Managers of the Royal Institution to be the first Lecturer in that establishment, which unfolicited honour he accepted, and delivered a very extensive and truly laborious course of lectures during the commencing fession of that corporate body, which were followed by the first ranks in audiences fo crowded, as perhaps were never before witneffed on a fimilar occasion.

Before he quitted Harrowgate he married a Miss Cleveland, whose talents and accomplishments were peculiarly adapted to his own. With this amiable friend and partner of his fortunes he passed a few short years happily, and became the father of two daughters. But the birth of the latter infant was fatal to her mother. It was foon after this event, and during the early exertion of his fortitude under fo great an affliction, that he became attached to the Royal Institution. His constant attention to his numerous occupations, supported by the hope of fecuring an independence for himself and the orphan representatives of their regretted parent, afforded the most rational means of confolation. But the energies of the mind cannot support the physical structure under accumulated labours, anxious reflections, and the folitary privation of that domestic fociety which choice and habit had rendered dear to the human heart. His health became impaired, but his determination to proceed, continued in all its vigor. He took an house in Marlborough street, increased his stock of apparatus, completed a lecture room, and his classes became highly respectable under him as the lecturer on his own private account. At this period it was, when he had applied the whole of his means and the utmost stretch of his powers to the well grounded plan of cftablifhing himfelf in the ufeful and honourable profession

Some accounts of Dr. Garnet.

he had adopted; at this period it was that death terminated all his hopes, and left his infants too young to feel the possible extent of their missfortune!

I have faid the possible, and not the probable extent of their misfortune. He himself in the latest moments of his life must have known-nay he did know-that the public he had endeavoured to serve and to instruct, have ever been alive to cases much less powerfully claimant on their feelings, their generofity, and their justice. These children cannot be forfaken by the numerous and affluent individuals who esteemed his virtues and revered his talents. When it shall be known that his Lectures on Zoonomia are to be published by a subscription destined to the purpose of affording them an humble, but independent support, they will most affuredly find the loss of their parent supplied in this respect, by that ready benevolence which is fo peculiarly characteristic of our country. is intended to address the Public at the season when the town fills; but in the mean time the Royal Institution have permitted the work to be printed at their press, and have ordered a donation worthy of the spirit and liberality of their views. Several gentlemen have also subscribed, and I am authorized to affure the friends of this laudable undertaking, that fubscriptions are received at the house of the Royal Institution. shall myself be very happy to afford any farther information to enquirers.

<sup>\*\*</sup> Wr. Notlem will have the goodness to point out the Volume and Page of the passage for which he withes to have an explanation.

### A

# JOURNAL

o F

# NATURAL PHILOSOPHY, CHEMISTRY,

AND

# THE ARTS.

OCTOBER, 1802

### ARTICLE I.

Objervations upon fome remarkable Wells near the Sea Coast at Brighthelmstone, and other Places contiguous. By the Rev. W. Pearson, P. R. I. From the Author.

IN Fisher's "Description of Brighthelmstone, and the adja- The wells near cent country," (p. 38, fourth edition) it is faid that "water is Brighthelmstone are said to be procured at Brighton from wells of considerable depth, and empty at high being filtered through chalk, of which these parts are one im-water, and full mense rock, it is of course persectly cleared of every species at low water, of foulness. The following phenomenon is observable in feveral of these wells; at the time of high water they are empty, and at low water they are full;" likewise under the article Rot- at Rottingdcan. tingdean of the same pamphlet we read, "On the road to Newhaven, at the distance of near four miles from Brighthelmstone, lies the pleasant and delightful village of Rottingdean. This place is remarkable for its wells, which are nearly empty at high water, but which rife as the tide declines. These affertions appeared on perufal fo fingular, that I formed a determination of availing myfelf of the opportunity which a temporary vifit to this fashionable bathing place gave me in the month of July last, of examining how far the statements in question are accurate; accordingly I obtained a plumb line,

and

Vol. III,-October, 1802,

This report is not accurate.

but was lately repeated in a

respectable

newspaper.

and measured the depths of the surface and bottom respectively of several wells in different weeks, and at different hours of the tide; the result of which measurements turned out to be a convincing proof that this traditionary report, though partially founded in fact, has been greatly exaggerated; but whether or not for the purpose of entertaining the visitors of this place, it would be presumptuous to determine. At the time, however of my vifit, it was not my intention to lay before the philosophical reader any observations which were made, but merely to register them by way of private amusement; which intention would have been adhered to, had I not met with a paragraph in the General Evening Post of Tuesday, Aug. 24, 1802, of which the annexed is an exact copy.-" Rottingdean, four miles from Brighton, on the road to Newhaven, is remarkable for the fingular variation of its wells, which rife as the tide declines, and are nearly empty at high water! The refearches of philosophy have not as yet been able to affix the cause of this apparent contradiction in the course of nature." This attempt to excite a more general aftonishment than was excited by the Brighthelmstone guide alone, and confequently to call the attention of the philosopher to investigate the subject before us, has induced me to recur to my notes, and to transmit for your Journal the deductions I had made from them, in order to show that there is no apparent contradiction in the phenomenon, as it has been called, of the Brighton Wells. The South Downs, which constitute a ridge of high ground

The South Downs lie on a bed of chalk floping towards Brighton;

to the north of Brighton, the fummit of which is parallel to the fea coast, have their gradual slope inclining towards the fea or fouth fide, and the declivity, which approaches more to a perpendicular line, facing the level country on the north; and the dip of the stratum of chalk, and consequently of its impermeable bed, is in a direction from the fummit to the fea, upon which the or from north to fouth; the rain therefore which falls upon these Downs may be expected to run down the bed of the rocks until it arrives at their lower extremity, where, if it meet with any opposing clay, or other impermeable substance, it will collect into a body of water, and, as it collects, rife through the crevices of the rock and superincumbent permeand form springs able earth, till it shews itself in the form of springs. foning, if my recollection be perfect, is agreeable to the theory of Mr. Pilkington, whose success in draining and tapping

forings

rains are conveyed;

at its lower extremity;

fprings is now pretty generally known; and in conformity to it I was led to fearch in the first place for the lower extremity of the rock, which I found was upon the shore, near the separation between the rounded stones and fand, at from thirty to forty yards by estimation below the bank. Here, as I had which make previously supposed, I found the fresh water continue to spring their way into or gush out constantly in a line extending parallel to the bank, distance of thirty both to the east and west of Brighton as far as I examined. or forty yards below the high-In this line then are fituated the natural springs of the place, water bank. which, if the fea were remote, would be not only perennial, but probably would emit an equal \* quantity of water at all times; but in every tide the fea covers this line of natural Thefe springs fprings, from three to fix feet we will suppose, according to are covered by the state of the moon, wind, &c. so that the fresh water is tide; prevented from making its escape by the heavier superincumbent falt water of the sea, the consequence of which is, that, from the time the falt water covers this line, the fresh water and consequentbegins to accumulate in the bed of the rock and inclined rock water is preitself, as high at least as the surface of the tide. During the vented from time therefore that the tide continues above this line, the fresh coming down; water is pent up in the rock at a distance sufficiently remote from the sea, to exhibit a perceptible rise in the artificial wells and the wells fouthward of the bank, which are dug down into the bottom become fuller; of the rock. That this should be the case to a certain extent it was natural to expect, but what period of time should elapse not at high wabetween the inftant of the tide's arriving at the line of the na-ter but later. tural springs, and the instant of a perceptible rise commencing in the artificial wells, at some distance from that line, can only be ascertained by actual observation, for both the distance of the wells, and number as well as capacity of the small natural ducts that convey the water must be taken into the consideration, if an attempt were made to inflitute a calculation, and I confess my memoranda are rather deficient in this respect; my attention having been directed more particularly to afcer-Observations as tain the states of the tide when the wells were at the two ex- to the facts. tremes of rifing and falling, which times are erroneously afferted to be about fix hours from being contemporary with the

The reason for this supposition is, that the quantity of inclined surface is very considerable, and is also contiguous to the sea, where showers are frequent.---W. P.

influx and efflux of the tide. The two wells, which were most convenient for examination, were, one immediately behind Fisher's library, and another in a little office at a small public house called the Dolphin, near the foot of East-street, which the fifthermen use for procuring water for tanning their nets; which two were nearly equally diffant from the bank, and subject to the same alternations of rising and falling.

Deductions.

The deductions which were made from a variety of meafurements were as follow:

The wells are fullest during the ebb.

1. The wells are most filled with water when the tide is retiring from the line of the natural fprings; evidently because the accumulation of fresh water continues as long as the falt water is above the faid line.

The fprings are

2. The whole continuance of the tide above the line of the hours by the fea; springs, or time of filling, appears on an average about sour

and uncovered fix hours. A shallow well becomes quite empty; a deeper not fo.

- 3. The whole time of emptying feems to be about fix hours.
- 4. The well at the Dolphin is about three hours quite empty in every tide, but the other, which is deeper, is never quite empty.

5. At two hours after high water on the 16th of July, viz. Particulars, well known. one day and a half after full moon, the water in the well at the Dolphin was two yards deep.

> 6. The fishermen who used the well at the Dolphin, informed me that they knew very well that the water in the wells rofe to the greatest height at two hours after high water, and were at the lowest two hours after low water, which information accords with my observations.

The water fresh.

7. The water always tafted fresh and pleasant.

A remote well fcarcely varies.

8. A well about 170 yards more remote from the fea than that of the Dolphin, in a direction towards the market, never varied more than three inches and a half in depth at the different times of observation.

Why the deeper well holds its water.

9. The well near Fisher's is eight yards and 30 inches deep, which depth is probably below the line of the natural springs, and therefore it contains water at all times; but the other is not fo deep, and therefore lofes all its water every tide.

The effect has thus b -n correctly stated and accounted for.

From these deductions it will, I presume, appear evident not only that the accounts which have been circulated respecting the wells at Brighton, Rottingdean, and other villages contiguous to the Suffex coast, are erroneous, but also that a fatisfactory.

fatisfactory reason may be given for their reciprocation, which, though not contemporary with the ebbing and flowing of the tide, is yet so far dependent upon it, as to follow it at an in- It resembles a terval of two hours, but in fuch a way that the immediate in-river tide, profluence of the tide upon it only lasts about four hours out of the fresh water by twelve, notwithstanding the effect produced in that time re-the sea: quires nearly fix more to put the wells into their original state later; and has of vacuity or rest.

Lastly, the same reasoning which accounts for the alternate than the flood. rifing and falling of the water in the artificial wells, makes it flowed down a more than probable, that if the line of the natural fprings had declivity higher than the highest been below the low-water mark, the artificial wells would tide, the fream have been not only required to be deeper, but would have would always been invariable in quantity; also, that if the faid line had been above high-water mark, it would have been in vain to have dug for water into the rock above, whilft there continued a natural issue below: and should it be asked, why the water that has an immediate communication occasionally with the fea yet continues fresh, the same answer may be given that will apply in the case of fresh water being driven back to Chelsea and Putney at every successive tide that enters the mouth of the Thames.

Parson's Green, Aug. 29, 1802.

### TT.

On the Theory of Galvanisin. By John Bostock, M. D. From the Author.

HE following Paper contains some remarks upon the elec-Galvanism and trical pile of Volta, principally made with a view to illustrate the fame. and confirm the theory of its action, which I proposed in my former communication.

1. With respect to the nature of the agent producing the phenomena of galvanism, there can scarcely remain any reafonable doubt of its perfect identity with the electric fluid. The apparent diffimilarity in the effects of an electrical machine and a galvanic pile has been fufficiently accounted for, without supposing any thing essentially different in their nature, and we feem enabled by proper management to produce from either

like that it is its ebt longer But electricity is produced in the pile.

either apparatus at pleasure every effect of galvanism strictly so called, or of electricity. There appears however to be a great difference in their mode of action; in the pile the fluid is actually generated, whereas in the machine it is merely transferred from one body, where it previously existed in a difengaged state, to another body. In support of this position it may be alledged, that the energy of the pile is found to be materially increased by being infulated, whereas in the common machine it is impossible to produce any considerable degree of excitement \*, unless there be a communication formed by a conductor between the rubber and the furface of the earth. Dr. Woollaston, who has thrown so much light upon this subject, feems to have carried too far the analogy between the methods of exciting the electric fluid, when he confiders the The pile is con- oxidation of the amalgam on the cushion, as more than a se-

as a collecting apparatus.

fidered as a gene-rating, and the condary cause in the operation of the electrical machine. electric machine confidering the pile as a generating, and the other as only a collecting apparatus, we may affign a probable cause for the vivid sparks, the bright flashes, and the more considerable marks of excitement which are exhibited by the common machine; in this latter case a quantity of electric fluid derived from the neighbouring bodies, is as it were forcibly detained in the apparatus, and is ready to escape to every substance for which it possesses any degree of affinity; whereas in the pile, though there is a rapid and conftant production of electricity, yet being derived from a change which takes place in its own composition, by which different parts of it appear to acquire different capacities for the electric fluid, it feems little disposed to enter into any combination with the furrounding bodies, and is only brought into action by the approximation of the different parts of the apparatus itself.

The effects of ascribed to its electricity at a low intenfity;

It was justly remarked by Mr. Nicholfon, that the peculiar the pile has been effects of the pile may be ascribed to the electric fluid existing great quantity of in it in a large quantity, but at a low degree of intensity. This confideration eafily explains all the fingular properties of the apparatus, but it still remains to be ascertained by what

> \* This is an overlight. In fact, the ground serves only to form a communication between the receiving conductor and the cushion, and the excitation is fully as strong when the current is made to fly from the plus to the minus conductor, when both are infulated .-- N.

means this difference of intenfity is produced. By different degrees of electrical intenfity it is generally understood, that the same quantity of uncombined \* electric matter may be diffused through a greater or less space, and in consequence of this, as it exists in a state of concentration or dilution, it will exhibit effects more or less violent. In the galvanic pile it might have been supposed from the small bulk of the apparatus, that a large quantity of electric matter could not have existed in it except in a concentrated state, but we find that the contrary is invariably the case; nor has any method been hitherto contrived by which the electricity of the pile can be made to exhibit the fame marks of intenfity with that of the common machine. Future experiments will probably enable us to de-but the nature termine with more precision in what the difference between a of the differences greater or less intensity consists; possibly the electric sluid, as not been exis the case with many other bodies, may differ in its nature plained. according to the method employed for extricating it from the fubstances with which it was previously united.

2. That the extrication of the electric fluid in the pile of The electricity Volta depends upon the rapidity with which the furface of the of the pile demetal is oxidated, after the experiments which have been pidity of oxidaperformed, and particularly those of Mr. Davy, feems to re-tion, &c. quire no farther proof. It is however not only necessary that the fluid should be extricated, but that when difengaged it be concentrated and carried forward in a uniform direction. This confideration clearly explains the reason why in the construction of a pile there must either be two metals employed, which are differently acted upon by the interposed fluid, or that the different surfaces of the same metal be subjected to the action of different finids. Dr. Van Marum conceives it impossible Dr. Van Marum that the extrication of the electric fluid can depend upon the denies this. oxidation of the metals, because he found that the pile acted more powerfully when he used the muriate of ammoniac, than when a diluted acid was employed; but the objection of the learned experimenter is fufficiently obviated by the confideration suggested above. He farther alledges in support of his particularly beopinion, that a pile with potash interposed between the plates cause potash afacted very powerfully, though no oxidation was produced. I without oxidahave repeated this experiment, and have on the contrary uni-tion;

but the contrary

proved to be the fact. formly

engagement of electricity is always attended with oxidation in a metal.

supposed to be that of Volta combined with that of oxidation.

exposed to faturated folution of muriate of ammonia caused some acid, and

became white.

in like circum-

flances was oxided and ab-

crust; in this case the water employed in dissolving the potash, oxidates the zinc in confequence of the attraction which subfifts between the oxide of zinc and the fixed alkali. We conclude it to be a general fact, that whenever a metal is oxi-Probably the dif- dated the electric fluid is difengaged, and it is probable that the reverse of this proposition will be found to be true, viz. that whenever electricity is difengaged from a metal it becomes It will appear therefore, that the experiments which were made with fo much accuracy by Sig. Volta, upon the different electricities produced in metals by mutual contact, though certainly inadequate to form a complete theory of the action of the pile, are yet not to be difregarded in our The true theory attempts to increase the power of the apparatus. In order that its confiruction be adapted for exhibiting the greatest energy, it will be necessary both that the metals employed be such as to produce a current of electricity when brought into contact, and also that the interposed fluid should rapidly oxidate one of the metals without exercifing any action upon the other. The combination of zinc and filver answers the first of these conditions, and accordingly affords the best materials for the formation of the pile. With respect to the interposed solution, the muriates, and particularly the muriate of ammoniac, have for the most part been found the most efficacious, probably in confequence of the attraction which fubfifts between zinc and the muriatic acid. The muriat ammoniac is known however to act readily upon copper as well as upon zinc, and yet we find that this falt in a ftate of folution powerfully excites the energy of a pile composed of zinc and copper, a fact which feems to militate against our hypothesis. With a view to illustrate this point the following experiments were performed: Exper. I. Zinc 1st. A piece of zinc was exposed to the action of a saturated folution of the muriate of ammoniac, a small quantity of hidrogen gas was disengaged, the fluid acquired an excess of alkali, and after fome time the metal became covered with a emission of hidrogen, absorbed white crust. 2. A clean plate of copper was placed horizontally in a folution of the muriate of ammoniac; after a few hours the upper furface of the metal was covered with a bright Exp. 2. Copper green crust, while the under surface became beautifully studded with small, transparent, and nearly colourless crystals. forbed acid, &c. These crystals after exposure to the air assumed a green hue,

fimilar

fimilar to that of the crust on the other side of the copper. This green crust was scarcely acted upon by water, but was rapidly diffolved by ammoniac, and the folution became of a deep blue colour. It feemed therefore to be a muriated oxide of copper. The crystals were only in part foluble in water, the infoluble part became brownish; it was quickly dissolved by ammoniac, and a deep blue was produced. The water in which the crystals had been digested was without colour or fmell, but upon the addition of a few drops of caustic potash. it exhaled a strong odour of ammoniac. These crystals therefore appeared to confift of muriate of ammoniac united to a quantity of the oxide, or muriate of copper. The fides of the glass were lined with a coating of the green oxide, and as the ammoniac continued to evaporate, the quantity of this oxide was increased. The fluid in which the copper had been digested exhibited alkaline properties, and was of a deep blue colour; it contained the ammoniated oxide of copper. 3. A Exper. 3. Zinc piece of zinc and a piece of copper fimilar to those employed and copper in and copper in a fiin the former experiments, were placed in a folution of the milar folution; muriate of ammoniac in contact with each other. The zinc acted on. was oxidated as in the former case, but the fluid remained colourless, and the copper acquired no green crust; though after being removed from the liquor and exposed for some time to the atmosphere, it gradually acquired a thin covering of oxide. The liquor appeared to confift of muriate of ammoniac with a fmall excess of alkali. In this experiment we may conjecture, that the zinc poffessing a stronger attraction for oxigen than the copper, abstracts it from the water, and the hidrogen which is then difengaged either prevents the oxidation of the copper from taking place, or reduces the oxide as quickly as it is formed. 4. When the zinc and copper were placed in sepa-Exper. 4. Zinc rate glasses of the solution of the muriate of ammoniac, and and copper exthe metals were connected by a filver wire, the zinc became but in separate oxidated as usual: the copper was in appearance scarcely vessels, and comacted upon, but the fluid acquired a blue tinge, which proves filver wire, the that the ammoniated oxide of copper is formed, and confe-copper was little quently that the muriate of ammoniac and water are decomposed, and the copper oxidated, though these effects take zinc and copper place to a much less extent than in the second experiment. may be confidered as if one The general refults are favourable to our hypothesis, as we of the metals

able by mur. acid.

only were oxid- learn from them that the action of the muriate of ammoniac upon copper is in a great measure suspended when the copper is in contact with zinc.

On the difference of the elect. at the ends of the pile.

3. Before we can arrive at a perfect theory of the galvanic apparatus, it is necessary to ascertain what is the essential difference between the condition of the electric fluid, as it exists in the two ends of the pile. It is established decidedly by the experiments of Van Marum, that the fluid exists in the positive state at what is called the zinc end, and in the negative at the When the materials are arranged in the opposite extremity. order of filver, zinc, and card, it is evident that the end from which the electric fluid is discharged in the positive state, is the one contiguous to the oxidating furface of the zinc. It is not altogether determined in what confifts the difference between the positive and negative electricities, but it is generally supposed to depend upon the fluid being contained in bodies in a greater or less quantity than their natural proportion. Let us examine how far this takes place in the pile of Volta. may confider each pair of metallic plates with the interpofed fluid as a complete apparatus for evolving and receiving the electric matter; and to whatever extent the pile is increased. we shall have merely a repetition of the same action. fig. 3, plate 8, represent the plate of zinc, F the interposed fluid, and S the plate of filver. At the surface of the zinc, cd, the electric matter is difengaged in consequence of the oxidation of the metal; this difengaged electricity is carried across the fluid, and enters the furface of the filver ef, and being then diffused through the body of this plate, the whole of the filver acquires the positive state, and consequently i becomes the pofitive and k the negative end of this little apparatus. plate of zinc in this case become absolutely or only relatively negative? It might be supposed that no other part of the zinc should have its electricity charged except that which is oxidated; yet from the experiments of Van Marum and others we are led to conclude, that the zinc becomes absolutely negative Dr. Van Marum informs us, that through its whole substance. when the wire of a Leyden phial was applied to the negative end of his large pile, the infide of the jar became charged with negative electricity; and as there does not appear to have been any connection between the external coating and the other end

of the pile, we must conclude that the zinc plate becomes absolutely negative through its whole fubstance. If the zinc became negative only in relation to the quantity of electricity accumulated in the filver, it would follow that the same effects should be produced whether the filver was connected with the negative end of the pile, or with any other metallic fubstance of equal bulk; but this is directly contrary to fact. Besides a Direction of the variety in the quantity or quality of the electricity, another ma-current. terial difference between the two ends of the pile is the direction of the current. While the electric matter is difengaged from the oxidating furface, it feems impossible to imagine that it can be received again by the fame furface from which it has been discharged, and which is continually discharging fresh quantities of the fluid. The electricity is therefore necessarily diffused through the interposed liquid, and from this is taken up by the opposite metallic surface. Upon the uniform direction of its current it is that the action of the apparatus essentially depends; and accordingly we find that its effects are fuspended when it is altogether furrounded with water, or even when the external edges of the plates are moistened. It is a very Limit of electricurious circumstance in the operation of this apparatus, that it zation. does not appear capable of having its power augmented by permitting the electricity to become accumulated in it. oxidating furfaces of the zinc are continually discharging a large quantity of electricity, all of which is carried forward to the positive extremity, and does not appear to escape from it; yet after the two ends of the pile have been brought to an equilibrium by means of an uniting conductor, the energy and intenfity of the apparatus are inftantaneously restored to as great a degree as before the discharge. Upon the whole it may be inferred, that though the two extremities of the pile differ from each other in the absolute quantity of electricity which they contain, yet there is reason to suppose that some farther alteration is effected, which at prefent we are unable to explain.

4. With respect to the operation which takes place at the ends Effects at the of the wires in the interrupted circuit, whatever idea we may ends of the adopt of the comparative states of the electric sluid in the two extremities of the pile, we can scarcely doubt that it proceeds from the positive to the negative end, or from the wire which emits oxigen to that which emits hidrogen. The oxigen which is disengaged from the positive or discharging wire, must neces-

farily

farily either proceed from the wire itself, or be difengaged from the fluid in which the wire terminates. If it proceed from the wire itself, we may conclude that in whatever fluid the wire is plunged, the oxigen will still be emitted, except the fluid should itself possess an attraction for oxigen. Upon the other supposition, however, that the oxigen proceeds from the decomposition of the fluid, in consequence of the electricity uniting itself to hidrogen in its passage between the wires, we may expect to be able to suspend the production of oxigen, by presenting to the wire a liquid which contains hidrogen united Immersed in am- to some other element. The liquid caustic ammoniae promises to answer all the requisite purposes, and we shall find that the experiments made upon it by Mr. Cruickshank and Mr. Davy, though not precifely fimilar in their refults, afford conclusions not unfavourable to our hypothesis. Mr. Cruickshank employed wires of platina, and when these were connected with the pile, and permitted to terminate in ammoniac, he found Azote and hidro- that the gases evolved were azote and hidrogen, with so small a mixture of oxigen as to render it probable that this latter fubstance proceeded rather from some accidental circumstance, than from the effential nature of the operation. Mr. Davy

monia.

gen evolved by platina wires in ammonia.

Gold wires;

corrotion:

used gold wires which terminated in two glasses of ammoniac, connected to each other by means of muscular fibres. found that the positive wire evolved gas only in small quantity, and that it was a mixture of three parts of oxigen and two of azote, while the other wire gave out hidrogen in confiderable quantity; the positive wire was visibly corroded. I was for fome time at a loss to reconcile the results of these able experimenters, but upon reflection I was led to conclude that the variation might be produced in part from the nature of the wires which they used, and partly from a difference in the energy of their respective piles. In order that the experiment should be unexceptionable, it is necessary that a metallic conductor be employed which possesses little attraction for any of the elements composing the fluid in which it terminates; as by this precaution any change of composition which takes place must depend altogether upon the affinities of the electric fluid. Gold is on this account less proper than platina. It may be conjectured

that in Mr. Davy's experiment with gold wires, the predifpofing affinity of the oxide of this metal for an oxigenated nitric acid, would cause the decomposition of a portion both of

the water and the ammoniac, and that the oxigen and azote thus liberated, would unite together in fuch proportion as to generate a folvent for the metal. We suppose the gas which is difengaged from the positive wire to be produced by the decomposition of the fluid in which it terminates, in consequence of the strong attraction which subsists between the electric matter and hidrogen. It may reasonably be inferred that there is a point of reciprocal faturation between these substances, or that a given quantity of electricity cannot pass between the wires unless it be united to a definite quantity of hidrogen. the electric matter in the positive wire exist there only in a fmall quantity, it acquires for itself a sufficient proportion of hidrogen by decomposing the ammoniac alone; but when the The solution of apparatus is large and operates with energy, the ammoniac is alkali decominadequate to afford the necessary supply of hidrogen, and the posed by a water becomes also decomposed. I endeavoured to submit the weaker power, and also water by truth of this conjecture to the test of experiment, for which a stronger. purpose gold wires were employed terminating in ammoniac, Experiment. and connected in different inflances with a greater or less number of metallic plates. The refults were upon the whole favourable, but on account of the minute quantities of gas procured, I was unable to ascertain its nature with as much accuracy as was defirable. When the process went on rapidly the water was decomposed as well as the ammoniac, and if it was continued for a great length of time, the ammoniac attracted carbonic acid gas from the atmosphere, and thus produced a fource of error in the refults. Several other metallic bodies which were interposed between the ends of the pile. were all of them more or less corroded in consequence of their strong attraction for oxigen. When copper and tin were used, the oxides were dissolved by the ammoniac; in the former case this was rendered visible by the blue colour which the liquor assumed, and in the latter by the precipitation of the oxide of tin effected by the addition of fulphuric acid. The flips of tin foil which were employed had their extremities changed to a brown colour, particularly that which was connected with the negative end of the apparatus.

After these observations upon the action of the positive wire, The minus wire there remains little to be added on the subject of the opposite of the pile. one; it merely receives the electric stud after its passage

through

through the water, and transmits it to the negative end of the pile. If the wire terminate in pure water, the hidrogen is difengaged in the form of gas; but if a metallic falt be diffolved in the water, the hidrogen is attracted by the oxide, and this last reduced to the state of a metal. The phenomena exhibited in this process are often peculiarly beautiful, and the extreme delicacy which the apparatus possesses in discovering even the minutest metallic impregnations, suggests its probable utility in the analytical operations of chemistry.

How far galvanour electric theories.

5. A very interesting subject of inquiry here presents itself, ism may modify viz. in what degree will our theoretical notions of the action and properties of the electric fluid be influenced by the galvanic discoveries? The theory of Dr. Franklin, as well as the elegant one proposed by M. AEpinus, seems to be adapted only to those cases in which the electric fluid previously existing in bodies, was altered either with respect to its disposition or quantity. But we are now in possession of an apparatus by which it is actually generated, a circumstance with which we were until lately unacquainted, and confequently our hypothefis, being a general deduction from facts, will at least require to be extended, perhaps in some instances new-modelled.

Chemical theory nefited by these discoveries.

The pile of Volta affords many illustrations of the modern is also much be-chemical doctrines, which appear the more striking as effected by experiments made with a different object in view, and produced by an agent which cannot be supposed capable of affecting the refults. The grand question respecting the decomposition of water, if it flood in need of farther support, derives powerful confirmation from the experiments first performed by Mr. Nicholfon and Mr. Carlifle, and fince repeated with various modifications. The experiments made with the galvanic apparatus also confirm our ideas respecting the composition of ammoniae and nitric acid, and confiderably add to our knowledge of the action which takes place between acids and metals. They beautifully exhibit the powerful effects produced by oxigen and hidrogen when in a condenfed or uncombined state, and promife to afford the chemist a valuable opportunity of. employing in his different operations these bodies in a state of absolute purity. Every one interested in the progress of philofophical knowledge, will regard with admiration the discovery of an apparatus, which has enabled him to develope fome of the

the most hidden operations of nature, and will consider the name of Volta as worthy to be classed with those of Lavoisier, Priestley, Cavendish, and the other illustrious sounders of the pneumatic chemistry.

#### III.

A Summary of the most useful Parts of Hydraulics, chiefty extracted and abridged from Eytelwein's Handbuch der Mechanik und der Hydraulik. Berlin, 1801. By Thomas Young, M. D. F. R. S.

(Concluded from Page 35.)

CHAPTER 9. Of the motion of water in pipes. Motion of water

The author has attempted to simplify this subject nearly in in pipes. the same manner as that of the motion of rivers, and apparently with considerable success. He observes, that the head of water may be divided into two parts, one of which is employed in producing velocity, the other in overcoming the friction: that the height employed in overcoming the friction must be as the length of the pipe directly; and also directly as the circumference of the section, or as the diameter of the pipe, and inversely as the content of the section, or as the square of the diameter; that is, on the whole, inversely as the diameter; this height too must vary, like the friction, as the square of the velocity.

Hence  $f = \frac{al}{d} \cdot v^2$ , f denoting the height due to the friction, To find the velocity; and a a constant quantity: therefore,  $v^2 = \frac{fd}{d}$ . Now the

v2 ==

height employed on the friction, corresponds to the difference between the actual velocity and the actual height, or  $f=h-\frac{v^2}{b^2}$ , where b is the coefficient for determining the velocity from the height; consequently,  $v^2=\frac{b^2dh-dv^2}{ab^2l}$ , and  $v^2=\frac{b^2dh}{ab^2l+d}$ . Now b=6.6, and from Buat's experiments,  $ab^2$  is determined to be .0211, which agrees the most accurately where the velocity is between 6 and 24 inches in a second. Whence we have

$$v^2 = \frac{43.6dh}{.0211l+d}$$
, or  $v=45.5$   $\sqrt{\left(\frac{dh}{l+47d}\right)}$ ; but it is fomewhat more accurate to make  $v=50$   $\sqrt{\left(\frac{dh}{l+50d}\right)}$ ; all the

measures being expressed in English feet.

allowance for flexure. When the pipe is bent into angles, or rather arcs, we must diminish the velocity thus found, by taking the product of its square multiplied by the sum of the sines of the several angles of insection, and then by .0038; which will give the degree of pressure employed in overcoming the resistance occasioned by the angles: and deducting this height from the height corresponding to the velocity, we may thence find the corrected velocity.

Compound pipes.

Mr. Eytelwein proceeds to investigate, both theoretically and experimentally, the discharge of water by compound pipes, with apertures of various dimensions between them: he allows at each orifice for the contraction of the stream, and calculates the height necessary to produce the increase of velocity in each instance, allowing also for the friction of the pipe. But the velocity thus found is somewhat smaller than the result of his experiments; probably because the whole of the force of the water accelerated at any orifice, is not immediately lost as soon as it arrives at a wider part of the pipe. The ascent of water in a compound pipe, to the level of a reservoir, is next considered, a case which often occurs in pump-work; and an approximation to the velocity of ascent is deduced from theory and compared with experiment.

lets of water.

Chapter 10. Of jets of water.

This chapter contains little that is new or interesting; it is well known, that the velocity of a jet is greatest when it springs through an orifice in a thin plate, and in this case, the height falls little short of that of the reservoir.

Impulse or hydraulic pressure of water.

Chapter 11. Of the impulse or hydraulic pressure of water. There are three principal cases of the impulse of water falling perpendicularly on plane surfaces: when a detached jet of water strikes the plane; when the plane moves in an unlimited extent of water, or is very small in respect to a stream that strikes it; and when the impulse takes place in a limited channel.

When the ftreum Supposing a stream of water to strike against a plane so as to loses all its velo-lose all its motion, it is obvious, that the force that destroys city;

the motion must be equal to the force that generates it; that is, to the weight of the column of water operating during the time necessary for its acquiring the given velocity: and the quantity of water arriving during this time, being equal to twice the column of which the length is the height due to the velocity, the hydraulic pressure must be twice the weight of such a column. The relative impulse against a plane in mo-or when the tion, must be determined from the difference of the velocities: plane recedes but when all the water of a stream strikes against a plane, the effect of the impulse may be more simply determined, as if a solid body struck the plane with the relative velocity; and this is nearly what happens in undershot water-wheels.

When a detached jet strikes against a plane, it appears, Detached jet, from the experiments of Bossiut and Langsdorf, that its effect is equal to the weight of an equal column of twice the height due to the velocity; but the plane must be at least four times as large in diameter as the jet; if it be only of the same size, the effect will be but one half as great. In an unlimited Unlimited stream, the impulse is also nearly determined from the height corresponding to the velocity; and it appears, that the effect is nearly doubled by confining the stream to prevent its divert-confined in a ing laterally from the float-boards.

For oblique surfaces, the effect of a detached jet in its own Oblique surfaces direction, appears to vary as the square of the sine of the angle opposed to a jet of incidence; but, for motions in open water, we must add to this square about  $\frac{2}{3}$  of the difference of the sine from the radius:

a correction which is tolerably accurate, until the inclination becomes very great. Mr. Eytelwein found the resistance to the motion of a sphere nearly  $\frac{4}{3}$  of the resistance to a circle equal to its section: perhaps it was a hemisphere, otherwise it is difficult to reconcile the result with other experiments in which it has appeared to be only  $\frac{2}{3}$ .

Mr. Eytelwein informs us, that at the temperature 14° of Weight of diftile Reaumur, or 63½ of Fahrenheit, a cubic foot of diftilled water led water weighs 66.0656 pounds of Cologne, or 65.9368 commercial pounds of Berlin. According to Sir George Shuckburgh's experiment, an English cubic foot of distilled water at 66° weighs 997 ounces avoirdupois; and water expands for every degree .000165: hence the pound of Cologne is 1.0312 Eng-

lish avoirdupois pounds, and that of Berlin 1.0332.

Chapter 12. Of overshot water-wheels.

The power which operates upon overshot wheels, is divided Vol. III.—October, 1802. G into

Overshot wheels;

into two parts, one derived from the weight of the water in the cells or buckets, the other from the impulse of the water falling on it: the effect of the first is constant, that of the second varies with the velocity: the maximum is found to be when the velocity is half that of the water received; but the variable part being the smaller, the rule is of little practical consequence, and the velocity of the wheel is generally greater than this. advantage of re- The author observes, that by turning the stream back upon the nearer half of the wheel, we remove the refiftance of the lower water, fince it runs off in the fame direction with that of the

verfing their rotation.

water-wheel.

Undershot

wheels.

Chapter 13. Of undershot water-wheels.

Floatboards should rife perpendicularly.

The author enters into a minute description of the parts of an undershot water-wheel: he observes, that the most advantageous position for the floatboards in a straight channel is, when they are perpendicular to the water at the time that they rife out of it: that only one half of each should ever be below the furface, and that from three to five should be immersed at once, according to the magnitude of the wheel. When there is fufficient fall, the floatboards should be divided and made into buckets, fo that the wheel may become a breast wheel; the position of the external portion being such, that a line drawn through it at the time when the water enters, may divide the vertical radius in the same proportion that it divides the quadrant of the circumference; that is, if the water is received, for instance, at one third of the quadrant from the bottom, the line must leave one third of the radius above it. A formula is laid down for calculating the actual force of a given stream of Undershot wheel water on a wheel, and it is shown, that half the velocity of the ftream is that which gives the maximum of effect, the theory

ought to work with half the velocity of the Aream.

agreeing perfectly with the experiments of Smeaton and others: for, fince the effect is estimated by the product of the force into the velocity of the parts upon which it acts, and fince the force is in this case simply as the relative velocity, because the quantity of water is given, and the whole of it is supposed in all cases to act: therefore, the effect will be expressed by the product of the relative and absolute velocity of the wheel, or e=va; but r=v-a, v being the velocity of the fiream, and ra=av-aa, which is obviously greatest when v=2a, as is evident either by taking the fluxion, or by confidering that the greatest ordinate of a femicircle is the radius.

To show the advantage of breast wheels over common un- Breast wheels dershot wheels, the author quotes Mr. Banks's experiments, much better than undershot. He also observes, that by placing two wheels after each other in the same stream, about one fourth more force may be obtained than either by a fingle wheel, or by two wheels fide by fide; but that a fingle wheel has less friction, and is generally less expensive.

Chapter 14. Of the properties of the air, as far as they relate to hydraulic machines.

What Mr. Eytelwein quotes as Mariotte's discovery of the Force of air increase of the air's density in proportion to the pressure, was against a plane; well known to Hooke and Boyle. From the experiments of Woltmann and Schober, he remarks, that the force of the wind against a perpendicular plane, is nearly equal to four thirds of the weight of a column of air, of a length equal to the height due to the velocity. The height of a column of water nearly equivalent to the force or refistance may be found, by taking Rule for finding the square of 1 of the velocity in a second, in English feet.

Thus, if the velocity were 1000 feet in a fecond, the refif-Example. tance would be equal to a column of water in the same surface, 25 feet in height; and the refistance to a sphere about half as much.

For another example, if we had a cubic foot of a fubfiance Instance of the equal in specific gravity to water, and were defirous of know-utmost velocity ing the greatest velocity that it could acquire by falling through by a body falling the air; the height of the column of water is here 1, and its in the air: fquare root 1, which multiplied by 200 gives 200 feet in a fecond for the velocity, when the refistance would be equal to the weight, which of course is the limit beyond which the velocity could never pass. Hence we may form an idea of the utmost velocity that a stone, of moderate size, could acquire in descending from the upper regions of the atmosphere, or even from the neighbourhood of the moon; a velocity that would be much much less than less than that of a bomb or a cannon ball, even when it may be ball. followed by the eye.

Again, Mr. Garnerin's parachute contains about 860 fquare Parachute of feet of furface, and weighs, together with the aeronaut fuf-Garnerin. pended from it, about 230 pounds. Here the weight is \$\frac{23}{86}\$ of a pound for each square foot, which is equivalent to  $\frac{1}{230}$  of a foot of water; multiplying the square root by 200, we have about 13 feet in a fecond for the utmost velocity; which is the

fame

fame as if one leaped from a height of between two and three feet. Mr. Garnerin, however, finds the mean velocity of defeent only eight feet, which agrees better with the experiments of Borda, in which the refistance appeared to be  $\frac{5}{3}$  of the weight of the column due to the velocity, and exceeded this proportion as the furface increased in magnitude.

Syphons.

Chapter 15. Of fiphons.

For estimating the discharge of a siphon, the head of water must be reckoned equal to the difference between the levels of the surface of the water, and of the lower orisice. The author observes, that the theory of waves has been treated in a new and improved manner by Lagrange in his Mécanique Analytique. The problem is, however, not yet completely solved: Lagrange's formula includes the depth of the water agitated as a given quantity, but it does not inform us how to determine this depth from theory.

Sucking pumps.

Chapter 16. Of fucking pumps.

The length of a fucking pump must never be greater than 30 feet below the moveable valve: and there may be a loss of time in the ascent of the water, unless it be made even a few feet shorter. The motions to be produced, and the resistances to be overcome, are considered in detail: but the author refers, for still further information, to Langsdorf's Treatise on Machinery.

Nature of the Broke.

The velocity of the stroke should never be less than four inches, nor greater than two or three seet in a second; the stroke should be as long as possible, to prevent loss of water by the frequent alternations of the valves. The diameter of the pipe should be about  $\frac{2}{3}$  or  $\frac{3}{4}$  of that of the barrel. The listing pump is also here described; it only differs from the sucking pump in having the lower valve moveable, and the upper one sixed. A number of valves and pistons are described in this chapter; chiefly from models of English manufactory.

Forcing pumps;

Chapter 17. Of forcing pumps.

In describing the different kinds of solid piston, the author gives the preference to that which has a conical leather projecting on each side; but remarks, that there is another form which has the advantage in the situation of the ring for receiving the rod, which is precisely in the centre of the piston, and is therefore sitter for communicating motion in each direction. He says, that where the barrel is well polished, the piston may be used without either wadding or leather.

made without leathering.

The

The first pump, invented above a century before Christ, by The first pump Ctefibius of Alexandria, to whom also music is indebted for was a forcer the organ, and whose name Mr. Eytelwein mentions in speaking of fucking pumps, was in reality a forcing pump, as may eafily be collected from its description by Vitruvius (L. X. cap. 12.)

Chapter 18. Of mixed pumps, or the combination of suck-Mixed pumps.

ing and forcing pumps.

When the lower valve is above the furface of the water, the forcing pump can only raise the water by suction, but the construction remains the same. Such is Mr. Buchanan's patent ship pump. De la Hire's pump is more complicated; both the afcending and descending strokes of the piston being made effective, by means of a double apparatus of valves and pipes.

Chapter 19. Of acting columns of water.

lumns acting by

The mechanism of a pump may be employed for convert-preffure. ing the weight of water descending in its barrel, to the purpose of working another pump. The author describes a machine of this kind invented by Mr. Höll, and improved by Langsdorf. A similar arrangement, used in Cornwall, has lately been described in Nicholson's Journal, by Mr. Trevithick. The only objection to it appears to be the magnitude of the friction.

Chapter 20. Of the spiral pump.

The spiral pump or horizontal he-

If we wind a pipe round a cylinder, of which the axis is lix forcing water horizontal, and connect one end with a vertical tube, while up a pipe. the other is at liberty to turn round and receive water and air in each revolution, the machine is called a spiral pump. It was invented about 1746, by Andrew Wirz, a pewterer in Zurich, and was employed at Florence with Bernoulli's improvement, in 1779. At Archangelsky, near Moscow, a pump of this kind was erected in 1784, which raifed a hogshead of water in a minute, to a height of 74 feet, and through a pipe 760 feet in length. The force employed is not mentioned, we may therefore conjecture that it was turned by water. Mr. Eytelwein enters very minutely into calculations of the effect of fuch a machine under different circumstances; and the results of theory, as well as of experiment, are fuch, as to induce us to expect that it will in time come into common use, instead of forcing pumps of a more complicated and expensive construction. The water-tight joint prefents the only difficulty: the

pipe may form either a cylindrical, a conical, or a plane fpiral, and it appears to be uncertain which is the most advantageous: the vertical pipe should be nearly of the same dimensions as the spiral pipe, which may without difficulty be made of wood.

Screw of Archimedes.

Chapter 21. Of the fcrew of Archimedes, or the waterfnail, and of the waterfcrew.

The fcrew of Archimedes confifts either of a pipe wound

spirally round a cylinder, or of one or more spiral excavations,

formed by means of spiral projections from an internal cylinder, covered by an external coating so as to be water-tight. But if the coating is detached, so as to remain at rest while the spirals revolve, the machine is called a waterscrew. Mr. Eytelwein enters into an accurate determination of the effects of these machines in given circumstances, and the results of the theory agree remarkably well with experiment. He observes, that the screw of Archimedes should always be so placed, as to sill exactly one half of a convolution in each turn; and that very unfavourable reports have sometimes been made of the machine from want of attention to this circumstance; for when the orifice remains constantly immersed, the effect is very much

diminished: this appears also to have happened in some late experiments in London. Where the height of the water is so variable as to render this precaution impossible, Mr. Eytelwein presers the waterscrew; although, in this instrument, one third of the water generally runs back, and it is easily clogged by accidental impurities in the water. The screw of Archimedes is generally placed so as to form an angle of between 45° and 60° with the horizon; but the open waterscrew at an angle of 30° only: for great heights, the spiral pump is preserable to

The waterfcrew.

Caution for placing the forew of Archimedes.

Bucket wheels.

Throwing wheel.

Chapter 22. Of bucket wheels and throwing wheels.

In the conftruction of wheels for raifing water in buckets, there is little room for refined theory; whether the buckets be fixed or suspended on an axis. It is sometimes convenient to raise water to the height of 3 or 4 feet by the revolution of a wheel with simple sloatboards; and such a wheel may be either in a vertical or an inclined position; it must of course be inclosed in a sweep.

Chapter 23. Of cellular pumps and paternoster works.

either.

Water has been fometimes raised by stuffed cushions con-Cellular pumps nected with an endless rope, and caused by means of two or paternoster wheels or drums to rise in succession in the same barrel, and to carry water with them: but the magnitude of the friction ap- The chain-pears to be an objection. From the resemblance of the appa-pump. ratus to a string of beads, it has been called a paternoster work. When stat boards are united by chains, and employed instead of these cushions, the machine may, without impropriety, be called a cellular pump: here the barrel is generally square, and placed in an inclined position. But these machines are very rarely employed.

Chapter 24. Of instruments for measuring the velocity of

ftreams of water.

The superficial velocity of a stream is ascertained without Instruments for difficulty, by observing in calm weather the motion of a body measuring the velocity of barely floating on it. But it is more difficult to determine the ftreams of water. velocity of a river at a confiderable depth. Pitot's tube, as improved by Buat, furnishes one of the easiest methods. A funnel is prefented to the stream, and the water in a vertical tube connected with it, is elevated above the level of the river, nearly to the height corresponding to the velocity: but the result will be more accurate, if the funnel be covered by a plate perforated only in the centre by a small orifice: in this case, the elevation in the tube will be half as great again as the height due to the velocity. Other inftruments for appreciating the impulse of the water against a flat board, require some previous comparative observation. In Woltmann's hydrometrical fly, the number of revolutions of a wheel, in a given time, indicates the velocity of the water, which strikes against two inclined planes, and carries round the arm to which they are fixed.

It is prefumed, that this abridged account of Mr. Eytel-Concluding adwein's book, will not only do justice to his diligence and ingenuity, but will convey to the English reader some matter perfectly new, and capable of frequent application in practical hydraulies; which is perhaps of the more value, as there is little probability that the work will be translated at length. To disseminate information of this kind must always be the principal object of the Journals of the Royal Institution.

7

#### IV.

Experiments and Observations on certain stony and metalline Substances, which at different Times are said to have fallen on the Earth; also on various Kinds of native Iron. By Edward Howard, Esq. F. R. S. From the Philosophical Transactions, 1802.

# (Continued from Page 263, Vol. II.)

Chemical examination of stones fallen on the earth. I PROCEED to consider the assistance to be derived from chemistry in distinguishing these stones from all other known substances, and in establishing the assertion, that they have fallen on the earth.

Objections to the analysis of the stone prefented by the Abbé Bachelay, The analysis made by the French academicians of the stone presented to them by the Abbé Bachelay, was in part conducted by the ever to be deplored Lavoisier; but it was performed before that celebrated author had enriched chemistry with his last discoveries, and before he had given birth to the system under which it slourishes. The result of this analysis might well induce the conclusion, that the subject of it was common pyritical matter. It was unfortunately made of an aggregate portion of the stone, and not of each distinct substance irregularly disseminated through it. The proportions obtained were, consequently, as accidental as the arrangement of every substance in the mass.

and to that of M. Barthold.

The analysis of M. Barthold, of the stone of Ensisheim, is subject to the same objections: but, after having the advantage of the foregoing descriptions, the researches which follow cannot be supposed altogether liable to a similar fatality.

# EXAMINATION OF THE STONE FROM BENARES.

Chemical examination of the stone from Bepares. This stone, as the Count de Bournon has already remarked, has the most distinguished characters. Indeed it is the only one of the sour sufficiently perfect (if I be allowed that expression) to be subjected to any thing approaching to a regular analysis.

The black coating deprived of magnetic parts

The crust, or external black covering, is the first substance to which the attention is naturally directed. When a portion

of

of this crust had been detached with a knife or a file, and finely was digested in pulverized, I separated the particles attractable by a magnet, nitric acid; and digested the unattractable portion with nitric acid, which which it decomwas prefently decomposed; but, owing to a strong adherence posed. of some of the interior and earthy parts of the stone, it did not difentangle the coating or metalline part without some difficulty. The acid being fufficiently neutralized, the folution was passed through a filter, and saturated to excess with ammonia. An abundant precipitate of oxide of iron was produced; and, The iron foluwhen this oxide was separated, I observed the saline liquor to tion was precipitated by amhave a greenish colour. I evaporated it to dryness, and re-monia. diffolved the dry falt in diffilled water. No precipitate was formed during the evaporation, nor was the colour of the folution entirely destroyed. It appeared to me like a triple Ammoniacal nifalt, described by Mr. Hermstadt \* as an ammoniacal ni-trate of nicke trate of nickel. By examination with prussiate of ammonia, it yielded a whitish precipitate, inclining to a violet colour; and, by various properties, I was foon confirmed in the opinion, that nickel was prefent. Since I shall have occasion more than once to treat of the triple compound, and fince it has been only mentioned by Mr. Hermstadt, it is necessary now to detail some of its distinctive characters. The same chemist informs us, that the three mineral Distinctive chaacids, with ammonia, enter into fimilar combinations with triple falt. nickel; and I have observed, that oxide of nickel can be diffolved by nitrate and muriate of ammonia. The muriate feems to take up the largest quantity. The colour of this falt is by no means uniform: it is fometimes grafs green, violet, rose colour, inclining to purple, and I have seen it almost colourless. It seems to be purple, and to incline to rose colour and violet, when all the oxide of nickel is not united to both acid and alkali, but, from the deficiency of falt, is held in folution by an excess of ammonia. In this cafe, evaporation, of courfe, precipitates the nickel in the state of oxide, which is of a whitish green colour.

The nickel cannot be precipitated from a perfectly formed The nickelis triple falt, by any reagent I have tried, except by a pruffiate, precipitable by a pruffiate or an or a hidrogenized fulphuret of ammonia. Potash and lime, as hidrogenized

\* Annales de Chimie, Tom. XXII. p. 108.

fulphuret of ammonia.

well as, I presume, other bodies, standing in the order of affinities before ammonia, decompose the falt; but the nickel is then continued in folution by the difengaged ammonia.

The liquid was always examined if it had been prefent.

As it may be imagined that I have occasionally met with to detect copper copper, when I describe a violet or purple ammoniacal solution, it is right to observe, that to avoid this error. I have either reduced the liquor to a neutral state, and endeavoured. without fuccefs, to obtain from it a precipitate, with a folution of fulphureted hidrogen gas; or, by adding an acid to flight excefs, and immerfing a piece of iron, I have not been able to detect a trace of copper. These, and many other trials. when they do not appear to be made before the estimation of the quantities of nickel, have been constantly made after-

The coating contained iron nearly metallic, and fome nickel.

But, to return to the incrustation or coating of the stone. the decomposition of the nitric acid shewed the presence of matter at least nearly metallic, although not attractable; and the examinations made of the liquor, from which the iron was precipitated, ascertained the presence of nickel beyond disrute. The difficulty of obtaining the coating of the stone, either distinct from matter not belonging to it, or in sufficient quantity, induced me to relinquish the idea of attempting to give the proportions of its constituent parts.

The fhining particles of the fione

The stone being deprived of its covering, the shining particles irregularly diffeminated, next demand examination. I first examined the pyrites. Their very loose texture made it exceedingly difficult to collect the weight of 16 grains, which was however effected by the dexterity of the Count de Bournon.

were digested in dilute muriatic acid.

It diffolved the metal, and fufpended the fulphur, and the earthy portion fubfided.

I digested these, at a low heat, with weak muriatic acid; which acted gradually, and difengaged a trifling but fenfible quantity of fulphureted hidrogen gas. After feveral hours, I found the acid discontinued its action. The whole metalline part appeared in folution; but fulphur and earthy particles were observable. The sulphur, from its small specific gravity, was suspended through the solution; whilst the earthy matter, which could not be separated by mechanical means, was fortunately left at the bottom of the digesting vessel. I decanted off the folution; holding suspended the sulphur; and, by repeated washing, separated every thing belonging to the py-

rites

rites from the infoluble earthy matter, the fubtraction of which reduced the weight of real pyrites to 14 grains. I next obtained the fulphur, by filtration. When it was as dry as I could make it, without fear of its being sublimed, its weight was two grains. To the filtrated liquor I added nitrate of barytes, by way of detecting any fulphuric acid which might have been present; but no cloudiness ensued. I then separated, by fulphate of ammonia, the barytes thus added, and precipitated the iron with ammonia. The liquor, on the fubfidence The metals were of oxide of iron, appeared of a violet purple colour: it con-iron and fome tained nickel, which I threw down with fulphureted hidrogen gas, there being already a sufficient excess of ammonia in the faline liquor to form an alkaline hidrogenized fulphuret. The oxide of iron, after ignition, weighed 15 grains; and the fulphuret of nickel, reduced to an oxide, weighed, after the fame treatment, fomething more than one grain. The proportions of the substances contained in the pyrites of the stone from Benares, may therefore be confidered nearly thus.

Sulphur

Iron
Since 15 grains of the oxide represent about that quantity of iron,
Nickel, nearly

Extraneous earthy matter

Grains.

Component parts, sulphur, much iron, nickel, and earth.

Ad Memory, Arrive of a resp. "Quality for a resp. "15

It is observable that, notwithstanding the loss appears to be Remarks. only half a grain, it was probably more, because the sulphur could not be reduced to the same state of dryness in which it existed when in combination with the iron; not to say that it was, in a small degree, volatilized with the hidrogen gas dis-

engaged during the folution.

The weight of nickel is a mere estimation. We are not yet sufficiently acquainted with that metal to speak of it with accuracy, except as to its presence. Upon the whole, however, it may be concluded, that these pyrites are of a very particular nature; for, although Henkel has observed that sulphur may be separated from pyrites by muriatic acid, it is by no means the usual habitude of pyrites to be of such easy decomposition.

The

ing particles being malleable iron.

The other shin- The other shining particles immediately seen, when the internal structure of the stone is exposed, are the malleable iron. Before I state the examination of this iron, I must remark, that preliminary experiments having shewn me it contained nickel, I treated feveral kinds of the most pure irons I could obtain, with nitric acid; and precipitated the oxide from the metallic falt by ammonia. The quantity of oxide I obtained from 100 grains of iron, was from 144 to 146. I may confe quently infer, that 100 grains of pure iron acquires, by fuch a process, 45 grains of oxigen; and that, whenever a metallic substance, supposed to be iron, does not, under the same circumstances, acquire the same proportionate weight, something is either volatilized, or left in folution. Hence, when a metallic alloy of nickel and iron prefents itself, a judgment may, at least, be formed of the quantity of nickel, by the deficiency of weight in the precipitated oxide of iron.

were distolved in nitric acid, and precipitated by ammonia,

This mode of treatment was not allowed me in the examination of the coating of the stone, because it was impossible to know in what state of oxidizement the iron existed. But, as the particles disseminated through the whole mass, are clearly metallic, a very tolerable idea of the quantities of nickel contained in them will be obtained, by noting the quantity of oxide of iron feparated, as above described. 25 grains of these metallic particles were therefore heated with a quantity of nitric acid, much more than sufficient to dissolve the whole. Some earthy matter, which, as in a former case, was not separable by mechanical means, remained after a complete folution of the metal had been effected. This earthy matter, after being ignited, weighed two grains. The real matter of the present examination, was therefore reduced to 23 grains, and was in complete folution. I added ammonia to a very The oxide of iron was thereby precipitated, fensible excess. and, being collected and ignited, it weighed 24 grains; whereas, according to my experiments, 33½ grains should have been produced from the folution, had it contained nothing but iron. which left nickel I examined the faline liquor, when free from ferruginous par-

of which the timated.

in triple folution ticles, and discovered it to be the triple falt of nickel. Hence, quantity was ef-allowing for loss, the quantity of nickel may be estimated, by calculating the quantity of iron contained in 24 grains of oxide. Thus, if 145 grains of oxide contain 100 of iron, about 161 are contained in 24 of oxide. This would suppose the 23

grain

grains of alloy to confift of  $16\frac{1}{2}$  iron and  $6\frac{1}{2}$  nickel; which, if the usual loss be added to the 161 grains of iron, and deducted from the nickel, may not be very remote from the truth.

I shall next examine the globular bodies, also irregularly The globular bodispersed throughout the stone. A number of them were reduced to sine powder; but nothing metallic could be separation, exhibited nothing metallic could be separation. rated by the magnet. As a preliminary experiment, I fought tic. for pyrites, by digestion with muriatic acid; but no hepatic fmell was in the least perceivable, nor was white carbonate of lead at all altered by being held over the mixture. I therefore conclude these globular bodies do not envelope either iron or pyrites. By way of analysis, I treated 100 grains with pot-Analysis. ash, in a filver crucible; and, after the usual application of a red heat, feparated as much filica as poslible, by muriatic acid and evaporation. The filica being collected on a filtre, carbonate of potash was added to the filtrated liquor; by which a precipitate, almost wholly ferruginous, was produced. This precipitate was collected in the common way; then boiled with potash, to extract alumina; and, by superfaturating the alkaline liquor with muriatic acid, and precipitating by carbonate of ammonia, an earth was gathered, which I afterwards found to be partly, if not intirely, filiceous. After redissolving, in muriatic acid, the portion of the ferruginous matter rejected by the potath, I precipitated by ammonia, what I took to be intirely oxide of iron; but, after igniting it, and again attempting to rediffolve the whole in muriatic acid, more filica was left. The non-existence of lime was proved, by the addition of carbonate of ammonia, immediately after the same alkali, pure, had thrown down what I took wholly for oxide of iron. I had now obtained every thing in the fubject of my analysis, except magnesia and nickel. The former, and a trace of the latter, were held by carbonic acid in the liquor, from which the ferruginous precipitate was, in the first instance, thrown down by carbonate of potath; and the latter

Under all circumstances, I am induced to state the propor- Constituent tions of conflituent parts thus;

was found in the last named muriate of ammonia. I disengaged the magnefia, by the affiftance of potash, and by evaporating to dryness. The oxide of nickel was precipitated by hidroge-

nized fulphuret of ammonia.

111

Silica

#### ON STONY AND METALLINE SUBSTANCES

Silica	-			11 =	1/1	Hita	1 10 50
Magnefia			-	od)			ud ile15
Oxide of iro	n			154		. 3	34
Oxide of ni	cke			syla			5. 120d 21
				3172	- 13	11	Samuel La

1012

#### Remarks.

The excess of weight, instead of the usual loss, is owing to the difference of oxidizement of the iron, in the stone and in the refult of the analysis; which will be found to be the case in all analyses of these substances; indeed it is always necesfary to reduce the oxide to the red state, as being the only one to be depended upon. To avoid future repetition, I shall also observe, first, that by preliminary experiments, I could not detect any other substance than those mentioned. Secondly, that the earth obtained as alumina, appeared to me to be mostly, if not intirely, filiceous; because, after it had been ignited, and again treated with potash and muriatic acid, I found it was very nearly all precipitated by evaporation. Thirdly, I examined, and judged of, the filica collected from the oxide of iron, in the same way. Fourthly, the weight of the magnetia is given, not immediately, as obtained by evaporation, but after a subsequent solution in an acid, and precipitation by potash. And, fifthly, the proportions are taken from the mean of two analyses.

The earthy matrix or cement.

Nothing remains to be examined, of the stone from Benares, except the earthy matter, forming a cement or matrix for the substances already examined. 100 grains of this matter were, by mechanical means, separated as perfectly as possible, from the pyrites, iron, and globular bodies, and analysed as above. The mean result of two analyses gave,

Its component	1	Silica -		-	4 1177	-	48
parts,		Magnefia	-	-	17-1400		18
		Oxide of iron		0.		-	34
	712 7	Oxide of nickel		-	****	-	21
	1						T RALLEN

1021

Stone from Sienna.

# EXAMINATION OF THE STONE FROM SIENNA.

The external coating of this stone appeared to have the same characters as that of the stone from Benares.

The

The pyrites, although certainly prefent, were not crystallized in fuch groups as in the preceding stone; nor could they be separated by mechanical means.

The attractable metal was easily separated by the magnet; Iron and nickel but  $8\frac{1}{2}$  grains only were collected. I treated them with nitric as in the former acid and ammonia, as in a preceding case. Nearly one grain the attractable of earthy matter was insoluble; the weight was therefore re- metal. duced to rather less than 8 grains. The oxide of iron, precipitated by ammonia, weighed 8 grains; and the faline liquor gave abundant indications of nickel. As 8 grains of this oxide of iron contain nearly 6 of metal, the quantity of nickel, in the bare 8 grains, may be estimated between 1 and 2 grains.

Some globular bodies were extracted, but too few to analyze.

Since the pyrites could not be feparated, I collected 150 Analysis of the grains of the ftone, freed from iron by the magnet, and as ex-ftone freed from empt as possible from globular bodies. These 150 grains, I and globular first digested with muriatic acid, that the pyrites might be de-Parts. composed, and every thing taken up which could be dissolved by that menstruum. A very decided disengagement of sulphureted hidrogen gas was occasioned. When the acid could produce no further action. I collected the undiffolved matter on a filtre, and boiled it with the most concentrate nitric acid. in hopes of being able to convert the fulphur, previously liberated, into fulphuric acid; but my endeavours were fruitless; for, upon the addition of nitrate of barytes to the nitric folution, rendered previously transparent, a very infignificant quantity of sulphate of barytes was obtained. The surplus of barytic nitrate was removed by fulphate of potash. I next completely edulcorated the mass which remained infoluble, after the action of the muriatic and nitric acids; and, adding the water of edulcoration to the muriatic and nitric liquors, evaporated the whole for filica. I then submitted the mass. undiffolved by the acids and the water, to the treatment with potash, muriatic acid, and evaporation, which was, in the first instance, applied to the stone from Benares. The first precipitation was, as in that analysis, also effected with carbonate of potath; but, instead of endeavouring immediately to extract alumina, I ignited the precipitate, that the alumina or filica remaining might be rendered infoluble. After the ignition, I feparated the oxide of iron with very concentrate muriatic acid; and the earths, which were left perfectly white, I heated

I heated with potash, until they were again capable of being taken up by the same acid. The solution so made, was slowly evaporated; and, as very nearly every thing was deposited during the evaporation, I conclude all was silica. The proportions resulting from this single analysis, without the weight of sulphur contained in the pyrites irregularly disseminated through the whole, were,

Its component

Silica -		10.70	70
Magnefia -	-	12 0	34
Oxide of iron		10 I	- 52
Oxide of nickel	-	4	- 3
			-
			159

# The stone from EXAMINATION OF THE STONE FROM YORK-Yorkshire. SHIRE.

The mechanical feparation of the fubstances in this stone being as difficult as in the preceding case, I was necessarily satisfied with submitting it to the same treatment. I collected, however, 34 grains of malleable particles; which, by the process already more than once mentioned, left 4 grains of earthy matter; and, by yielding  $37\frac{1}{2}$  of oxide of iron, indicated about 4 grains of nickel.

Component parts.

150 grains of the earthy part of the stone were, by analysis, resolved into.

Silica -	-	* (1)			75	
Magnefia -	4	0 2		olas	37	10
Oxide of iron	-		1		48	
Oxide of nickel	. 1	10.120		6	2	
				111		

162.

The stone from

# EXAMINATION OF THE STONE FROM BOHEMIA.

The probability of never being able to obtain another specimen of the very remarkable fragment of this substance, did not allow me to trespass more on the liberality of Mr. Greville, than to detach a small portion. I sound it of similar composition to that of the three preceding stones; and the Count de Bournon has already shewn the proportionate quantity of the attractable metal to be very considerable. 16½ grains, lest 2½

of extraneous earthy matter; and yielded, by the treatment with nitric acid and ammonia,  $17\frac{1}{2}$  grains of oxide of iron. This would feem to induce an estimation of 11 of nickel in 14 grains, or about 9 per cent.

55 grains of the earthy part of the stone, by the analytical Component treatment of the two former, afforded, hand the parts.

Silica	-		0,01	Toy-	2019		1.63	25
Magnefia	nt-	411			1- /		40.7	91
Oxide of iro			<b>-</b> (£ ¹)				1 ,:	23
Oxide of nic	kel	Cell		Set 1	1-100		- 1711	1 1
main T. Leads	10 Art		. 17	1115.	. Fall Di	Sec.		

with Salaya large to most of the space of the soft sell

The unufual increase of weight in the result of the three last analyses, notwithstanding the intire loss of the sulphur in the pyrites, is obviously owing to the metallic state of the iron combined with the fulphur, as was shewn in a former instance.

Is have now concluded the chemical examination of these Remarks on four extraordinary fubstances. It unfortunately differs from these analyses the analysis made by the French Academicians, of the stone those formerly presented to them by the Abbé Bachelay. as well as from that made by others. made by Professor Barthold, of the stone of Ensisheim. It is at variance with that of the Academicians, inafmuch as they found neither magnefia nor nickel. It differs from that of Mr. Barthold, as he did not find nickel, but discovered some lime, with 17 per cent. of alumina. With regard to these differences, I have to submit to the chemical world, whether magnefia might not have eluded the action of an acid, when the aggregation of the integrant parts of the stone was not destroyed by treatment with potash. As to the existence of alumina, I do not absolutely deny it; yet I must observe, that the whole of the earth which feemed to have any refemblance, however small, to alumina, was at most 3 per cent. and there feems good reason to consider it as silica. Respecting the exiftence of lime in the stone of Ensisheim, I must appeal to Professor Barthold, whether, supposing lime a constituent part, fulphate of lime flould not have been formed, as well as fulphate of magnesia, when sulphuric acid was generated by igniting the carths and pyrites. And, as to the proportion of alumina, in the same stone, I would ask, at least, whether it would have been so considerable, if the solutions formed by acids, after the treatment with potasi, had been evaporated VOL. III.-OCTOBER, 1802,

The ftriking conformity of character in these stones, and femblance to all other minerals, join with the historical testimony to prove their origin.

to the requisite dryness: not to observe that no mention is tuade of any examination of the properties of the earth called alumina. In the proportion of magnefia, I have the fatisfaction to find my analysis correspond very nearly with that of Profesior Barthold; and, if what he considered alumina were supposed filica, the stone presented to the French Academy, the stone of Ensisheim, and the four I have examined, would agree very nearly in filiceous proportions. With respect to the nickel, I am confident it would have been found in all, had the metallic particles been feparately examined. their want of re- whatever be these variations, the mineralogical description of the French Academicians, of Mr. Barthold, and of the Count de Bournon, all exhibit a striking conformity of character, common to each of these stones; and I doubt not but the fimilarity of component parts, especially of the malleable alloy, together with the near approach of the constituent proportions of the earths contained in each of the four stones, the immediate subject of this Paper, will establish very strong evidence in favour of the affertion, that they have fallen on our globe. They have been found at places very remote from each other. and at periods also sufficiently distant. The mineralogists who have examined them, agree that they have no refemblance to mineral fubstances, properly so called; nor have they been described by mineralogical authors. I would further urge the authenticity of accounts of fallen stones, and the similarity of circumstances attendant on such phenomena; but, to the impartial it would be superfluous, and, to those who disbelieve whatever they cannot explain, it would be fruitless. Attempts to reconcile occurrences of this nature with known principles of philosophy, it is true, are already abundant; but (as the Earl of Bristol has well expressed) they leave us a choice of difficulties equally perplexing. It is however remarkable, that Dr. Chladni, who feems to have indulged in these speculations with most success, should have connected the descent of fallen frones with meteors; and that, in the narrative of Mr. Williams, the defcent of the stones near Benares, should have been immediately accompanied with a meteor.

Luminous appearance attending the fall of feveral.

No luminous appearance having been perceived during the day on which the stone fell in Yorkshire, it must be admitted, rather militates against the idea, that these stones are the subflances which produce or convey the light of a meteor, or that

a meteor must necessarily accompany them \*. Yet the stone from Sienna fell amidft what was imagined lightning, but what might in reality have been a meteor. Stones were also found after the meteor feen in Gascony, in July, 1790. And Mr. Falconet, in the memoir I have already quoted, relates, that the stone which was adored as the mother of the gods, was a Bœtilia; and that it fell at the feet of the poet Pindar, enveloped in a ball of fire. He also observes, that all the Bostilia had the fame origin.

I ought not perhaps to suppress, that in endeavouring to Electricity renform an artificial black coating on the interior furface of one ders these stones of the stones from Benares, by sending over it the electrical black. charge of about 37 square feet of glass, it was observed to become luminous, in the dark, for nearly a quarter of an hour; and that the tract of the electrical fluid was rendered black. I by no means wish to lay any stress upon this circumstance; for I am well aware, that many substances become luminous by electricity.

But, should it ever be discovered that fallen stones are actually the bodies of meteors, it would not appear fo problematical, that fuch masses as these stones are sometimes reprefented, do not penetrate further into the earth: for meteors move more in a horizontal than in a perpendicular direction; and we are as absolutely unacquainted with the force which impels the meteor, as with the origin of the fallen stone.

Before I close this subject, I may be particularly expected Meteor which to notice the meteor which, a few months ago, traverfed the lately croffed the county of Suffells. It was first that a first the county of Suffells. county of Suffolk. It was faid, that part of it fell near Saint folk. Edmundsbury, and even that it set fire to a cottage in that vicinity. It appeared, from inquiries made on the fpot, that fomething, feemingly from the meteor, was, with a degree of reason, believed to have fallen in the adjacent meadows; but the time of the combustion of the house did not correspond with the moment of the meteor's transition.

A phenomenon much more worthy of attention, has fince been described in the Philosophical Magazine. On the night Prodigious meof the 5th of April, 1800, a body wholly luminous, was feen, teor in Americae in America, to move with prodigious velocity. Its apparent

<sup>\*</sup> In the account of the stone which fell in Portugal, no mention is made, either of a meteor or lightning.

fize was that of a large house, 70 feet long; and its elevation above the furface of the earth, about 200 yards. The light produced effects little fhort of fun-beams; and a confiderable degree of heat was felt by those who saw it, but no electric fensation. Immediately after it disappeared in the north-west, a violent rushing noise was heard, as if the phenomenon were bearing down the forest before it; and, in a few seconds after, there was a tremendous crash, causing a very sensible earth-Search being afterwards made in the place where the burning body fell, every vegetable was found burnt, or greatly fcorched, and a confiderable portion of the surface of the earth broken up. We have to lament, that the authors of this account did not fearch deeper than the furface of the ground. Such an immense body, though moving in a horizontal direction, could not but be buried to a confiderable depth. Should it have been more than the femblance of a body of a peculiar nature, the lapfe of ages may perhaps effect what has now been neglected; and its magnitude and folitary fituation become the aftonishment of future philosophers.

Concerning the native iron of S. America.

This leads me to speak of the solitary mass of what has been immense mass of called native iron, which was discovered in South America, and has been described by Don Rubin de Celis. Its weight was about 15 tons. The same author mentions another infulated mass of the same nature. The whole account is exceedingly interesting; but, being already published in the Philofophical Transactions for the year 1788, it needs not be here repeated.

> Mr. Prouft has shewn the mass particularly described, not to be wholly iron, but a mixture of nickel and iron. tees of the British Museum, who are in possession of some fragments of this mass, sent to the Royal Society by Don Rubin de Celis, have done me the honour to permit me to examine them; and I have great fatisfaction in agreeing with a chemist fo justly celebrated as Mr. Proust.

and that deand confidered by the Tartars as fallen from keaven.

The connection which naturally exists between one mass of scribed by Pallas, native iron and another, immediately turns our attention to the native iron in Siberia, described by Pallas; and this, we are told, the Tartars confidered as a facred relic, which had dropped from heaven. The nickel found in the one mass, and the traditional history of the other, not to compare the globular bodies of the from Benares with the globular concavities

concavities and the earthy matter of the Siberian iron, tend to the formation of a chain between fallen stones and all kinds of native iron. How far any real affinity exists between these several substances, very obliging friends have afforded me an opportunity to form some judgment. I am indebted to Mr. Greville and Mr. Hatchett for portions of almost every known native iron: and the Count de Bournon has done me the savour particularly to describe them as follows.

(To be concluded in our next.)

#### v.

On the Colours obtained from Metallic Oxides, and fixed by Means of Fusion on different Vitreous Bodies. By ALEX. BRONGNIART, Director of the National Manufactory of Porcelain at Sévres, Engineer of Mines\*, &c.

THE art of employing metallic oxides to colour the different The colouring vitreous matters has been long known. It is well known that of glaffes by oxthe ancients made coloured glafs and enamels, and that this cient. art was much practifed by the Egyptians, who were the first that imitated precious stones by these means.

In modern times the practice of this art has been brought to The theory negating high degree of perfection, but its theory has been neglected, lected, It is almost the only chemical art to which the new principles

of this science have not been applied.

A CONTRACTOR

di di -

In the numerous works which treat of the method of using or erroneous, and preparing metallic vitrisiable colours, the authors either give no theory, and consequently no general principles, or the explanations are founded only on the absurd hypotheses which formerly composed much of the theory of chemistry.

One of the best publications, as it is the work of an en-Montamy's lightened practician, is The Treatife on Painting in Enamel, of treatife com. Montamy. The archives of the national manufactory at Sévres, mended. likewise contain simple and good processes for the fabrication of colours; Bailly, Fontelliau, and Montigny are the authors; but they are simple descriptions, without any observations that lead to general principles.

<sup>\*</sup> Journal des Mines, No . 67.

Kunckel's work and the Encyclopædias are of little value.

Kunckel's work, the manuscripts of Hellot, in the possession of the manusactory at Sevres, and the two Encyclopædias, present only an undigested collection, and a multitude of processes collected together from all quarters, without choice or reason. With a slight knowledge of the art it is easier to invent a new process of sabrication, than to discover which among this multitude of receipts we ought to preser.

Science is confiderably advanced when the facts compose a body of doctrine.

It has been remarked, that one of the most certain criterions of the progress made to the persection of a science, is the possibility it leaves of a re-union of the sacts that compose it, in a body of doctrine from which general principles may be deduced. At the present period only does it deserve the name of a science, and it is to the exposition of these principles, that the striking, though inaccurate name of the philosophy of science has been given.

The arts would be greatly advanced by fimilar treatment. The arts, which oftener compose a branch of science, than the simple application of one of its parts, present facts equally capable of being united in a body of doctrine; they will attain this valuable degree of persection if practised by men accustomed to trace the connections, and deduce the consequences of the events that pass under their observation.

Philosophers might acquire them. Philosophers, whose more elevated speculations seclude them from the practice of these arts, would then perceive their principles with greater ease; they could more directly apply their researches to the progress of the art, which being then directed by reason, would become more certain, direct and rapid in its advancement.

The prefent memoir aims to make this difpofition as to the art of vitrifiable colours.

Being convinced that the art of preparing and using vitrifiable colours is susceptible of receiving the application of the means of persection; and that the sacts of which it is composed are sufficiently numerous and accurate to be presented in a general outline, I have concluded that the precise knowledge of these facts, and of the principles which unite them, which naturally lead to an explanation of a number of the results, might be interesting to chemists, who, being occupied by more general and important researches, cannot attend to all the details of a complicated art.

I wished likewise to give an accurate account of the principles of this art to chemists, that they might determine with certainty on the new operations to which the processes submitted to their judgment might lead.

Lastly,

Lastly, I thought it would be advantageous to the progress Exhibition of of the art, and that it was incumbent on the national manufact colours at Sevres. tory at Sévres to disclose the pretended secret of the compofition of colours for porcelain which do not change in the fire. These colours were presented to the Institute in the year 6, by a manufacturer of porcelain, justly admired for the beauty of the works produced at his manufactory. I should not have published this secret had it been given to me in confidence, but as it is known, I hope I shall no longer be suspected of the least breach of integrity.

From what has been faid, it will be feen that my object is not to give the exact composition of all the vitrisable colours at full length. Such a work could not be the subject of a sim-

ple memoir, 100 an what is a set with throw would to see

It is known that metallic oxides are the bales of all vitrifia- Vitrifiable coble colours; but some of the metallic oxides are not proper for by met. oxides. this use; and as they are not vitrifiable without some admix-

ture, they can feldom be employed alone.

The very volatile oxides, and those that do not adhere Not those which firongly to the oxigen they contain in abundance, either can are volatile, or eafily change not be employed in any manner, as is the case with the oxides their oxigenaof mercury and arfenic, or elfe they can be employed only as tion. agents. The colour they afford cannot be depended on, be-

the flightest heat. Such are the puce and red coloured oxides

of lead, the yellow oxide of gold, &c. Those oxides, in which the proportion of oxigen is capable of being varied too easily, are seldom employed. The black oxide of iron is never employed for that colour. The green oxide of copper is in many circumstances very uncertain in its

canfe they lofe it, in proportion as they lofe their oxigen by

effect.

I have remarked that the oxides do not melt alone; though Oxides cannot when placed in thin layers on vitrifiable fubflances they can be used alone; be made to adhere by means of a violent fire; but, with the exception of the oxides of lead and bifmuth, they afford only dull colours. The very violent heat which is frequently required to fix them, changes or totally destroys their colours. Some flux is therefore added to all the metallic oxides.

This flux is either glass of lead and filex, or glass of borax, but require a or a mixture of both.

flux. Lead and filex, or lead, Its filex, and borax.

The flux fixes the colour at a moderate heat; less than would destroy the colour.

Its general effect is to give brilliancy to the colours after their fusion, to fix them upon the piece that is painted by foftening its furface, to envelope the metallic oxides, to preferve their colour by defending them from the contact of the air, and particularly to facilitate the fusion of the colour at a flightly elevated temperature, not capable of destroying it.

The observation which proves this latter use of solvents is taken from delicate colours, fuch as the carmines produced from gold; these colours require a much greater proportion of flux than the others.

colours are flux without previous fusion.

Very transient Sometimes the oxides are directly employed, and fimply mixed with the mixed with their folvents, without having been previously melted with it; such are the colours which the violence or repetition of heat would alter. It is easy to conceive that a stronger and more lasting heat is required to suse a crucible of coloured glass than a layer of colour, which is not the tenth of a millimetre in thickness. this white that he there are a

I shall return to this subject when treating of the red colours obtained from gold. The state of the state o

Others are fuled ground.

In many cases the oxides are previously sused with their solwith it and then vents, and afterwards ground. When I fpeak of colours in particular, I shall mention those which are subjected to this fusion.

These general principles are too simple to require further explanation.

This treatise is confined to vitrifiable colours on glazes or glafs.

I shall only speak of the application of metallic colours to vitreous bodies or furfaces.

These grounds are three.

These bodies may be divided into three very distinct classes, from the nature of the substances that compose them, the effects produced on them by the colours, and the changes they undergo. These are,

r. Enamel.

1st. Enamel; fost porcelain, and all the glazes, enamels, or glasses, which contain lead in any considerable quantity.

2. Hard porcelain. 3. Common

2d. Hard porcelain, or fuch as is glazed with feldfpar. 3d. Glass in which there is no lead, such as the common

window glass. glass.

I shall successively examine the principles of composition of these colours, and the general phenomena they present on these three grounds or supporters.

Colours

\*niColours for enamel painting have been longest known: the Enamel colours receipts in the works I have mentioned at the beginning of have been long

this essay all relate to these colours.

It is known that enamel is a glass rendered opake by oxide Enamel is glass of tin, and very fusible by the oxide of lead. It is this last rendered suffile which in particular gives it properties very different from opake by ox. tin. those of the other excipients of metallic colours. Hence all the glasses and glazes that contain lead, have the properties of enamel, and what we may affert of the one, will apply to the other with very little difference.

Such are the white and transparent glazes of Dutch or Delft It is the glaze of ware, and the glaze of the porcelain called foft (tendre.) Delft ware, &c.

This porcelain, the first made in France, particularly at Soft porcelain of Sévres, and indeed for a long time almost exclusively at this Sévres. manufactory, has for its base a vitreous frit nearly opake, capable of being acted upon by marl, and its glaze is a very transparent glass containing much lead.

The colours made use of are the same as those for enamel. Colours the ling, confequently the changes these colours undergo in ena-fame as for ena-melling. mel, must take place in this species of porcelain, because, as we shall foon see, the causes of the change are the same in

bothelt to their are a lines The colours for enamel and tender porcelain require less These require flux than the others, because the glass on which they are less flux because the glaze is very placed, foftens fufficiently to be penetrated by them.

This folvent may be either the mixture of glass of lead, and The flux before pure filex called rocaille, or this fame glass mixed with that mentioned.

Montamy fays that glass of lead ought not to be used in the flux for enamel; he employs borax alone. He then dilutes or makes up his colours in a volatile oil.

On the contrary, the painters of the manufactory at Sévres, Glass of lead is use only colours without borax, because they dilute them with a good material. gum, and borax does not dilute well this way. I am convinced that both methods are equally good, and that Montamy is not justified in excluding the fluxes of lead, as they are employed without inconvenience every day, and even render the management of colours more easy.

I have remarked, that in the baking of these colours the The soft glaze glaze is foftened fo much as to be easily penetrated by them; very much dithis is one great cause of the change they undergo. They be-lours.

come diluted by the mixture with the glaze, and the first fire changes a painting apparently finished, into a very slight fketch.

And the oxide mically on them.

The oxide of lead contained in the glaze is a more powerful of lead acts che-cause of the great changes these colours undergo. Its destructive action is principally exercised on the reds of iron, and is very remarkable. I shall relate some experiments that particularly prove this.

Thefe colours themselves are fcarcely changeable, except they are placed upon.

It has already been shewn that the two principal causes of the change which colours on enamel and tender porcelain undergo, do not relate to the composition of these colours, but from the ground intirely to the nature of the glass upon which they are placed. The affertion that the colours of porcelain are subject to considerable change, relates to the colours of foft porcelain, (porcelain tendre) a species of ware now almost totally abandoned.

The paintings touching up.

It follows from what I have faid, that the paintings of before mention-ed require much porcelain require to be feveral times re-touched and burned, in order to possess the necessary strength. Though these paintings have always a certain foftness, they are constantly more brilliant, and never subject to the inconvenience of scaling off.

Hard porcelain; with glaze of feld fpar,

The hard porcelain, according to the division I have laid down, is the fecond species of ground or excipient for the metallic colours. It is known that the base of this porcelain is a very white argil, called kuolin, mixed with a filiceous and calcareous folvent, and the glaze of which is nothing but feld fpar fused without an atom of lead.

or of Saxony, receives two kinds of colour; or painting colours,

This porcelain, which is that of Saxony, is of much later date at Sévres than the foft or tender. The colours employed viz. on the glaze are of two kinds, the first used for representing different objects, are baked with a very inferior fire to that required for the baking of the porcelain itself. They are very numerous and varied.

or beneath it; or ground colours.

The others, which require to be fused at as great a heat as that for baking the porcelain, are laid on the general furface. They are much less numerous.

The colours for painting are nearly the same as for the foft porcelain; but contain more Aux.

The colours for painting are made up very nearly of the fame materials as those for tender porcelain; they only con-This flux is composed of the glass of lead, tain more flux. I have not yet met with any (called rocaille) and of borax. work that treats of the composition, use and effects of these colours. In fact, it has no where been afferted in print, that all these colours except one are unchangeable in the fire; The flux is glass whereas it has been often afferted in books, that paintings in of lead and boen amel are subject to considerable change.

When the porcelain is put in the fire to bake the colours, the The hard glaze feld spar glaze dilates and opens its pores, but does not become does not run nor soft. As the colours do not penetrate it, they are not subject lours, to the changes they undergo on tender porcelain. It must however be observed, that they lose a little of their intensity, by acquiring the transparence given them by the susion.

When works of little importance are made they need not so that only fine be re-touched; but this is necessary when a painting is to be work needs rebighly finished. This re-touching is not distinguishable in the paintings on porcelain from that of any other species of painting.

One of the great inconveniences of these colours, is the faci- The colours are lity with which they scale or sly off when the fire is often ap- apt to scale.

plied.

This has been particularly remarked at Sévres, on account Hard porcelain is of the folidity and infufibility with which porcelain is there and franch manufactured. But these qualities cause it to resist the alter-changes of heat nations of heat and cold for a longer time, and gives its ground and cold better than the soft.

a more brilliant white colour. On the other hand, the porcelains of Paris being more vitreous, transparent, and of a blue-ish cast, generally crack if boiling water is frequently poured into them.

In order to remedy this evil without altering the quality of The scaling is the body, I softened the glaze a little, by introducing more prevented by softening the solution or calcareous flux according to the nature of the feld glaze. This method succeeded, and for this twelvemonth past the colours have passed two and three times through the fire without cracking, provided there be not too much flux, and they be not laid on too thick.

It has been remarked, that when foda and potash were in-Alkalies are too troduced the colours scaled; so that they cannot be used as volatile to be used in this art. It was. These alkalies being volatilized abandon the colours, which cannot adhere to the glaze by themselves.

I have observed that other colours are likewise prepared, Colours to be which being laid upon the general surface, are sused by the laid on and baked by the same fire as bakes the porcelain. These colours are but sew; first fire are sew. because there are sew metallic oxides that can support such a

fire without being volatilized or discoloured. Their solvent is the feld fpar. As they incorporate with the glaze they never crack, and are more brilliant, whitesa on with the base

The third receptacle of metallic vitrifiable colours is glass I for glass chates and open its peres, but do bal with white the se-

on glass,

Art of painting The application of these colours constitutes the art of painting on glass; an art much practifed in former ages, but which is supposed to be lost, because out of fashion. It however too immediately depends on the art of painting on enamel and porcelain to be intirely loft. Descriptions of the processes may also be found in many different works. fill tral : bade set a paid

Books in which it is treated. Original.

is not loft.

A book intitled, L'Origine de l'Art de la Peinture fur Verre. published at Paris in the year 1693, and Le Traité de l'Art de la Verrerie, by Neri and Kunckel, feem to be the first works containing complete descriptions of this art. Those published

only a compilation.

Leviel's work is fince, even the great work of Leviel, which constitutes part of Les Arts et Metiers of the academy, and of the Encyclopédie Methodique, are only compilations from the two former works.

The others afford very loofe instruction.

It is fomewhat remarkable, that if we follow the processes exactly as they are described in these works, as I have done with some of them, the colours of which they pretend to give the receipt would never be fabricated. They only ferve to shew an able practician the method, and leave it to him to correct or make additions. This was found to be the case by Citizen Meraud, who was engaged to prepare them for the manufactory of Sévres. He was obliged to make the colours for painting on glass rather from his own experience, than from the instructions in the works I have just mentioned.

The colours for glass are nearly the fame as for enamel; but dull for want of the opake ground.

The limits of a memoir will not permit me to enter into hiftorical details on the art of painting on glass; its history is given at length in Leviel's work; the materials and fluxes some of them are which enter into the composition of the colours for painting on glass, are in general the same as those applied to porcelain. They vary only in their proportions; but a great number of the colours used for enamel and porcelain cannot be applied to glass; many of them when seen by transmitted light intirely change their aspect, and exhibit an obscure tint which can be of no use when deprived of the white ground which throws them out. We shall point out these when we treat of the colours in particular. Those colours which can be used on

this

this body, fometimes change in the baking, and acquire a great transparence. They are generally beautiful only when placed between the eye and the light, and then they answer the pur-

pose intended in painting on glass.

There is more difficulty in baking plates of coloured glass Difficulties of than is commonly thought. The bending of the piece, and baking glafs alteration of the colours, are to be avoided. All the treatifes we have consulted recommend the use of gypsum. This me- On the usual thod fometimes fucceeded, but generally the glass became fupport of gyp-fum, the alkali white, and cracked in all directions. It appears that the glasses of the glass is atthat are too alkaline, and which are far the most common in tacked, and the clear white glasses, are attacked by the hot sulphuric acid of piece broken. the fulphate of lime. We were able with ease to bake much Porcelain biscuit larger glaffes than any before painted, by placing them on very is a much better fupport. smooth plates of earth or unglazed porcelain.

Concerning the feveral particular Colours.

After having collected the general phenomena which each Particular coclass of vitrifiable colours offers with regard to the bodies on lours. which they are placed, I must shew the particular and most interesting phenomena which every principal species of colours employed on tender porcelain, on glass, and in the fire that bakes the porcelain, prefent.

Concerning the Reds, Purples, and Violets obtained from Gold.

The carmine red is obtained from the purple precipitate of Carmine red Caffius. It is mixed with about fix parts of its flux, and this from gold, mixture is directly employed without being first fused. It is then of a dirty violet, but acquires the beautiful carmine by baking. It is however very delicate; a little too much heat is very perishor carbonated vapours easily spoil it; yet it is more beautiful able in the fire, when baked with charcoal than with wood.

This colour and the purple which differs little from it, as as well as its well as all the shades which are obtained from their mixture with shades. other colours, really change in all porcelains and in the hands of all operators. But this is the only one that changes on hard It is the only co-porcelain. It may be replaced by a substitution of rose colour on hard porcefrom iron, which does not change; fo that by excluding from lain; the pallet the carmine made from gold, and substituting the

and may be changed for oxide of iron in this use, though this oxide fails on enamel.

rofe-coloured oxide of iron here spoken of, we have a pallet composed of colours none of which are subject to any remarkable change. This rose-coloured oxide of iron has been long known, but was not employed on enamel, because it is then subject to considerable change. Or perhaps when the painters on enamel became painters on porcelain, they continued to work according to their ancient method.

We might suppose that by previously reducing the colour named carmine, already mixed with its folvent, into a vitreous matter, the last tint would be obtained. But as I have already proved, the fire which must be used to melt this vitreous mass destroys the red colour. Besides, it is sound that, to obtain this colour in perfection, it is necessary to pass it through

the fire as little as possible.

Fabrication of the gold carmine.

The carmine of tender porcelain is made with fulminating gold gently decomposed, and muriate of filver; there is no tin in it, which proves that it is not necessary for the fabrication of a purple colour, that the oxide of this last metal and that of gold should be combined.

Violet from gold stands better on foft porcelain.

Violet is likewise obtained from the purple oxide of gold. This colour proceeds from having a greater quantity of lead in the flux, and it is nearly of the fame tint whether crude or baked.

But they all fail on hard.

These three colours totally disappear in the strong fire proper to bake porcelain.

The carmine and beautiful.

Carmine and purple afforded us upon glass, only tints of a purple are not good upon glass; dirty violet. The violet, on the contrary, has a very beautiful but the violet is effect; but it is subject to change to a blue. I cannot yet afcertain the cause of so singular a change, as I observed it for the first time only a few days ago.

(To be concluded.)

## the said him VI. of Ser Rectification

Examination of the supposed igneous Origin of the Rocks of the Trupp-formation. By Mr. Robert Jameson. From the Author.

Sheriff Brae, Leith, Sept. 8, 1802.

That the Rocks of the Trapp-formation have been deposited from a Aqueous origin State of Solution and Suspension in Water, is a Truth, which has trapp-formation been demonstrated by the Wernerian Geognosie.

At fection 64 of his illustrations of the Huttonian theory, he Prof. Playfair's

IT is not at present my intention to state the proofs of this opinion: the subject of the following observations is the examination of Mr. Playsair's account of these rocks.

observes, "The columnar structure sometimes found in that the columnar species of whinstone called basaltes, is a fact which has given form of basalt is rise to much discussion; and it must be confessed, that though because it is seen one of the most striking and peculiar characters of this fossil, is in lavas. not that which gives the clearest and directest information concerning its origin. One circumstance, however, very much in favour that basaltic rocks owe their origin to fire, is, that the columnar form is fometimes affumed by lava actually erupted from volcanos. Now, it is certainly of no small importance to have the fynthetic argument on our fide, and to know that bafaltic columns can be produced by fire; though no doubt, to give absolute certainty to our conclusion, it would be necessary to shew, that there are in nature no other means but this by which these columns can be formed. This fort of evidence can hardly be looked for; but fince the power of fusion to produce the phenomena in question is perfectly established, and fince the production of the same phenomena in the humid way is a mere hypothesis, if there be the least reason to suspect the action of subterraneous heat \* as one of the causes of mineralization,

The phenomena of volcanos and hot springs are considered by Professor Playsair as proofs of a subterranean heat, unconnected with the decomposition of any mineral substance. The Wernerian geognosie, however, has shewn us, that volcanos and hot springs originate in the newest slotz trapp-formation, therefore we have suffi-

cient

Reply. That the descriptions lanzani and others, are inappearance and aqueous origin of bafalt is deducible from fiture in other foffils to formed.

ralization, every maxim of found philosophy requires that the basaltic structure, in all cases, should be ascribed to it." The of lavas by Spal- accounts given by Spallanzani and other travellers of columnar lava, are most evidently incorrect; the appearances they have correct as to the described are accidental rents, such as we observe in fandstone. limestone, and even in basalt. Many of the prismatic lavas the matter itself; described by Spallanzani, are either basalt, greenstone, or porphyry. That bafalt columns have been formed by chrystallization from a state of solution in water, is rendered probable from milarity of ftruc- the analogy of their structure with that of other fossils which have been certainly fo formed. Bafalt we know occurs in the fame mass in tabular and columnar distinct concretions, corresponding in this particular with the lamellar and columnar diftinct concretions of truly chrystalline fossils \*.

P. The presence of pyrites urged as the proof of tion;

At fection 66 we are informed, "That a mark of fusion, or at least of the operation of heat, which whinstone possesses in igneous produc- common with many other fossils, is its being penetrated with pyrites, a substance, as has been already remarked, that is of all others most exclusively the production of fire."

but thefe are very abundant among matters never subjected to fire.

Mr. Playfair probably did not know that the great beds of bituminated wood which are found covered with fand, clay, &c. in the alluvial hills, are very frequently much impregnated with pyrites; and that lately Mr. Hatchett had also demonstrated its formation in the humid way.

P. Insulated maffes of indurated fandstone in whin, urged as a proof of igacous fusion.

He continues, " Another mark of fusion, more distinctive of whin, is, that both in veins and masses it sometimes includes pieces of fandstone, or of the other contiguous strata, completely infulated, and having the appearance of fragments of rock floating in a fluid sufficiently dense and ponderous to suftain their weight. Though these fragments have been too refractory to be reduced into fusion themselves, they have not remained intirely unchanged; but are, in general, extremely indurated, in comparison of the rock from which they appear

cient proof of the existence of quantities of inflammable matter, equal to the support of all the volcanoes that have, or can exist on our globe.

\* When basalt, having a columnar structure, occurs in veins, we find that these columns are perpendicular to the walls of the vein. This phenomena I have frequently observed, and consider it as supporting in fome degree the opinion mentioned above.

to have been detached." That the pieces of fandstone are not It is denied that insulated, I have proved in a former paper. The occurrence lated or induof masses of fandstone in veins of basalt, grunstone, and wacken, rated. is well known; but that there are masses more indurated than the other fandstone of this formation, must be denied, as we know that fandstone beds occur in this formation of all degrees of induration.

Section 67. " Similar instances of extraordinary induration P. Extraordiare observed in the parts of the strata in contact with whinstone, of the strata in whether they form the fides of the veins, or the floors and contact with roofs of the masses into which the whinstone is distributed. whinstone; The strata, whether fandy or argillaceous, in such fituations are usually extremely hard and consolidated; the former in particular lofe their granulated texture, and are fometimes converted into perfect jasper. This interesting remark was first made by Dr. Hutton, and the truth of it has been verified by a great number of subsequent observations."

The flinty fandstone which is characteristic for the newest accounted for by flötz trapp-formation, is usually covered by clay, wacken, ba-the deposition of basaltic folution falt, or grunftone, in the order here mentioned: fometimes the in the pores of clay is wanting, when the fandstone is covered by wacken, or fofter strata. when the wacken has been carried away, or not deposited, by basalt or grunstone. We have thus, to use the Huttonian language, the fandstone equally indurated under a bed of clay as under one of basalt or grunstone: it is therefore plain, that no argument can be drawn from the situation and nature of this fandstone in favour of the igneous fystem. When the walls of veins are indurated, we can easily trace it to the deposition of a portion of the basaltic solution in the pores of the softer strata.

Section 68. "To the same excellent geologist (Dr. Hutton) P. The appearwe are indebted for the knowledge of an analogous fact attend- ance of charring or coaking of coal ant on the passage of whinstone veins through coal strata. As in the vicinity of the beds of stone where they are in contact with the veins of veins of whin, adduced as a whin, seem to acquire additional induration, so those of coal, in proof of their like circumstances, are frequently found to have lost their fusi-former flate of bility, and to be reduced nearly to the state of coke or char-fusion. coal. The existence of coal of this kind has been already mentioned, and has been confidered as a proof of the operation of fubterraneous heat. In the inflances here referred to, that is, where the charring of the coal is limited to those parts of the firata which are in contact with the whin, or in its immediate VOL. III.—OCTOBER, 1802.

vicinity, the heat is pointed out as refiding in the vein; and this is to be accounted for only on the supposition of the melted whin, at a period subsequent to the confolidation of the coal, having flowed through the openings of the firata. heat has been powerful enough in many places to drive off the bituminous matter of the coal, and to force it into the colder and more diffant parts. Few facts in the history of fossils are more remarkable than this, and none more directly affimilates the operations of the mineral regions with those that take place at the furface of the earth."

In reply. to the percolation of water.

This. An eminent Irish geognost is of opinion, that the appeareffect is ascribed ances described by Professor Playsair may have been produced by the percolation of water along the fides of the basalt vein. This explanation will be fatisfactory to those who have witnessed the effects produced by the weather on coal. This altered coal, however, must not be confounded with the glanz and columnar coal, because these differ from it in oryctognostical and geognostical characters.

> The argument contained in fection 69, is refuted by what Werner mentions in his admirable book on the formation of veins.

Section 71. "Again, if it be true that the masses of whin-

P. That the whinstone was introduced berated.

ftone, thus interposed among the ftrata, were introduced there tween the strata, after the formation of the latter, we might expect to find, at is deducible from least in many instances, that the beds on which the whinstone the similarity of the parts so sepa- rests, and those by which it is covered, are exactly alike. these beds were once contiguous, and have been only heaved up and separated by the irruption of a fluid mass of subterranean lava, their identity should still be recognised. Now this is precifely what is observed; it is known to hold in a vast number of instances, and it is strikingly exemplified in the rock of Salisbury Crag near Edinburgh." If this argument was correct, it should follow, that the beds of foliated limestone containing shells, which are found between strata of fandstone,

had been ejected from below. This fact cannot be evaded

by faying, that the limestone has been only softened; it is plain if the foliated fracture (according to the Huttonian fystem) is a

This argument affirmed to prove too much, because it would apply to other middle strata.

character of fusion in one stone, so must it in all others. P. The nodules At fection 73 we have the following observations on the forof agate in whin-mation of agates: "Some of the species of whinstone are the stone appear to baye been fluid; common matrices of agates and chalcedonies, which lie inclosed

in them in the form of round nodules. The original fluidity of these nodules is evinced by their figured and sometimes chrystallized firucture, and indeed is fo generally admitted, that the only question respecting them is, whether this sluidity was the effect of heat or folution. To answer this question Dr. Hutton observes, that the formation of the concentric coats of which the agate is usually composed, has evidently proceeded from the circumference toward the center, the exterior coats always impressing the interior, but never the reverse. The same thing also follows from this other fact, that where there is any vacuity within the agate, it is usually at the center, and there too are found the regular chryftals, when any fuch have been formed. It therefore appears certain, that the progress and to have conof consolidation has been from the circumference inwards, and without: that the outward coats of the agate were the first to acquire folidity and hardness. Now it must be considered that these coats are highly confolidated; that they are of very pure fili- They are pure ceous matter, and are utterly impervious to every substance filiceous matter, which we know of, excepting light and heat. It is plain there-any other subfore, that whatever at any time, during the process of consoli-france; dation, was contained within the coats already formed, must have remained there as long as the agate was intire, without the least possibility of escape. But nothing is found within the coats of the agate fave its own fubftance; therefore no extraneous fubstance, that is to fay no solvent, was ever included within them. The fluidity of the agate was therefore fimple, whence this fluidity of the agate was therefore fimple, whence this fluidity of the agate was therefore fimple, whence this fluidity of the agate was therefore fimple, whence this fluidity of the agate was therefore fimple, whence this fluidity of the agate was therefore fimple, whence this fluidity of the agate was the t and unaffifted by any menstruum. In this argument nothing idity is inferred appears to me wanting that is necessary to the persection of a simple, without physical, I had almost faid of a mathematical demonstration. any menstruum. It feems, indeed, to be impossible that the igneous origin of fossils could be recorded in plainer language, than the phenomenon which has been just described."

It is well known to geognosts, that during the deposition of Reply. The certain kinds of strata, considerable quantities of air have been cellular structure of the strata formed, and this in endeavouring to escape has given them a ascribed to gas; cellular structure. The almond-stone (mandelstein) in which agates are most frequently found, has acquired its cellular structure in this manner. These cells appear to have been the cells are fucceffively filled with a stony solution, out of which thin coats stated to have been fucceffively of jasper, carnelian, chalcedony, &c. have been precipitated, filled with a according as the precipitate altered or was finer. We are na-ftony folution;

turally led to the idea of an increasing fineness in the solution, because we observe most distinctly that the outer coat is composed of the coarsest precipitate, and that as it approaches the middle, it is always finer and finer; fo that at last, out of the finest solution, amethyst shoots, and, when there is room enough,

dules took place.

whence the pre-chrystallizes. That the solution out of which the precipitation cipitation of no-took place was truly chemical, is demonstrated not only by the character of the substances, but also by the disposition of the coats; for we can observe that they have followed the attraction of the walls of the cell, and are fo accurately deposited upon it, that an inequality, or elevation upon the outer wall, continues to be observed through all the superincumbent coats to the center of the agate. That the folution had come from without, and by infiltration penetrated into the cells, is most diffinctly to be feen in specimens of agate when they are properly cut. If the cutting has been judiciously done, we can always observe the opening by which the solution has entered. This elegant explanation is an outline of that delivered by the illustrious Werner in his geognostical lectures, and is a demonstration of the aqueous origin of agate.

The opening of infiltration obfervable.

Flinty fossils not impervious to water.

That flinty fossils, as stated by Mr. Playfair, are impervious to water, has never been proved; on the contrary, it is well known to collectors of fossils, that if the cellular rock chrystal, which contains water, is kept in a cabinet for some years, the water gradually disappears.

Other obf. :

Sections 249, 250, 1, 2, are occupied with an unfuccessful attempt to reconcile the appearances on the hill of Scheibenberg with the Huttonian theory. Without infifting upon the Wernerian doctrine of formations, which gives the death-blow to all partial explanations like those proposed by the Huttonian theory, I shall merely mention a few facts respecting the individual relation of these rocks, to shew the insufficiency of the explanation attempted by Professor Playfair.

Situations of wacken admitted by Prof. P.

- 1. Sandstone is sometimes observed covering wacken.
- 2. Grayel, covered by flaty clay and basalt, has been obferved lying on wacken.
- 3. Veins of wacken fometimes traverse gneifs, porphyries, fandstone, &c.

Now, if wacken is clay hardened by the super-position of adduced to prove melted basalt, we must suppose the same effect to have been that it was not hardened by fire. produced

produced by the flaty clay, fandstone, or gravel. It is equally. impossible to explain wacken veins by this hypothesis of Professor Playfair.

From fection 255 to 259 is principally occupied in endea-Wedge-shape irvouring to shew, that from the wedge-shape irregularities in regularities of basalt, thickness, and the frequent great inclination of beds of basalt, they would not have been deposited from a state of solution in water.

To explain the appearances described in these sections will accounted for by be no difficulty to the Neptunian, they are what he daily meets aqueous deposiwith, and are illustrative of the deposition of these strata from equal surface; water. Let us conceive a bed of basalt deposited upon an unequal furface, that the water breaks it (either before it has be-broken by water; come folid, or afterwards) into shapes resembling those described by Professor Playsair, afterwards that sandstone is de- and sandstone deposited upon this broken bed; and we have a complete pic-posited thereon. ture and explanation of the wedge-shape, irregularities in the thickness, and every variety of inclination of the strata of basalt. This is illustrated by figure 1 and 2, Plate VIII.

A very strong objection has been made to the igneous origin Objection to the of basalt and grunstone, which has always appeared to me igneous formacompletely unanswerable: it is this: If these rocks had been that it is not formerly in a state of complete sussion, they should either ap-simple, but conpear as glass, or as a simple mass, not containing any substance dies. of a different nature from the mass itself.

Mr. Playfair, at the conclusion of his observations on the P. Sir James trapp-formation rocks, remarks, "Notwithstanding all this Hall's experiaccumulated and unanswerable evidence for the igneous for- in proof, that the mation of bafaltes, (the evidence I have examined in the preceding stony character pages) a great objection would still remain to our theory, were fusion. it not for the very accurate and conclusive experiments concerning the fusion of this fossil, made by Sir James Hall. strong prejudice against the production of any thing like a real ftone by means of fusion, had arisen even among those mineralogists who were every day witnesses of the stony appearance assumed by volcanic lava. They still maintained, on the authority of their own imperfect experiments, that nothing but glass can ever be obtained by the melting of earths and stones, in whatever way they are combined.

An ingenious naturalist, after describing a block of basaltes, in which he discovered such appearances as inclined him to admit its igneous confolidation, rejects that hypothesis, merely from the imaginary inability of fire to give to any substance a stony character: "Quelque mélange," fays he, "de terres que l'on suppose, quelque soit le tems que l'on emploie, il est très certain que l'on n'obtiendra pas, par la seul fluide igné, ni basalte, ni rien que lui resemble."

Sir James Hall's experiments have completely demonstrated the contrary of what is here afferted; they have added much to the evidence of the Huttonian fystem; and independently of all theory, have narrowed the circle of prejudice and error."

It is answered, that Sir James Hall's chrystallite and the original stone are different.

Sir James Hall, however, has only flown, that when bafalt, grunftone, or wacken, are completely melted and then flowly cooled, the glaffy character disappears, and the mass assumes that of indistinct chrystallization. This chrystallized mass he denominates christalite, as bearing a most striking refemblance to the natural stones. To the oryctognost, however, the glass and the indistinctly chrystallized mass are equally different from the original stone, therefore the arguments drawn from these experiments, in favour of the igneous origin of the rocks of the trapp-formation, are unfounded, and the objection which has been now flated remains in full force.

Objections to Sir Tames Hall's opinion, that lava has been in a fusion, &c.

Sir James Hall's opinion, that lava has been in a flate of complete fusion, and by flow cooling has acquired its present character, is probably incorrect. The well known facts, that ftate of complete lava has run into the fea, rivers, and lakes, without affuming the glassy character; that islands composed of cellular lava have been formed in the fea; and lastly, the low degree of heat possessed by lava streams, shew, that it has not been in a state of complete fusion; on the contrary, the following facts, which are to be found in the best writers on volcanos, render the Wernerian opinion, viz. that the fluidity of lava has rarely nion, that lava is exceeded the state of viscidity, extremely probable.

Wernerian opifeldom more than viscid; supported by its flow descent,

1. Lava flows very flowly, even over confiderably inclined planes.

2. In the European volcanos glass has hardly ever been observed. The Obsidian of Iceland, as described by Bergman and Olassen; that of Lipari; the glass of Vesuvius mentioned by Brierlac, are most distinctly products of water, as can be demonstrated by the Wernerian geognosie.

3. The chrystals of hornblende and felspar, which are fo freand the fcarcity of glaffy matter, quently found in lava, being wrapped up in it (eingewickelt), together with prove,

prove, as Werner has well observed, that these are the unalter- the unaltered ed chrystals of the mother-stone, which the volcanic fire has not tains.

had power to melt.

4. When porphiritic basalt is brought to the state of visci- Quick cooling of dity, the chrystals of felspar float in it unaltered, and the mass falt from its viswhich is now fufficiently liquid to flow, however quickly cool-cid state does not ed, does not assume the glassy character, but has the character character. of the commonest kinds of lava.

# ROBERT JAMESON.

## VII.

Description of the Blow-Pipe Apparatus of HAAS. From a Correspondent.

# To Mr. NICHOLSON.

DEAR SIR,

AFTER I had communicated to you the apparatus for the Letter. blow-pipe by means of alcohol, as used by Professor Pictet, which you favoured with a place in the last number of your excellent Journal, I was ftruck with the idea that the apparatus of Mr. Haas, which I mentioned in my letter, though made in the midst of London, is almost entirely unknown in this country. As this last is very ingenious, and affords many advantages in the use of the blow-pipe, I think you will render a fervice to your chemical readers by making them acquainted with its nature and construction. In consequence of this, I here subjoin a description and drawing of it, given by Mr. Orstein in Dr. Scherer's Chemical Journal, Vol. II. page 454. I leave it to your difcretion whether you think it deferving a place, and continue with the highest esteem,

SIR.

Your most humble and obedient,

N..N.

P. S. Concerning the older confiructions of the blow-pipe, Dr. Scherer refers in that paper to Bergman's Opusc. II. Writers on the Engström's

Engström's description of a mineralogical pocket laboratory (which may be seen in the second volume of Magellan's translation of Cronsted's Mineralogy, 8vo, Lond. 1788), and to Weigel's Versuch einer Geschichte Essay towards a history of the blow-pipe, and its use, in Crell's addition to the Chemical Annals, Vol. IV. page 262—283, and page 393—419; also, Vol. V. page 6—24, and page 198—226.

Description of the apparatus. THE delineation here added is upon the whole so distinct, that a few parts only require a detailed description. I begin with the candle tube, as it is the most essential part of this apparatus, on which all the others in some manner depend.

Plate VII. A is this candle-tube, which may be flided backwards and forwards upon the plate B. The rack in the tube ferves to raise the candle. The ball s, together with its tubes c c, may also be raised and lowered in order to direct the point e accurately towards the flame. The possessions of this apparatus will undoubtedly thank Mr. Haas for the ingenious arrangement of the inner parts of the ball, by means of which no moisture can enter into the point e.

The charcoal-stand C is likewise moveable upon B; but at the same time, as the drawing shews, its pillars k k are likewise, by means of the sliding-pieces i i, capable of a lateral motion; besides that at o the coal can be vertically turned upon the pillars.

Peculiar conftruction of fnuffers.

The conftruction of the fnuffers, and the reclangular fituation of their point m with their flaps to the handle n, will be found particularly important and ufeful. For, by reason of this form, the wick can, without any inconvenience to the hand, be always cut off in the same direction in which the stame is blown. and by stroking the wick with the point m (turned towards the body of the operator), it may be fo managed as to be always kept broader in the direction of the flame. This will be found more expedient than the usual method of bending the wick according to the flame's direction. The handle is laid hold on with the thumb and middle finger at rr. The fore-finger, for the purpose of snuffing the candle, opens or shuts the moveable flap by pushing the piece V to the right or left. It is because a common pair of fnuffers, in the use of this apparatus, would be not only extremely incommodious but even detrimental, that Mr. Haas invented this commodious and useful instrument.

The following articles which make this apparatus complete Additional appafor use, were fold by Mr. Haas \* in a neat box: 1. Various ratus. candles: 2. Points of different bores; 3. A hammer; 4. A round excavated piece of fteel to bound the products of the operation; 5. A pair of tongs; 6. A file; 7. A knife; 8. A magnifying lens; 9. A spoon made of platina; 10. A steel to strike fire, serving at the same time as a magnet; 11. A quadrangular magnet bar; 12. A thermometer; 13. An hydro-

### VIII.

Note respecting the Instruments by which the two Kinds of Electricity are distinguished, or its Direction ascertained. W. N.

ONCERNING the phenomena denominated electric, the Electricity is yet facts have hitherto afforded us little more than a few general understood. laws, which certain ingenious philosophers have connected by those hypotheses which constitute the theory. Among these Franklin's thethe two principal are, 1st, that a peculiar fluid, called the ory of plus and minus. electric matter, exists in or upon all bodies; capable of being accumulated by friction, eafily and most rapidly passing through or along metals, water, and charcoal, and difficultly or not at all through glass and other bodies not containing fluid water; capable of existing at the opposite confines of non-conductors. fo as to be in excess or plus at one surface, and in defect or minus at the other, in much higher degrees than either state could fingly exist; exhibiting the phenomena of ignition and found when it passes through the substance of non-conductors: and capable by its rapid motion of raifing the temperature of conductors to the highest intensity hitherto observed, as well as of exciting muscular action in all degrees, to the entire destruction of animal life.

2. The fecond received theory of electricity admits the ex- Nollet's theory istence of two sluids capable of being separated by friction; of two sluids. having the fame habitudes, as to conductors and non-conduc-

\* Mr. Haas is no longer in London: Five sets of this apparatus were fent to Portugal, where he is now established. The blowpipe and rest of the adjutage are made of brass, except the snuffers. -Tranfl.

militar

tors, as the simple fluid; strongly attracting each other, and not perceptible when combined in a due proportion; capable of existing separately at the confines of non-conductors in very high condensation; causing light and sound when they rush together through the fubstance of a conductor; and producing the highest temperature when they pass through or meet in non-conductors, &c.

According to the theory of plus and minus, that instrument kind will also distinguish the city.

Without entering farther into these theories, or the additions they require in order to fatisfy the facts, fuch as the attraction supposed to exist between conductors and the electric matter, the which shews the atmospheres of electricity, &c. the subject of the present notice requires we should use the language of one of the two. course of electri-former is most commonly adopted; and accordingly we should fay that an instrument which shall distinguish the plus from the minus state of electricity, will also shew the current or direction in which that matter is moved or carried.

Proposed means of doing this;

The fupporters of the theory called Franklin's, from the name of the philosopher who invented or at least explained it at an early period, have always been aware of the advantage it would be to their fystem if they could adduce any experiment by which the direction of their electric fluid might be aftertained. Dr. Franklin himfelf very modeftly offers conjectural reasons, why the one kind of electricity seemed to be a redundancy and the other a deficiency, chiefly grounded on the luminous appearances at the extremity of the wires in the electric states. Henly likewise observed the lights of the two electricities in the figure of the spark, of which the stem is always at the plus conductor and the ramifications at the negative; fo that the forked extremity of lightning will always denote the receiving body, whether it be the cloud or the earth.

from electric light;

figure of the fpark;

Lightning;

exhausted tube; also used his exhausted glass tube with good success, as an inftrument for the same purpose in which a receiving ball has a luminous atmosphere, and a giving ball throws out streams of

flame of a candle; light. The same philosopher first noted, that the slame of a fmall candle is blown towards a negative conductor, and from a positive one; though this experiment is not quite unequivocal: And he made feveral other experiments which cannot with confistent brevity be here detailed, having this leading

object in view.

Among the experiments made with this view, every one of None of these afford any decision which, it must be confessed, leave the great question of the

nature

nature and direction of the supposed electric matter undeter- of the real namined, there feems to be only two which afford the simple apture and course paratus we are in the habit of distinguishing by the name of instruments. As both instruments appear to deserve more im-Two instruments mediate notice than they have yet had, I have inserted them for the ordinary purpose; viz. in Plate VIII, where Fig. 4 represents the Galvanic tube, the Galvanic which, by the skilful management of Dr. Wollaston, is ren-tube; dered capable of diftinguishing the course of electricity. ingenious Mr. Acum, of Compton Street, Soho, who supplies chemists and philosophers with apparatus and materials of commodiously every kind, made the inftrument where this drawing was taken. Two fine gold wires are fixed in separate small glass tubes by fusion; and the ends of these tubes are ground away till the very fection or extremity of each wire is laid bare: at the other end of each tube appears a larger wire with a ball, which wire communicates with the gold wire within. two tubes are fixed by fusion in a larger tube, provided with a conical drawn aperture, through which it can be filled with water, all but a small vacuity or bubble, and then sealed. The apparatus is then complete; and if a stream of electricity or Galvanism be passed through it, the water is decomposed, and of the gas thus extricated, the larger stream of hidrogen shews the minus fide. It is probable this inftrument might be of use in atmospherical observations, in case large streams of electricity should pass through an elevated conductor at a low intenfity.

The other instrument, Fig. 5, was constructed by myself - and an appaabout twenty years ago, from an observation I made that low ratus with a low points cease to act as such with the plus at higher elevations, than with the minus electricity. So that if two balls be infulated, and a low point proceeds from one of them, electricity will fly through the interval in a spark if the point be plus, but will be filently transmitted if the point be minus.

This instrument, like the other, may be applied to distinguish the atmospheric changes; but it will be effectual only when the electricity is strong enough to give sparks.

#### IX.

Section and Plan of a moveable Crane, capable of heaving four Tons with four Men. Defigned, and executed for the Pier at Ramsgate, by Order of the Trustees for the Management of the Harbour at that Place. By Mr. Peter Keir, of St. Pancras. Communicated by Mr. P. Keir.

#### To Mr. NICHOLSON.

SIR,

Account and defcription of the crane.

THE crane represented in the annexed drawings was confiructed at the request of the Ramsgate Harbour Trust. It consists of a cast iron frame, marked A, Plates V. and VI. weight two tons, and 9 feet 7 inches square, supported on sour cast iron wheels, two of which traverse on a centre at B, by the rack work C, to steer it when moving from place to place. A cast iron upright shaft, D, weight 23cwt. supported by oak braces; the weight of the framing and wheel work is supported by a steeled pivot on the top of the shaft at F, round which the whole swings, including the men that work it.

The drawing is sufficiently plain, to require no farther explanation. It has been executed, and may be inspected at any time on Ramsgate Pier.

I am, SIR,

Your obedient humble fervant,

PETER KEIR.

Further particulars.

Upon inquiry, I find that the original inftructions were to make a crane capable of raifing four tons, and that the centre piece, or upright flaft, was at first made of oak: but that the strain for disengaging goods from their stowage requiring the occasional exertion of a greater power, it was found adviseable to substitute the present iron piece. The cost of this engine was four hundred pounds; but from the above circumstance, and particularly because the expence of patterns, &c. will not again be requisite, I am informed that others might now be made for three hundred pounds each.

W. N.

#### X.

Experiments on the Separation of Light and Heat by Refraction.

In a Letter from Sir H. C. Englefield, Bart. F. R. S. to Thomas Young, M. D. F. R. S. From the Journal of the Royal Institution, p. 202.

## DEAR SIR,

IN pursuance of your desire, I communicate to you the ex-Letter to Dr. periments which I made in consequence of Dr. Herschel's Young. most interesting discovery of the separation of solar heat and light by the prism. They carried with them, to my mind, a complete conviction of the truth and accuracy of the Doctor's affections on that subject. Should you judge them worthy of insertion in the Journals of the Royal Institution, I shall think myself honoured by their publication.

I am, Dear Sir,
Your faithful Servant,
H. C. ENGLEFIELD.

As I was defirous not only to ascertain, by actual observa-Introductory tion, the very curious discovery of Dr. Herschel, but to obvi-remarks. atte the objection made by Mr. Leslie \* to the mode in which his experiments had been made, I contrived an apparatus totally different in its arrangement from that which had been used by the Doctor; and disposed in such a manner that it was impossible that the thermometers should be affected by any foreign heat reslected from any part of it. As to the heat reslected from the floor, it could not influence the results, since it was the same, whatever colour was thrown on the ball of the thermometer.

As I had nothing to do with light, it was not necessary to A prism was darken the room; and as I wished to accumulate as large a placed in the portion of solar heat as possible, I placed the prism in an open room not dark-window, instead of admitting a beam through an aperture in ened. a shutter, as is the usual practice. The prism I used is a very good one; and it was lent to me for this purpose by Mr. Walker, of Conduit-Street. It is three inches long, and equilateral, each side being 1.15 inch broad.

<sup>•</sup> In his communications to the Philosophic Journal (quarto)
Vol. IV. p. 344, 416.

It was supported by an horizontal arm, projecting from a pole like that of a fire-screen. The arm could be adjusted to any height by a screw; and the prism was likewise capable of being turned on its axis to any required position.

The spectrum fell on a lens.

Through a flit which fuffered one colour only to pass.

The coloured spectrum was thrown on a very good lens, of four inches in diameter, and about 22 inches focal length. This lens formed part of a machine well known, and used for viewing prints; it stands on a foot adjustable in any direction, and to any height; and the mounting being of wood. and merely fufficient to support the lens, scarcely any heat can be accumulated in any part of it. The whole lens, as well as its mounting, was covered with a thick white pasteboard fcreen, in which was cut a flit of 3 inches long, and half an inch wide; this flit was over the centre of the lens; and through it any one of the colours required was admitted on the lens, while all the rest of the spectrum was totally excluded. A light wooden arm, of two feet long, projected at right angles from the lower part of the mounting of the lens. This arm carried a fmall screen of polified card, which received the image formed in the focus of the lens. This was found necessary, in order to find with certainty where to place the thermometer; when the focal distance was found, the fcreen was moved back, about a diameter of the bulb of the thermometer used, which was then held by the hand in the focus of the lens, which was done with great ease and certainty, as nothing more was necessary than to fix the eye on. the card screen, and keep the ball of the thermometer in the centre of the luminous image. The whiteness and polish of the screen totally precluded any accumulation of heat in it; nor indeed would fuch accumulation have been of any detriment to the experiments had it existed; for, as it must have been entirely owing to the ray under examination, it would have done nothing more than increase the effect of it on the thermometer.

A mercurial thermometer was held in the focus.

The thermometers used were mercurial, and very sensible. The scales were ivory tubes, embracing the stem, and graduated within. The ball of the instrument was therefore unconnected with any mounting; and no salse heat could possibly affect it. The balls were mostly blackened with Indian ink, carefully laid on; but some were used naked, and one, covered with white water-colour paint.

The

conf. r. = 1 5 1

beyond = 18

The lens, with its apparatus, was placed about three feet Positions of the from the prism; and, as the sun was pretty high in the greater lens and prism. part of the experiments, and the descending spectrum was in general used, the floor under the stand of the lens was in shade, from the wall below the window, and had been so all the day, as the window at which most of the experiments were made, fronted the south

Some part of this detail might feem supersuous, had not such stress been laid on the supposed accumulation of heat in Dr. Herschel's experiments, that it was necessary to obviate any objection on that head; or to show, at least, the utter improbability of any cause of that nature having affected the results of these experiments.

I now proceed to relate the experiments themselves: which I shall transcribe from the original notes made at the moment. April 6, 1801.

The apparatus being difposed as above described, the co-Experiment I. loured rays of the descending spectrum of the prism were suc-Maximum rise cessively thrown on the slit in the screen, covering the lens; of the therm, and the thermometer with a blackened ball, placed in the so-in blue = r cus of the lens, rose as follows:

| Comparatus being disposed in the prism were successively thrown on the slit in the screen, covering the lens; of the therm, and the thermometer with a blackened ball, placed in the so-in blue = r cus of the lens, rose as follows:

In the blue in 3', from 55° to 56°. or 1°.

Green in 3', from 54° to 58°. or 4°.

Yellow in 3', from 56° to 62°, or 6°.

Full red in  $2\frac{t}{2}$ , from  $56^{\circ}$  to  $72^{\circ}$ . or  $16^{\circ}$ .

In the confines of the red in  $2\frac{1}{2}$ , from 58° to  $73\frac{1}{2}$ °. hr  $15\frac{1}{2}$ °. Quite out of visible light in  $2\frac{1}{2}$ , from 61° to 79°. or 18°.

Between each of the observations the thermometer was placed in the shade so long as to sink it below the heat to which it had risen in the preceding observation: of course its rise above that point could only be the effect of the ray to which it was now exposed \*. A thermometer placed constantly in the shade near the apparatus, scarcely varied during the experiments.

April 17th, 11, A. M.

Three thermometers, used afterwards in the experiments, were exposed to the sun's rays until they became stationary.

\* In all the experiments the thermometer was continued in the focus long after it had ceased to rise; therefore the heats given are the greatest effect of the several rays on the thermometer in each observation.

The

Experiment II.	The apparatus being placed all as before, the blackened
	thermometer .
Black ther. rose	In full red ray in 3', from 58° to 61°. or 3°.
in red = 3° dark = 5	In quite dark in 3', from 59° to 64°. or 5°.
White therm.	The whitened thermometer
in red 3°.	In full red ray in 3', from 55° to 58°, or 3°.
dark of Black ther.	In quite dark in 3', from 58° to $58\frac{1}{2}$ °. or $0\frac{1}{2}$ °.

The thermometer with Naked ball . Whitened ball Blackened ball .

The blackened thermometer was now again placed in the quite dark, and rose in 3', from 58° to 61°. or 4°.

This is what I expected, as a thick fmoky haze had come on fince the experiments were begun, and increafed much towards the end of them.

Experiment III. Black ther. rofe in confine of red 120

in dark 4?

April 18, 11, A. M. Sun clear. All the apparatus as before-In the confines of the red.

Black thermometer in 3', from 59° to 71°.

White thermometer in 3', from  $57\frac{1}{2}$ ° to  $60\frac{1}{7}$ °. Clouds White do. do. 39 came on, and put a stop to the experiments.

Experiment IV. Black ther. rose in full red 120 White do. do.

3½°.

149

April 19, 31, P. M. Sun clear.

In the full red ray.

Black thermometer in 3', from 66° to 82° White thermometer in 3', from 66° to 693°.

In the confines of the red. In confine of red Black th. 1210

Black thermometer in 3', from 67° to  $79\frac{1}{2}$ °. But thin Ditto, quite dark ftreaky clouds had come over the fun.

In quite dark, half an inch out of the red. Black thermometer in 3', from 70° to 84°.

Transitions from fioned changes of temperature.

When the thermometer was carried into the faint red light, red to dark occa- it funk quickly; and rofe again as quickly when carried into the dark focus: but when carried into the dark on the other fide of the red light, it funk very rapidly, and did not appear to receive any heat at all. Thin clouds increased, and rendered the fun's light too faint for further experiments.

April 20, from  $10\frac{1}{2}$  to  $11\frac{1}{2}$ , A. M. Sun quite clear. Experiment V.

Although it could not be supposed that effects of the refracted light could differ in the two spectra, yet, in order to afcertain the fact, the horizontal spectrum was used in the subfequent experiments.

The

The apparatus all the fame as in the former.

In the full red ray.

Black thermometer in 3', from 67° to 71½°

Quite out of the ray.

Black thermometer in 3', from 68° to 77½°.

Black therm. full red 4° ½ dark 9½°

The ray was now so far removed from the flit in the screen, Darker to 10. that scarce any light was perceptible in the focus of the lens. The black thermometer was now placed near half an inch from the bound of the visible light in the focus, and rose in 3', from 69° to 791°.

The utmost edge of the prismatic spectrum was now removed Most dark 9°. an eighth of an inch from the edge of the flit in the fcreen; and no light was now visible in the focus of the lens.

Black thermometer in 3', from 70° to 79°.

Mr. Cary, optician in the Strand, and Dr. Hunter, were The difference present at these experiments, and repeatedly saw the thermo- perceivable by the touch. meter, in the fecond experiment, fink when carried into the light, and rife again when removed back into the dark. Dr. Hunter also received the focus on the palm of his hand, where the heat was fenfibly felt; and on shutting his eyes, and pointing with a long pen to where the heat was greatest, he always touched his hand beyond the vifible light.

As the red image has been continually mentioned in the Description of course of the above recited experiments, it may not be im- the red image. proper to describe it more particularly. The diameter of the red spot, formed by the ray in the socus of the lens, was just two tenths of an inch in diameter, at right angles to the length of the spectrum, and well defined: in the direction of the spectrum it was elongated, as might be expected, and less well defined.

When the whole visible spectrum from the prism was re- Very faint red in ceived on the fcreen which covered the lens, and the utmost the focus when the heat was edge of the red rays was removed a full eighth of an inch greatest. from the edge of the flit in the screen, there was still a faint blush of red, of a semioval form, visible when the focus of the lens was thrown on a white screen; and it was in these circumstances that the greatest effort of heat was constantly produced on the thermometer, not by placing it in the red light, but out of it, in the axis of the lens.

I have only to add, that in the course of the month of June Repetition of 1802, I repeated most of these experiments with the same experiments,

Vol. III.—OCTOBER, 1802. K apparatus,

apparatus, in presence of Mr. Davy, with the most complete fuccess; the sun's altitude being greater, the effect of his rays was fo great as to raise the thermometer in the invisible ray to 98°, while the vifible red never raifed it above 87°. At the fuggestion of Mr. Davy, we tried several experiments with respect to the power of the several coloured rays in ren-The blue rays il- dering Canton's phosphorus luminous; and we found, without a possibility of doubt, that the blue rays possessed that

luminate Canton's phosphorus

more than the red power in a much higher degree than the red. There was great reason to suspect that this power, like that of blackening the nitrate of filver, extended beyond the visible blue ray; but our apparatus was not prepared for the more delicate part of these experiments, which are only mentioned with a view of exciting further refearches on this very interesting subject, and of giving to Mr. Davy the credit due to him for having first thought of the experiment.

#### XI.

On the Expansion of the Elastic Fluids by Heat.\* By Mr. JOHN DALTON.

Essay.

THE principal occasion of this essay is another on the same Experiments and subject by Messirs. de Morveau and du Vernois in the first vol. of the Annales de Chimie. It appearing to them that the refults of the experiments of De Luc, Col. Roi, de Sauffure, Prieftley, Vandermonde, Berthollet and Monge did not . fufficiently accord with each other; and that it would be of importance to determine not only the whole expansion of each gas from two distant points, such as the freezing and boiling, but likewise whether that expansion be uniform in every part of the scale, they instituted a set of experiments Experiments of expressly for those purposes. The result of which was, that betwixt the temperatures of 32° and 212°, the whole expanfion of one gas differs much from that of another, it being in one instance about  $\frac{4}{10}$  of the original, and in others more than 12 times that expansion; and that the expansion is much more for a given number of degrees in the higher than in the lower

Guiton and Du Vernois.

part of the scale. These conclusions were so extremely discordant with and even contradictory to those of others, that I could not but suspect some great fallacy in them, and found it in reality to be the fact: I have no doubt it arose from the want of due care to keep the apparatus and materials free from moisture.

My method of experimenting on this subject is simple, and The author's therefore less liable to error. A straight manometer tube, simple manomefuch as has been mentioned, is duly divided into equal por-ter tube. tions of capacity; it is then dried by a wire and thread, and the open end inferted through a cork into a phial containing fulphuric acid, in order that the aqueous vapour may be drawn out of the tube; this is effential if we operate in temperatures lower than that of the atmosphere, otherwise not. For want and air dried by of this attention, Col. Roi, in his valuable paper in the Philos. fulph. acid. Trans. vol. 67, has been led into some erroneous conclusions. A fmall column of dry mercury is then let down to a proper point in the manometer, and it is ready for experiment with common air.

It requires some address to fill the manometer with any other Method of filling gas.—I succeeded best as follows: filled the tube with dry the manometer. mercury; then pushed down a wire with thread, so that when the wire was got to the end of the tube, a thick-covering of thread just entered the open end, and held the mercury like a cork, fo that the tube could be inverted without lofing the contents; then having a glass funnel with a perforated cork over the water apparatus, containing the gas, I flipped the manometer through the hole in the cork, and putting my hand into the water under the funnel, drew the wire out of the manometer, and with it the mercury; upon which the gas entered the manometer. For carbonic acid gas, I opened the Carbonic acid fealed end of the manometer, drew it out to a capillary bore. gas. and forced a ftream of the gas through the tube; then putting my finger on the other end, fealed it again by a blow-pipe, and let down a small column of mercury to the proper point.

When the manometer was to be exposed to a heat of 212°, Simple method I used a Florence flask, with a long glass tube corked into it, of applying the boiling heat. in fuch fort that as much of the manometer as was necessary to be exposed to the temperature might be in the tube; then the water at the bottom of the flask was made to boil violently, so that a constant stream of vapour issued out of the top of

the glass tube, which was found to raise the thermometer to 212°. Small specks of white paint were put upon the divisions of the manometer together with numbers which were difcernible through the containing tube. For lower temperatures, a deep tin vessel containing hot water was used, in which the manometer was immerfed, the water being well agitated previoufly to each observation.

The conclusions Berthollet, &c. are accurate; but those of Morveau and du Vernois not fo.

From a great many experiments made in this way on comof De Luc, Roy, mon air, and likewife upon hydrogenous gas, oxygenous and nitrous gafes, and carbonic acid gas, I can affert that the conclusions of De Luc, Roi, Saussure, Berthollet, &c. are nearly accurate throughout, and that those of de Morveau and du Vernois are extremely inaccurate in the higher temperatures.

Expansion of common air.

I have repeatedly found that 1000 parts of common air of the temperature 55° and common pressure, expand to 1321 parts of the manometer; to which adding 4 parts for the

Intermediate expansion.

corresponding expansion of glass, we have 325 parts increase upon 1000 from 55° to 212°; or for 157° of the thermometric scale. As for the expansion in the intermediate degrees, which Col. Roi's experiments shew to be a slowly diminishing one above the temperature of 57°, but which de Morveau's on the contrary shew to be a rapidly increasing one in the higher part of the scale; I am obliged to allow that Col. Roi is right, though it makes in some degree against an hypothesis I have formed relative to the fubject; he has certainly however made the diminution too great from 72° downwards, owing to his not perceiving that he actually destroyed a portion of the clastic fluid he was operating upon (aqueous vapour) in reducing its temperature fo low; if his air had been previously dried by fulphuric acid, &c. he would not have found fo re-

Other gafes agree with com-

mon air.

The refults of feveral experiments made upon hydrogenous gas, oxygenous gas, carbonic acid gas and nitrous gas, which were all the kinds I tried, agreed with those on common air not only in the total expansion, but in the gradual diminution of it in ascending: the small differences observed never exceeded 6 or 8 parts on the whole 325; and differences to

markable diminution below 72°. My experiments give for  $77\frac{1}{2}$ ° above 55°, 167 parts; for the next  $77\frac{1}{2}$ ° only 158 parts; and the expansion in every part of the scale seems to be a gra-

dually diminishing one in ascending.

this amount will take place in common air, when not freed from aqueous vapour, which was the fituation of all my factitious gales.

Upon the whole therefore I fee no sufficient reason why we The law is gemay not conclude, that all elastic stuids under the same pressure neral for all expand equally by heat, - and that for any given expansion of mercury, the corresponding expansion of air is proportionally something lefs, the higher the temperature.

This remarkable fact that all elastic fluids expand the same Supposed cause quantity in the same circumstances, plainly shews that the ex- they have to pansion depends folely upon heat: whereas the expansion in attraction. folid and liquid bodies feems to depend upon an adjustment of the two opposite forces of heat and chemical affinity, the one a constant force in the same temperature, the other a variable one, according to the nature of the body; hence the

general laws respecting the absolute quantity and the nature of heat, are more likely to be derived from elastic sluids than from other fubftances.

unequal expansion of such bodies. It seems therefore that

In order to explain the manner in which elastic fluids ex-Hypothesis to pand by heat, let us assume an hypothesis that the repulsive explain the law force of each particle is exactly proportional to the whole of expansion. quantity of heat combined with it, or in other words to its

temperature reckoned from the point of total privation; then, fince the diameter of each particle's sphere of influence is as the cube root of the space occupied by the mass, we shall

have \( \frac{1000}{1000} \): \( \frac{1325}{1325} \) (10 : 11, nearly) :: the absolute quantity of heat in air of 55°: the absolute quantity in air of 212°. This gives the point of total privation of heat, or Singular coinciabsolute cold, at 1547° below the point at which water freezes, dence as to the Dr. Crawford (On Animal Heat, &c. page 267) deduces the faid point by a method wholly different to be 15320, -So near a coincidence is certainly more than fortuitous.

The only objection I see to this hypothesis is, that it neces- An objection if farily requires the augmentation of elastic fluids for a given we admit the expansion of merquantity of heat to be greater in the higher temperatures than cury to be as the in the lower, because the cubes of a series of numbers in heat; but not so if mercury follow arithmetical progression differ more the larger the numbers or the law of all roots: but it has just been shewn that in fact an augmentation other fluids. of a contrary kind is observed. This refers us to the con-

fideration whether the mercurial thermometer is an accurate measure of the increments of heat; if it be, the hypothesis fails; but if equal increments of heat cause a greater expansion in mercury in the higher than in the lower temperatures, and that in a small degree, the fact noticed above instead of being an objection will corroborate the hypothesis.—Dr. Crawford determines the expansions of mercury to be very nearly in proportion to the increments of heat: M. De Luc makes them to be less for a given quantity of heat in the lower than in the higher part of the scale; and in a ratio that agrees with this hypothesis. Now as every other liquid we are acquainted with is found to expand more in the higher than in the lower temperatures; analogy is in favour of the conclusions of De Luc, that mercury does the same.

Short account of fimilar refearches by Mr. Gay Luffac.

In the Bulletin des Sciences, there is a notice of a memoir of Cit. Gay Lussac on the dilation of the gases and vapours; and the memoir itself is given in the \$\frac{4}{3}\text{d}\$ volume of the Annals de Chémie, whence I shall, as early as convenient, give either a full abstract or a translation. This able author gives the same conclusion as that of Mr. Dalton, and, like him, attributes the errors of preceding experimenters to the humidity of the gases they examined.

His method.

His experiments were made by expelling the gas from a glass ball properly fitted up, and determining the expansion by weighing the apparatus, after water had been fuffered to come in during the cooling, to occupy the space of the extruded gas. He very candidly informs us, that Cit. Charles had deduced the general refult of the equal and fimilar expanfion of all the gases by heat, under like pressure, fifteen years before, though he did not publish it; and he gives an historical account of the labours of his predeceffors; fo that we must neceffarily conclude that he could not have feen the late publication of the Manchester Society, in which the extensive refearches of Mr. Dalton are inferted; and wherein it appears that the preceding memoir was read about a year before that of Mr. Gay Luffac was read to the Institute. The French Philosopher finds the expansion or increase of dimension, between the freezing and boiling water points, to be 0.375, which

Mr. Dalton's experiments were earlier.

which gives for the space between 55° and 212°, or 157°, How far they the quantity 0.327 instead of 0.321, as Mr. Dalton finds it. agree. This difference of somewhat less than one fiftieth part of the expansion, may perhaps have arisen from a difference in the scales of operation, of perhaps the difficulty of producing a given temperature through the whole mass of a stuid, &c.

#### XII.

Account of some Experiments made in the Laboratory of the Royal Institution, relating to the Agencies of Galvanic Electricity, in producing Heat, and in effecting Changes in different stud Sub-stances. By H. Davy, Prof. Chem.

I. IT has been shown, by a very interesting experiment made Discovery of in France, by Messrs. Fourcroy, Vauquelin, and Thenard, Fourcroy, Vauquelin, and Thenard, quelin, and Thethat the power of galvanic batteries containing large plates, tonard, of the ignite metallic substances, is much greater than that of batte-greater ignition ries composed of an equal number of small plates; though their nic plates. agencies upon water, and upon the human body, are nearly the same.

In examining the circumstances of the action of a galvanic The trough with apparatus, or trough, constructed in the Royal Institution, and the same relation containing twenty series of plates of copper and zinc, square, of chemical and thirteen inches in diameter, I observed that the same relations between chemical agency and the production of galvanic electricity existed as in other cases. When pure water was used for filling the cells, the sparks, as well as the shocks, were extremely indistinct; and the battery was capable of ig-Water had little niting only about a line of iron wire of  $\frac{1}{170}$  of an inch in diame-effect; falts and acid much more. With solution of muriate of soda it acted better; and diluted nitric acid was still more efficacious. With this last substance, it became capable of rendering white-hot three inches of the iron wire of  $\frac{1}{170}$ , and of causing two inches to enter

In comparing the effects produced by a folution of nitrous Nitrous acid did acid, of the specific gravity of 1.4, in about fixty parts of wa-not act by ats conducting power, ter, with those occasioned by a concentrated solution of carbo- For it is much nate of potash, the acid was found to produce by much the more powerful than carbonate of greatest intensity of action; which can hardly be ascribed to potash, though is any conducts much less.

into fusion.

galvanism.

any other cause than its chemical agency; for, with regard to conducting power, it appeared very much inferior to the other There is every reason to believe, that with pure water, that is, water deprived of air and of all faline substances, no action would be produced in this battery. I was unable to ascertain the fact by a direct experiment; but I found

Water appears to repeatedly, that a pile, composed of thirty-fix series of square act, not as a con-plates, of copper and zinc, of five inches in diameter, lost-its ductor, but by activity in nitrogene and hydrogene gases, in about two days; its gas. and it was conftantly reftored by common air; and rendered

more intense by oxigene gas.

II. When the galvanic battery, with large plates, was in Water and oil made to boil by full action, it was found that a w of an inch in diameter, and two feet long, when placed in the circuit, was rendered fo hot as to cause a small quantity of water, brought in contact with it, speedily to boil. It continued warm for many minutes; and, by an occasional momentary interruption and completion of the circle, the heat was permanently kept up. When three or four inches of the wire of  $\frac{1}{170}$  were placed in any part of the conducting chain, they continued red-hot for more than a minute; and, by a succession of interruptions and contacts, they were kept partially ignited for five or fix minutes. When that part of the communicating chain containing the small wire was introduced into a small quantity of ether, alcohol, or oil, the fluid foon became warm; and olive oil, the only substance that was exposed for a sufficient time, was made to boil.

Charcoal made red hot under water, oils, &c. tion of gales.

III. When two small pieces of well burned charcoal, or a piece of charcoal and a metallic wire, were made to complete water, oils, ac. the circle, in water, vivid sparks were perceived, gas was given out very plentifully, and the points of the charcoal appeared red-hot in the fluid, for fome time after the contact was made; and, as long as this appearance existed, elastic sluid was generated, with the noise of ebullition. The fensible phenomena were nearly the same with the volatile and fixed oils, ether, and alcohol; and, by means of charcoal, the spark could be produced in concentrated fulphuric and nitric acids, which are amongst the best of the less perfect conductors.

the The gases produced from different fluids by the galvano-Nature of electric spark, were examined; and as the results were, in gafes ;

most

most cases, what might have been expected from theory; the analysis of them was not made with very minute attention.

When water was acted upon by sparks taken from two from water; pieces of charcoal, the elastic products evolved were about  $\frac{1}{8}$  of carbonic acid,  $\frac{1}{8}$  of oxigene, and the remainder an inflammable gas, which required a little more than half its volume of oxigene for its combustion. With gold and charcoal, the gold being on the zinc side, the gas produced appeared to be chiefly a mixture of oxigene and hidrogene, for it diminished  $\frac{7}{10}$  by the electric spark.

The gas difengaged from alcohol, the spark being taken by from alcohol; gold connected with the zinc end, and charcoal, was a mixture of nearly two parts of oxigene and eleven parts of inflammable gas, which appeared to be partly light hidrocarbonate.

Ether, in the same method of operating, gave four parts of ether;

oxigene and twelve parts of inflammable gas.

From fulphuric acid, oxigene and hidrogene were produced fulph. acid; very rapidly, (the oxigene being more than sufficient for the saturation of the hidrogene by combustion,) and the acid became blue.

The gas from nitric acid detonated with great violence by nitric acid. the electric spark, and the residuum was oxigene, mixed with a little nitrogene.

The products from the acids, there is every reason to believe, were evolved chiefly in consequence of the decomposition of the water they contained. And, in operating upon these substances, as well as upon pure water, a portion of the elastic shuids must have been produced at the time of the filent transmission of the electricity, during the momentary interruptions of contact. The apparent ignition of the charcoal in the Ignition of the different shuids depended, probably, in some measure, upon its charcoal accounted for. being surrounded, at the moment of contact, by globules of gas, which prevented the heat, produced at the points of it, from being rapidly carried off by the shuid.

When the spark was taken by means of iron wires, in phos-Gas from sufed phorus rendered sluid by heat, under a stratum of water, per-phosphorus. manent gas was produced from it, but in a quantity too small to be examined, after a process that continued an hour. I purpose to repeat the experiment, with conductors of dry charcoal.

The large plates gas; provided power be good.

IV. When gold wires, connected with the ends of the batdifengage most tery, were made to act upon fluids in the common method of the conductor's communication, being placed at a diffance from each other, it was found that the rapidity of the evolution of the gases was much more influenced by the conducting power of the fluid, than it is in common cases with small plates. In comparing the action of a battery of twenty plates, of five inches in diameter, upon sulphuric acid, nitric acid, and various saline solutions, with that of the large battery, it was observed, in feveral experiments, that the gas was difengaged much faster, and in larger quantities, from the wires connected with the large plates, whilft the action of the two arrangements upon water was nearly the fame. This fact, combined with other facts of the same kind, feems to show, that the quantity of electricity excited in the arrangements with large furfaces, is much greater than that produced in those with small surfaces; and that it is capable of passing with facility through the more perfect conductors, whilst, from the nature of the series, its circulation is impeded, comparatively to a great extent, by imperfect conductors; a conjecture that has been already formed by different philosophers.

Attempt to procoal.

V. As the great quantity of electricity made to circulate duce change in through perfect conductors, by means of the large apparatus, muriatic gas, by the galvanic ig-increases their affinity for oxigene more perhaps than any nition of char-known agent, and as charcoal by means of it can be rendered white-hot, and kept in constant combustion in oxigene gas or atmospherical air, I thought of trying the effects of the electrical ignition of this fubstance, upon muriatic acid gas confined over mercury.

Experiment.

The experiment was made by means of a small glass tube \*, containing a flip of platina hermetically fealed into it, and having a piece of charcoal attached to its lower extremity: the communication was effected by means of iron wires; and the charcoal was made white-hot, by fuccessive contacts continued for nearly two hours. At the end of this time, the muriatic acid gas had diminished a very little in volume: much white matter had formed upon the charcoal, which was not fenfibly confumed. When the gas was examined, 3 of it were infantly absorbed by water, and the remainder proved to be

<sup>\*</sup> For a description of this apparatus, see p. 214.

inflammable. The process was repeated three times; and, Effects; when the spark was most vivid, a white cloud was always perceived at the moment of its production, I am inclined to attribute this phenomenon, and the other phenomena, to the decomposition of the water held in solution in the gas, by the charcoal and the mercury adhering to it; and the white matter was probably muriate of mercury. The acid gases are rapidly absorbed by charcoal; and this substance, when well made, will take up more than 30 times its volume of muriatic acid gas; so that, in the process of ignition, a part of the water and of the acid must have been acted upon in a very condensed state.

The want of fuccess in this experiment, the results of which very similar to are very similar to those obtained by Mr. William Henry \*, those of Mr. W. in his trials with common electricity, prevented me from car-mon electricity. rying on the process upon fluoric acid gas, as I had at first in-No decompositended. Many of the compound gases that are decomposable by heated charcoal, might probably, however, be analysed in a very simple manner, by means of the ignition of that substance by galvanic electricity; and this mode of operating may be conveniently applied, for ascertaining the relations of the affinities of charcoal for the constituent parts of compound gases at very high temperatures.

#### SCIENTIFIC NEWS, &c.

#### NATIONAL INSTITUTE OF FRANCE.

Prize proposed at the Public Sitting of the 15 Germinal, in the 10th Year. (April 5, 1802.)

THE general conditions are as usual; namely, that the Conditions. name of the author shall be concealed, and his rank diffinguished by a sentence or device, which shall also be written on a sealed billet, containing his name and address. The works are to be addressed (carriage paid) to one of the Secretaries of the Institute. Neither the treatises, nor any draw-

<sup>\*</sup> Phil. Trans. 1800, p. 188, or Philos. Journal, quarto, iv. 209, 245.

ings or machines thus forwarded, will be reftored; but the authors may take copies, and may again receive their models or machines, upon substituting regular drawings in place of them. The Commissioner of the Funds of the Institute will deliver the gold medal to the bearer of the Secretary's receipt, if given for the treatise, &c.; but where no receipt exists, the author must himself appear, or fend his procuration.

#### Subject of the Prize of Chemistry,

The class of mathematical and physical sciences having proposed in the year 8 the following prize question, to be decided upon at the sitting above mentioned:

Prize question respecting ferments. What are the characters which distinguish in regetable and animal matters those which serve as ferments, from those in which they cause or excite fermentation.

And none of the memoirs having fulfilled the conditions of the program, the class proposes the same subject again for the year 12.

The prize will be a gold medal, of the value of one kilogramme, which will be given at the fitting of the 15 Germinal of the year 12 (April 5, 1804).

Prize Questions from the Batavian Society at Haarlem.

Prize questions.

The Batavian Society at Haarlem has published a number of subjects of prizes at their sitting of May 1, 1802; among which are the following:

Respecting ma-

1. "How far it is known, from the latest discoveries in the physiology of plants, in what manner the disserent manures on different foils are favourable to the vegetation of plants; and what indications may we deduce from the knowledge required on this subject, to direct our choice as to manures, and the fertilization of uncultivated and arid lands?"

This question is continued to the 1st of Nov. 1802.

- motion of the

2. "What is the actual state of our knowledge respecting the motion of the sap in trees and plants? In what manner can we acquire a more complete knowledge respecting the obfcure and doubtful parts of the subject? And is it possible to deduce, by decisive experiments, such indications as may be useful for the cultivation of trees and plants?"

The Society has determined to repeat this question, to which answers are to be sent before November 1, 1803.

3. " The

3. "The Society demands a theory, or physical explication, — the ascent of which shall clearly and distinctly show the causes of the ascent smoke. of smoke in chimnies, or those which prevent its rising. Together with rules, deduced from this theory, for the construction of chimnies, which shall point out the circumstances necessary to be attended to in the several cases to prevent smoke from entering the apartments?"

This question is here again repeated, and the answer is demanded before the 1st of November, 1803, in order that the authors of memoirs may be enabled to correct them.

4. "What indigenous plants, hitherto not used, may, from — dying plants. well confirmed experiments, afford good colours, which may be prepared and used with profit? And what exotic plants may be cultivated with profit upon the less fertile or uncultivated lands of the Republic, in order to the extraction of colours?"

The Society has determined to repeat this question, which is continued without limit of time.

The questions for the present year are the following:

5. "What do we learn, from the latest observations, re-effect of oxfpecting the influence of the oxigen of the atmosphere, whe-igen on colours ther combined or not, together with the action of light upon the changes of colour? And what advantages may be derived from this knowledge?"

The Society desires that it may be shown concisely, and with precision, what is well proved by observations or experiments, in order that the actual state of the science with regard to this subject may be more readily apprehended, and greater advantages obtained in trade, or in the other branches of economy.

The term of concurrence is November 1, 1803.

6. "What light has been thrown upon the manner in which — nourishment plants acquire their nourishment, in consequence of the disco- of plants. veries respecting the decomposition of water and atmospheric air? And what deductions can be made from this knowledge for the improvement of useful vegetables?"

The term of concurrence is November 1, 1803.

7. "What facts are well proved with regard to the purification of corrupted water, and other impure substances, by charcoal? How far is it possible to explain this effect by the principles of chemistry? And what further advantages may be thence derived?" The term of concurrence is November 1, 1803.

The following questions were proposed formerly, but the term of concurrence is fixed to the first of November, 1803.

physiology of the . human body;

1. "What light has the new chemistry thrown upon the physiology of the human body?"

discases;

2. "To what extent has this light been of use in rendering us better acquainted with the nature and causes of certain diseases? And what useful consequences, more or less grounded on experience, may be thence deduced for improving the practice of medicine?"

remedies;

3. "To what extent has the new chemistry been of use to afford precise notions concerning the action of certain external and internal remedies long in use or lately recommended? And what advantages may be obtained from a more exact knowledge in this respect in the treatment of certain disorders?"

explanations.

As many philosophers have mixed hypotheses of slight foundation with the applications they have made of the principles of the new chemistry to the functions of the human body. which is no doubt very hurtful to the progress of science, which might be so highly improved, if, according to the rule of Lavoifier, nothing were adopted either in chemistry or the application of its principles, but what shall be founded on decifive experiments; the Society defires that those who shall answer these questions should precisely distinguish facts from hypothefes; and that with regard to the latter, they should be merely pointed out, and their flight foundation briefly shown. For the principal aim of the Society in these questions, is to procure those who exercise the practice of medicine and surgery in the Batavian Republic, and are not fufficiently acquainted with the progress of the new chemistry and the application of its well established principles to physiology, pathology, and theurapeutics, fuch memoirs as may eafily show them what light the new chemistry has in effect thrown upon these sciences; and also those doctrines which, being of flight foundation, too hastily adopted, or in themselves doubtful, are not entitled to confidence.

Judgment will be given feparately upon the memoirs respecting each of these questions. They who propose to answer more than one, are therefore requested to do it separately.

The above were proposed in 1799.

Philosophy of fire in the ma4. "What are the principles of the philosophy of fire, respecting the production, the communication, and confinement of heat, which are necessary to be known in order to form a nagement of right judgment of the methods of using combustibles; and how &cc. can we, according to these principles, improve the fires for warming apartments and stoves for kitchen use, in order to fave as much as possible the combustibles in use in this country?"

The above was proposed in 1801.

5. "What is actually known respecting the causes of the stagnant water, corruption of stagnant water, and is it possible to deduce from what is known or may be proved by decifive experiments, what are the most effectual and safe means to prevent the corruption of stagnant waters?"

Prepared in 1801.

The prize for each question is a gold medal of thirty ducats. Prize, a medal of memoirs may be written in Dutch, French, or German; of 30 ducats. but only in Italic characters. They must be accompanied with a fealed billet, containing the name and address of the writer, and fent to M. Van Marum, Secretary of the Society.

The Batavian Society has proposed as one of the prized sub. Fermentation. jects to be decided on the 1st of May 1802, the following queftion: "What do we learn from the latest discoveries in chemistry respecting the nature of fermentation, and what advantages may refult with regard to certain trades in which fermentative matters are used?" It was not thought fit to adjudge the prize; but without opening a new concurrence, the Society has engaged the author of a memoir, of which the device is, Tout est important dans les voïes de la nature, to support his theory upon more decifive experiments, and to extend its application to a greater number of manufactories; with promife of the crown if he shall accomplish his purpose before the end of 1803 or 1804.

#### Sixth Descent of Garnerin with his Parachute.

On the 21st of September 1802, at seventeen minutes past Account of Garnerin's descent five in the evening, the aeronaut Garnerin fent off from the by a parachute. Parade in North Audley Street, an exploring balloon, which took a north-east direction until it attained a very considerable elevation, and foon disappeared. At fifty-eight minutes after five the great balloon itself ascended with the parachute depending, and beneath it the cylindrical basket in which Garnerin himself stood and repeatedly waved his slag. Its slow ascent

afcent in the calmness of a beautiful evening, with the fixed and still attention of the inhabitants of a great metropolis, to the amount of many hundreds of thousands, produced a fingular and very impressive effect on the mind. This pause of expectation which lasted nine minutes and a quarter, and carried the aeronaut fo high as not to be perfonally diffinguished, was fuddenly terminated by the separation of the parachute from the balloon. It fuddenly defcended for near a fecond, and then expanded. The balloon proceeded immediately to the fouth, and was found the next day in Kent, about twelve miles out Admiration and furprize now engaged the minds of all the spectators; which soon gave place to a marked sentiment of terror from the extreme vibrations of the apparatus. The descent appeared flow and regular, but its vibrations (performed in about fix feconds each) were fo violent as frequently to carry the basket nearly as high as the level of the parachute itself, which collapsed at its lower edge every time, so as to be nearly half closed, and opened again when it refumed its perpendicular fituation. However firm the confidence of this daring adventurer may have been in the truth of his combinations, he must have have found himself in a situation requiring the exertions of all his courage. He descended in safety in a field near Pancras, having been very nearly fix minutes in his descent. He seemed to have been upwards of a mile high, or probably about fix thousand seet; and on this supposition he fell at the rate of fixteen feet in a fecond. This is the velocity that would have been produced by jumping from the height of about four feet, and must therefore have been very fafe.

Velocity of his

# cities at front thele or its confingent parts; it placks

menticoner " entrine of coffee position in a large

# NATURAL PHILOSOPHY, CHEMISTRY,

which be found in a sequence by founds, that the round or nimes that the round of similar to be comes for this test by some

## THE ARTS. 12 1 in the state of

July aninhmed NOVEMBER, 1802, was a life We mark the to

of it can be be be seen as a first of the second of the se Mr. Gough's Letter, at Page 36 of the prefent Volume. On the Phenomena of Sound. Adams will been midanidano and of

#### motions. The comparison uppears to me to be periodically appoints, and Tomme, NICHOLSON. ald it is made

what enafe there is left for further conference. . R I &

I beg leave once more to recomin ad the grave lamments AM not certain that any answer, that I can make to Mr. Further discus-Gough's reply to my letter, will add very materially to the fion apparently elucidation of the subject in dispute; a part of my object is unnecessary. already attained, for Mr. Gough feems to be no longer difposed to contend for Dr. Smith's infallibility, and he appears to have formed in some measure a more complete idea of my original opinions, while he imagines that I have introduced fome new modifications of them. I cannot however avoid remark- But Mr. Gough ing, that Mr. Gough has wholly omitted to notice the funda- has not noticed mental fact, which I stated as affording the most satisfactory of the grave harmonics. all proofs of the coalescence of sounds; that is, the production The facts. of a faint, but very audible, graver found from the union of two acuter ones, a phenomenon fo well know to musicians, that I can scarcely suppose a person, who is ignorant of it, properly

....

Vot. III. - NOVEMBER, 1802.

qualified

install since " The

The grave harmonic is not a mixture, but a compound.

qualified to discuss the subject of harmonics; and so important with regard to the question at present in dispute, that I can fcarcely conceive how a person, acquainted with it, can omit to endeavour to reconcile it with his opinions, especially when challenged by his opponent to take it-into confideration. Here is a product of combination which polletles properties totally different from those of its constituent parts; its pitch is always much lower, its quality of tone is perfectly fingular, and the hearer is fometimes almost inclined to imagine its direction to be different from that of the original founds; a circumstance which he fometimes expresses by faying, that the found "rings in his ears." Mr. Gough cannot controvert this fact by any reasoning: if he can persuade me that such a sound is a mixture and not a compound by partial coalescence, we shall then be nearly of the same opinion.

Illustration from dulum.

Will Mr. Gough deny, that when a fmall pendulum is fufa compound pen- pended to the heavy weight of a larger one, its motion is compounded of larger and smaller vibrations? Or will he insift that the result is a mixed motion, and not a compound one? The eye certainly follows each vibration separately, although the vibrations may be fo proportioned that a joint order and fequence may be observed, which may produce effects peculiar to the combination, and discoverable in neither of the original motions. The comparison appears to me to be perfectly apposite, and if Mr. Gough allows it, I can scarcely imagine what cause there is left for further controversy.

I beg leave once more to recommend the grave harmonics One in the to Mr. Gough's notice, and to intreat him to suspend any further reply to my remarks until he has made himself thoroughly acquainted with these interesting phenomena.

- when and o tod Dam, Sir, Topoco . id. not have id. to

to an atrial trail the said survey of

Your faithful humble fervant,

THOMAS YOUNG.

Royal Institution, was a war to any allow worth OA. 12, 1802.

esp 13 (12.5)

restance of the comments a filing brook rimal and some of

On the Galvanic Effect of very minute Particles of Zinc and Copper in Water. In a Letter from Mr. WILLIAM WILSON.

## To Mr. NICHOLSON.

2, 1,

SIR, London, October 11, 1802.

I WAS lately grinding some small plates of copper upon some Copper and zins plates of zinc (with fine emery), and after having washed them were ground with emery. in a cup of water to cleanfe them, I left them about five hours. The mud after When I returned I found the water in the cup covered with a fubfidence emitdirty froth, and upon taking up the cup to examine it, the agitation caused innumerable bubbles of air to rise from the sediment at the bottom of the water. By continuing to agitate the and at remote water for feveral minutes, they decreased in quantity, and at intervals. last disappeared. I now left it about four or five hours, and then examined it again, when the bubbles were nearly as copious as ever, and, with nearly the fame quantity of agitation, again ceafed to be produced.

I examined this water every four or five hours for five days, The effect lafted and the appearance was the fame, though weaker and weaker. five days. After the fifth day I could not find that any bubbles were generated; I therefore (supposing this appearance to be occa- Copper and zins fioned by the particles of copper and zinc forming a kind of filings were galvanic feries) took equal quantities of filings of copper and tried, zinc and put them into a glass of water, and stirred it about to mix them as much as possible. In about three hours afterwards there were some bubbles of air adhering to the filings, which by agitation rose to the top of the water; and the same pheno- with the same menon was repeatedly exhibited. In short these metallic par-effect. ticles acted precifely fimilar to the former, with only this difference: It required more violent agitation; the air bubbles were Explanation of larger, and the property disappeared much sooner, for there the difference. were no bubbles generated after the third day. This difference I supposed to be owing to the difference in the fize of the particles of the metals; for in the first case the particles, by reason of their smallness, would form a more numerous series and extensive surface of contact, than the filings could in the latter.

The impalpable powder of the metals.

Lastly, I repeated the first experiment by washing some emery very fine, and flightly ground with it a plate of zinc on another of copper; and then washed them in a glass of clear water. It was feveral hours before the fediment had all fallen to the bottom of the glass; but when it had, it acted nearly with the same power as in the first case, and continued to act quite as long \*.

. If, Sir, you should judge the above account worthy of a place in your Philosophical Journal, I shall consider myself very much obliged by its infertion.

I am, Sir, respectively to the control of th

the standard below of the warts, "It continuing on finite life and at amora

Your obedient humble fervant,

ing with the will wave by do and gu guider WM. WILSON.

### water for the parties of the same of the contract of the same of the same.

Paragraph with an analy to be a policy to the capacita for On the Colours obtained from Metallic Oxides, and fixed by Means of Fusion on different Vitreous Bodies. By ALEX. BRONGNIART, Director of the National Manufactory of Porcelain at Sévres, Engineer of Mines, &c.

(Concluded from Page 101.)

Concerning the Red, Rose, and Brown Colours obtained from Iron. State of a state of parties.

billion ?

put S, sup of oding life Red colours from 1 HESE colours are made from red oxidated iron prepared the oxide of iron. with nitric acid. The oxides are calcined still more by exposing them to the action of fire. If too much heated, they change to a brown.

Composition.

Their flux is composed of borax, fand, and minium in small quantity.

They may be Tobstituted for the oxide of gold.

These are the oxides which afford the rose and red colours that may be substituted instead of the same colours made from oxide of gold. If properly applied on hard porcelain, they never change. I have made rofes with these colours, and there

\* These galvanic facts appear to constitute one of the cases detail. ed with larger pieces of metal, and obscurely explained by Fabbroni in his curious paper in the Philof. Journal, quarto feries, IV. 120. -W. N.

was no difference between the flower before and after the baking, except that brilliancy which colours naturally receive by fusion.

The colours may either be previously fused or not at plea-

In a violent fire they either partly disappear, or produce a dull How affected by and brick-dust red colour, which is not at all agreeable.

Their composition is the same either for tender porcelain or They are good for glass. They do not change on the latter, but on the for tender porcelain; mer they almost intirely disappear by the first fire, and they must be laid on very heavily in order to have any part visible.

It is to the presence of lead in their glaze that this fingular but must be heavily laid on the effect must be attributed. I have ascertained this by a very latter. fimple experiment. I placed this colour on window-glass and fired it very strongly, and it did not change.

I then covered some parts of it with minium, and again ex-Experiments; posed it to the fire. The colour totally disappeared in those places where the red oxide of lead had been applied,

When I performed this operation on a larger scale in closed vessels, a large quantity of oxigen gas was disengaged.

This observation I think clearly proves the effect of oxided which shew that lead as a discolourer of glass: we see that it does not operate, colours glass as has been supposed, by burning combustible impurities in the glass, but by dissolving, discolouring, or volatilizing the oxide of iron which may affect its clearness.

#### Concerning the Yellows.

Yellows are colours that require much precaution in fabricat-Yellows are ing, on account of the lead they contain; which fometimes, oxides of lead by approaching to the metallic state, produces black spots.

The yellows of hard and tender porcelain are the fame. They are composed of oxide of lead, white oxide of antimony, and fand.

Oxide of tin is fometimes added; and when it is required and fometimes very lively and refembling the colour of the marigold, red ox-tin-ide of iron is added, the very deep colour of which disappears during the previous fusion they undergo, on account of the lead contained in this yellow. When these colours are once They are not made, they do not change; they disappear almost intirely in changeable, the porcelain fire.

Thefe

Not applicable to

A yellow for gials equal in beauty to the ancient.

These yellows cannot be applied to glass, they are opaque and muddy. That employed by the ancient painters on glass is, on the contrary, beautifully transparent, very brilliant, and of a colour approaching to gold. The processes they give indicate that it contained a mixture of filver; but when exactly followed, they afford nothing fatisfactory. Citizen Meraud, whom I have before quoted, has fucceeded in making it as beautiful as that of the ancient painters on glass, by employing muriate of filver, oxide of zinc, white clay, and the yellow' oxide of iron. These colours are applied to the glass simply ground, and without flux. The oxide of iron gives the yellow nearly the same tinge as it ought to have after the baking, and contributes, with the clay and oxide of zinc, to decompose the muriate of filver without difoxidating the filver itself. A powder remains after baking which does not penetrate the glass, and may be eafily cleared off.

This yellow, when employed in greater quantity, affords deeper shades, and produces a reddish yellow.

#### Concerning the Blues.

Blues from cobalt require nothing but care.

These are known to be obtained from the oxide of cobalt. Their preparation is known to every chemist. The superiority at Sevres, so justly reputed for the beauty of its blues, is owing merely to the care taken in its sabrication, and to the quality of the porcelain, which appears more proper to receive it on account of the violent fire it can support.

Oxide of cobalt is volatile.

I have observed one fact respecting the oxide of cobalt, which is perhaps not known to chemists: It is volatile in a violent heat; to this property must be attributed the bluish tint which the white (bordering upon the blue) always receives. I purposely put into the same case a white piece next to a blue; the side of the white piece that was turned towards the blue, became very bluish.

Blue grounds.

The blue of hard porcelain, prepared for what is called a blue ground by strong fire, is fused with feld-spar: the solvent for tender porcelain is silex, potash, and lead; it is not volatilized like the preceding, because the fire is much inferior to that of the hard porcelain.

These colours being previously fused, do not in the least change when applied.

Blue for glass.

The blues on glass are the same as for tender porcelain.

Concerning

Concerning Greens.

The greens employed in painting are made with the green Greens; from oxide of copper, or sometimes with a mixture of yellow and copper, blue. They must be previously melted with their flux; without this precaution they would become black; but they do not change after the first fusion.

They must not be treated with a violent fire, or they would not very fixed. totally disappear. The green grounds by strong heat are made with the oxides of cobalt and nickel, but it is only a brownish

green.

The bluish greens named sky-blue, formerly a colour very Blue-green. much in esteem, can only be used on tender porcelain; they always scale off from hard porcelain, because there is potash in

their composition.

These greens cannot be used on glass, because they afford a These greens do dirty colour: It is necessary to put a yellow on one fide and a not work well on more or less pale blue on the other, in order to produce a green. This colour may likewise be fabricated by mixing a blue with Blues for glass. the yellow oxide of iron. I hope to obtain a green from the oxide of chrome; and the experiments I have made promife to be attended with fuccefs. The pure chromate of lead, fixed on porcelain by means of a strong fire, has already afforded me a very deep and very fixed blue of confiderable beauty.

#### Concerning Biffres and Brown Reds.

These are obtained by mixtures of different proportions of Brown reds from manganese, brown oxide of copper, and the oxide of iron called mixed, umber. They are likewife previously fused with their solvents, so that they do not in the least change on tender porcelain; lead not having the fame action on the oxide of manganefe as it has on that of iron. I am convinced of this by an experiment fimilar to that I have already related.

This colour may be employed very well on glass.

The brown red grounds by strong heat, known by the name Grounds. of fonds caille, are made in the same manner. Feld-spar is their flux. There is no titanium in their composition, though generally afferted in books. Titanium was not known at Sévres when I first came to that manufactory. I have treated this fingular metal in various ways, and I never obtained any grounds but a flight obscure yellow, and very uncertain in its quality.

Concerning

answer on

Concerning the Blacks.

Blacks:

Black colours are the most difficult to be obtained very beautiful. There is no metallic oxide that singly affords a fine black. Manganese gives the best. Iron an opaque, dull, blistered black, which easily turns to red; the makers of colours have therefore combined several metallic oxides which singly do not afford blacks, and they have obtained a very beautiful colour, but it is subject to scale and become dull.

Oxides for compoling black.

These oxides are those of manganese, the brown oxides of copper, and a little of that of cobalt. Grey is obtained by suppressing the copper and increasing the quantity of flux.

Fine black at Sévres, The Sévres manufactory is the only one which has as yet produced beautiful blacks with a strong fire. This is more owing to the quality of the biscuit than to any peculiarity of process. It is by a mixture of blue with the oxides of manganese and iron, that they make this very brilliant black.

for glafs.

The blacks for opaque glasses are made the same as for painting, by giving different doses of solvent.

Mixtures of co-

I have shewn the principles of sabricating each principal colour: it is clear that by mixing these colours together, all possible shades may be obtained: and also, that care in the preparation, choice of materials, and just proportions of doses, must exhibit very sensible differences to the experienced eye of a painter. A knowledge of the composition of colours does not give the requisite care and neatness in making them up.

Recapitulation.

On recapitulating the facts I have just stated, in order to present them in a general view, we see, first, that amongst the colours usually employed for hard porcelain, one only is susceptible of change; namely, the carmine; and this may be replaced by the reds of iron, and then no colour changes.

Unchangeable fpecimens.

I have presented to the Institute an unbaked head made in this manner, and a painting of two roses, the one baked, the other in its first state. There was not any difference between them.

Secondly, That amongst the colours of soft porcelain and enamel several change considerably, particularly the reds of iron and gold with the yellows, greens, and browns. None have been substituted instead of them, this species of painting being almost abandoned.

Thirdly,

Thirdly, That feveral of these colours change likewise upon the glass by becoming persectly transparent, particularly the

yellows and violets.

Fourthly, That neither an additional calcination, nor a previous fusion, as has been suspected, will prevent them from changing. For this method alters the colours that change, and does nothing to the rest. The change which several colours undergo on tender porcelain and on glass, does not therefore relate to the nature of their composition, but rather to that of the body on which they are placed.

Consequently by suppressing the carmine of gold from the colours of hard porcelain, we shall have a series of unchangeable colours, which would be absolutely similar to those pre-

fented to the Institute in the year 6 \*.

#### IV.

Experiments and Observations on certain stony and metalline Substances, which at different Times are said to have fallen on the Earth; also on various Kinds of native Iron. By Edward Howard, Esq. F. R. S. From the Philosophical Transactions, 1802.

(Concluded from Page 101.)

Description of various Kinds of native Iron. By the Count de Bournon.

THE great number of particles of iron, in a perfectly me-Remarks on the tallic state, contained in the stone from Bohemia, and the said manner in which particles being so near each other, naturally lead to some re-have been steetions respecting the existence of native iron, which, by formed. many mineralogists, is still considered as problematical. Let us suppose for a moment, that these particles of iron were to

\* The preparation of colours for lively grounds by strong fire is under the care of Cit. Chanon, and that of the colours for painting to Cit. Meraud already quoted. It is to their care and intelligence in this chemical art, that the manufacture of Sévres is indebted for the preservation of its beautiful colours, which they have very much varied and improved.---B.

approach still more nearly to each other, so as absolutely to come into contact, and in that manner to form a kind of chain, folded upon itself in the interior part of the substance, and leaving a great number of cavities between the links of the chain fo folded. Let us then suppose, that the earthy substance with which these cavities are filled, being very porous, and having but a small degree of consistence, should (as may happen by a variety of causes) be destroyed. It is plain, that if fuch a destruction were to take place, the iron alone would remain; and, being thus left bare, it would appear in the form of a mass, more or less considerable, of a cellular texture, and as it were ramified; fuch a form, in short, as that in which most of the native irons we are acquainted with have been found. May it not be fair to attribute to fuch an origin. the native iron found in Bohemia, a specimen of which was prefented by the Academy of Freyberg to Baron Born, and which came, with the rest of his collection, into the hands of Mr. Greville? May not fuch also, notwithstanding the enormity of its bulk, be the origin of the mass of native iron found in Siberia, near Mount Kemirs, by the celebrated Pallas?

The stones examined by Mr. Howard contained nickel,

and so do the S. American iron. We have already feen, in the refults of the analyses made by Mr. Howard, of the various stones above described, that he constantly found a certain proportion of nickel mixed with the iron they contained. This circumstance recals to our notice the observations that were made by Mr. Proust, some time ago, respecting the mixture of nickel in the native iron of South America; and tends to give some additional support to the opinion hinted at in the foregoing paragraph.

The circumstances just mentioned, naturally gave to Mr. Howard, as well as to me, a defire to know whether the native iron from Siberia, and that from Bohemia, were also mixed with nickel. Mr. Howard, consequently, lost no time in proceeding upon this important investigation. The native iron of Siberia presents some very interesting peculiarities, and has often been referred to, but has not yet been properly described; it is therefore with great pleasure that I add the sollowing description of it, and of some other kinds of native iron, to the description I have already given of the various stones said to have fallen on the earth.

The Siberian native iron;

I feel the greater fatisfaction in doing this, as the noble collection of Mr. Greville contains two specimens of this iron, in perfect condition; one of which weighs feveral pounds, and was fent to Mr. Greville by Mr. Pallas himself: on this account, therefore, I enjoy an advantage that many of the authors who have spoken of this iron probably wanted.

. One of these pieces has a cellular and ramified texture, ana described, cellulogous to that of some very porous and light volcanic scoria: lar, and mixed with a yellowish this is the usual texture of the specimens of this kind of iron, green transpawhich are preserved in the prious mineralogical collections in rent matter. Europe. When it is attentively examined, there may be perceived in it, not only empty cells, but also impressions or cavities, of greater or left depth, and fometimes perfectly round, which appear evidently to be the refult of the compression of hard bodies, which were fituated there, and which, when they came away, left the furface of these cavities quite smooth, and having the luftre of polished metal. Here and there, in some of these cavities, there remains a transparent substance, of a yellowish green colour, of which I shall treat more particularly, when I come to the description of the second of the specimens above mentioned. It is very clear, that the cavities here spoken of owe their existence to this transparent fubstance; and that the polish of the cavities arises merely from the compression of the said substance. and is the natural consequence of its surface having been in persect contact with that of the iron.

This iron is very malleable: it may be easily cut with a This iron very knife; and may be as eafily flattened or extended by means malleable. of a hammer. Its specific gravity is 6487; which, however, is very much under that of iron which has been merely melted, and has not been forged. The specific gravity of the native iron of Bohemia, which is nearly as malleable and as eafy to be cut, is still less: I found it not to exceed 6146. This low degree of gravity, appears to be owing partly to the oxidizement of the furface of the iron, and partly to there being, in the interior part of its substance, a number of small cavities. which are often rendered visible by fracture, and which have their furfaces also oxidized. The fracture of this iron, pre-Its fracture is fents the fame shining and filvery white colour as the common white, east iron, known by the name of white cast iron; but its grain is much fmoother and finer: it is also much more malleable when cold. Bergman fays that this iron is brittle, when heated to a red heat. I have frequently tried it in that state, but it is not and red-short.

and have confrantly found it to be malleable. The fame remark may be applied to the native iron from South America; and also to that from Senegal.

Another fpecimen; more compact,

The fecond of the two specimens mentioned above, and which weighs feveral pounds, prefents an afpect that differs, in some respects, from that of the preceding specimen. The most confiderable part of it forms a folid compact mass, in which there is not to be perceived the smallest appearance of pores or cavities; but there arises upon its surface, a kind of ramified or cellular part, fimilar, in every respect, to the specimen already defcribed, and every where completely connected with the fubstance of the mass itself.

confisting of iron ter, which fills ets cavities.

If the compact part of this piece is examined with attention, and the greenish it will be perceived, that it is not entirely composed of iron in the metallic state, but that it is mixed with nearly an equal quantity of the transparent substance of a yellowish green colour, (fometimes also of a greenish yellow,) already spoken of in the description of the other specimen. This substance is mixed with the iron, in fuch a manner, that if the whole of the former could be removed, the remaining part would confift merely of iron in the metallic state, and would present the fame cellular appearance as the preceding specimen, and the ramified or cellular part of the specimen now described.

The green stony part looks like glafs :

This flony part, separated from the iron, appears in the form of small nodules, generally of an irregular shape, but fometimes nearly globular: they have a perfectly smooth and shining surface, so as very often to present the appearance of fmall balls of glass; a circumstance that has led many persons to suppose them the result of a real vitrification. Some of these nodules have several irregular facets, produced by the compression of the iron in which they were inclosed; but I have never observed in them, any appearances that could lead me to suspect they had the slightest tendency whatever to alfume a determined crystalline form.

is always tranfparent; cuts glass, but not quartz; fracture conchoidal; electric by friction;

This substance is always more or less transparent. fufficiently hard to cut glass; but has no effect upon quartz. It is very brittle: its fracture is usually conchoid; but I could not perceive that it broke in any particular direction, in fuch a way that I could confider the fracture as a natural one. becomes electric by friction. Its specific gravity is from 3263 to 3300. It is very refractory: I kept it, for some time, exposed

posed to a degree of heat sufficiently strong to oxidize, to a considerable depth, the iron crucible in which it was placed, not suffile in an without its having undergone any alteration, except that of having acquired a greater degree of intensity in its colour. Its transparency was not at all diminished. I think, therefore, there is not the smallest reason to allow any probability to the opinion that it ought to be considered as a kind of glass.

Of all substances hitherto known, that with which it seems it resembles the to have the greatest analogy, is the peridot, (the chrysolite of peridot; Werner,) to which some mineralogists have referred it. The result of Mr. Howard's analysis of it, is nearly the same as that

of the analysis of the peridot, made by Mr. Klaproth.

The hardness and infusibility of this substance are nearly the in hardness and fame as those of the period; but it feems to have a rather less infusibility. degree of specific gravity: that of two very perfect crystals of peridot, I found to be from 3340 to 3375. The crystalline forms of the substance here described, if ever we should be able to determine them, would clear up our doubts respecting the analogy between the two fubftances. If we confider the compact part of the specimen now treated of, particularly the strong connection that appears to exist between the iron and It is curious that, the transparent substance, and the great resistance we expe-it should be de-fructible by the rience when we attempt to separate them, we cannot help weather; being furprised, that almost all the specimens of this mass of metallic iron that have been brought to Europe, are in the cellular state already described, owing apparently to the total, or almost total, destruction of the transparent substance. But, befides the fragility of this substance, the specimen in question helps very much to explain the above cirumstance, inasmuch as many of the nodules of the transparent substance belonging to it, are in a state of real decomposition. In that state, they are changed into a white opaque substance, which, upon being lightly pressed or squeezed between the singers, crumbles into a gritty dry powder. This decomposition may be observed to but it undoubthave taken place in various degrees: in many of the nodules, edly is fo. the fubstance is merely become friable, without being much altered in its appearance: whereas, some of those which are in a state of complete decomposition, are of an ochreous reddish yellow colour; it is, however, easy to distinguish that this colour does not belong to them, but is owing only to the oxiidizement of the adjacent particles of iron.

From

This fact exture of native iron, &c.

From the above observations, it will not be difficult to conplains the firuc-ceive the possibility of the total, or nearly total, destruction of the transparent substance; and also, the appearance the pieces of iron must naturally present, when deprived of it. I cannot help observing likewise, that there appears to exist a very interefting analogy, between thefe transparent nodules and the globules I described as making part of the stones said to have fallen on the earth. This analogy, though not a very firong one, may lead us to suppose that the two substances are similar in their nature, but that the globales are less pure, and contain a greater quantity of iron.

The Bohemian iron in fome measure resembles the Siberian.

The native iron from Bohemia is a compact mass, fimilar to the compact part of the large specimen of iron from Siberia, which has just been described: like that, also, it contains a number of globular bodies or nodules; but they are not in fuch great proportion as in the Siberian iron. They are befides perfectly opaque, and very much refemble the most compact of the globules belonging to the stones said to have fallen on the earth.

#### EXAMINATION OF THE IRON FROM SOUTH AMERICA.

S. American iron contains 10 per cent. of nickel.

I have already observed, that my experiments coincided with those of Mr. Proust. He obtained 50 grains of sulphate of nickel, from 100 of this mass. The process I have so frequently mentioned, yielded me 80 grains of oxide of iron from 62 of the metal; which indicates about 7 of nickel, or about 10 per cent.

#### EXAMINATION OF THE SIBERIAN IRON.

100 grains of this iron, gave 127 of oxide of iron: hence, Siberian iron 17 it should contain about 17 per cent. of nickel. per cent. &c.

The yellow mat-

The yellow substance belonging to this iron, was analyzed in the same way as the globular bodies, and the earthy parts, of the stone from Benares.

The proportions, refulting from the analysis of 50 grains, and from fome previous experiments on other particles, were.

Silica

Silica -	i u		i Jiv		-		27
Magnesia -	100	-	Titu	0 4		-	131
Oxide of iron	1 50		-14				- 8 <u>I</u>
Oxide of nickel				-		- 1	- 13t1 - 1
The service			13				-
and Pilotty			13.7		1		494

#### EXAMINATION OF THE BOHEMIAN IRON.

261 grains of this metal, left about 11 grain of earthy mat-Bohemian iron ter, infoluble in nitric acid; and, by ammonia, afforded 30 17 per cent. do. grains of oxide of iron, inducing an estimation of nearly 5 of nickel.

#### EXAMINATION OF IRON FROM SENEGAL, BROUGHT BY GENERAL O'HARA, AND GIVEN TO ME BY MR. HATCHETT.

In this experiment, 199 grains of oxide were produced Senegal iron-5 from 145 grains of metal; hence, there may be an estimation or 6 per cent. of 8 grains in 145, or between 5 and 6 per cent. of nickel.

It will appear, from a collected view of the preceding pages Summary of the and authorities, that a number of stones afferted to have fallen characters of the under similar circumstances, have precisely the same characters. the earth. The stones from Benares, the stone from Yorkshire, that from Sienna, and a fragment of one from Bohemia, have a relation

to each other not to be questioned.

1st. They have all pyrites of a peculiar character.

2dly. They have all a coating of black oxide of iron.

3dly. They all contain an alloy of iron and nickel.

4thly. The earths which serve to them as a fort of connecting medium, correspond in their nature, and nearly in their proportions.

Moreover, in the stones from Benares, pyrites and globular bodies are exceedingly distinct. In the others they are more or less definite; and that from Sienna had one of its globules transparent. Meteors, or lightning, attended the descent of the Meteors or lightstones at Benares, and at Sienna. Such coincidence of cir-ning accompacumstances, and the unquestionable authorities I have ad-nying them. duced, must, I imagine, remove all doubt as to the descent of these stony substances; for, to disbelieve on the mere ground of incomprehensibility, would be to dispute most of the works of nature.

Pyrites, oxide of iron, nickel, earths correfpondent.

Native irons.

Respecting the kinds of iron called native, they all contain nickel. The mass in South America is hollow, has concavities, and appears to have been in a foft or welding state, because it has received various impressions.

The Siberian iron has globular concavities, in part filled with a transparent substance, which, the proportional quantity of oxide of iron excepted, has nearly the composition of the globules in the stone from Benares. O Valla 110433

The iron from Bohemia adheres to earthy matter studded with globular bodies.

The Senegal iron had been completely mutilated before it came under my examination.

From these facts, I shall draw no conclusion, but submit the following queries.

Have thefe all and from meteors?

1st. Have not all fallen stones, and what are called native the same origin? irons, the same origin?

2dly. Are all, or any, the produce or the bodies of meteors? And, lastly, Might not the stone from Yorkshire have formed a meteor in regions too elevated to be discovered?

Specimens of the Benares and Yorkshire stones have been deposited, by the President, in the British Museum.

Experiments and Observations on the Heat and Cold produced by the Mechanical Condensation and Rarefaction of Air \*. By JOHN DALTON.

Well known facts that the thermometer rifes in condenfed and falls in rarefied air.

IF a thermometer be inclosed in a receiver and the air suddenly condenfed, the thermometer rifes a few degrees above the temperature of the atmosphere; and if the air be exhausted from a receiver inclosing a thermometer, the mercury finks a few degrees immediately; but in both cases after some time it refumes its former station. These facts are well known to philosophers of the present age, but they do not all agree in the explanation of them. Thinking the subject worthy of elucidation. I was induced to inftitute a feries of experiments for the purpose, which I apprehend have led to a clear demonfiration of the cause of the phenomena, and moreover makes

<sup>\*</sup> Manchester Memoirs, V. 515.

the facts themselves appear in a somewhat different point of view from what they are seen in at the first moment.

One circumstance is very remarkable, that whether the This rise and mercury rises or falls in these instances, it is done very rapidly; fall are very rapidly; whereas in the open air, if a thermometer be only two or three degrees above or below the temperature, it moves very slowly.

This seems to have suggested to every one the idea, that the whence it was elasticity of the glass bulb of the thermometer has a principal supposed to arise from pressure on thare in producing the effect, by causing the bulb to yield a the glass bulb; little to the pressure of the air. It has however been found upon trial that the same effects take place whether the thermometer is sealed or not. My experiments accord with this, have but it happens ing made a thermometer and left it unsealed for the express whether the tube be sealed or purpose; in all the experiments with condensed and rarefied not.

air, there was no sentible difference observed to arise from the inequality of pressure on the external and internal surfaces of the bulbs, the sealed and open thermometers varying the same in kind and also in degree, except from circumstances to be

It being certain then that a real change of temperature takes Confeq. a real place, it remained to determine the quantity and manner of change of temperature takes that change. Having chosen a small and consequently sensitive takes ble thermometer, with a scale of degrees sufficiently large to admit of distinguishing one tenth of a degree, I proceeded to ascertain several facts experimentally.

### EXPERIMENT 1.

noticed hereafter.

Took a receiver, the capacity of which was about 120 cu-Exp. I. A very bic inches, and suspended the thermometer with its clear bulb small thermometer in the central part of it; then letting the whole acquire the receiver sell by temperature of the room, which was without a fire, I ex-exhaustion, and hausted the air and afterwards restored it, marking the effects the contrary apeny support the thermometer. The medium of several trials nearly agreeing with each other was as under:

The thermometer in the air of the room flood at - 36°.8

funk upon exhaustion to - - 34 .7

rose when the air was restored to 38 .9

The fuddenness of the fall and rise puzzled me most: after reslecting upon it for some time, I conjectured that the real change of temperature of the air or medium was much greater Vol. III.—November, 1802.

whence it was inferred, that the real change of temperature was very great; but not very effective because to momentary.

than the thermometer indicated, but that the inequality existed only for a few seconds of time, because the receiver, &c. immediately impart heat to or abstract it from so small a quantity of air as 120 cubic inches, which are only equal to 40 grains in weight. The phenomena of the thermometer feemed very well to accord with the supposition of great heat or cold acting upon it for a few feconds only.

#### EXPERIMENT 2.

Exp. 2. A large and a small thermometer were cooled differently.

Pursuing this idea, I imagined that if two thermometers whose bulbs were very unequal in magnitude were inclosed together, the smaller bulb ought to give the greater variation: accordingly I inclosed two, the diameters of their bulbs being .35 and .65 of an inch respectively; and having exhausted the air and restored it again repeatedly in succession, and found a mean of the variations, that of the small bulb was 2°.8, and that of the large, 20.20 and a ligable helicited a limit and in home and affection was every home to an amount

#### EXPERIMENT 3.

Exp. 3. And the fame therm. of heat.

Repeated the exhaustion with the small thermometer inwas more cooled closed in three different circumstances successively; 1st with the more remote the bulb in the centre of the receiver; 2d with the bulb restfrom conductors ing on the wet leather of the plate; and 3d with the bulb refting against the side of the receiver.

1st Case-	-Reduced	by exhauftion	n -			2°.45
2d Cafe	•	A TACEL	2-14	-	1	2.15
3d Cafe	1 - mm	accepte who		PIVE S	-1 4 5	1.2
Ift Cafe-	-Raifed by	reftoring the	e air	All big	4	4 .05
		di seditar se				
		vi aw- 'aidi				
		ati long top of	JID ST TIT	8 5 19	11	haufted

#### EXPERIMENT 4.

Exp. 4. A cubic inch of water was fenfibly cooled.

Inclosed a wine glass with about a cubic inch of water in it, containing the bulb of a thermometer, in a receiver; and, exhausting the air, the thermometer funk half a degree fuddenly, and then continued flationary; upon restoring the air it fuddenly rofe half a degree.

From all which it follows that the real change was much

All these experiments confirmed my conjecture of a much greater degree of heat and cold being produced in these cases than the thermometer points out, but that its continuance is fo short as not to effect a material change in the temperature greater than of the mercury. The following experiments were made to them by the thermometer. afcertain what may be the real degree of heat and cold generated in those operations.

#### EXPERIMENT 5.

The same receiver and small thermometer as above being Exp. 5. The used, I sound the exhaustion was effected by working the time of substitution one minute. The thermometer sunk nearly 2° in the than of subsefirst half minute, and the remainder, a few tenths of a degree, quent rise in the in the latter half minute. The operation being stopped, and vacuum; things remaining in the fame state, it required some minutes of time before the thermometer recovered one degree of the heat loft. Upon opening the cock, the receiver filled with air in five feconds, and the greatest velocity of the rising mercury was about the end of that time. The rifing continued and the time of for 30 or 40 feconds from its commencement, but  $\frac{3}{4}$  of the duction of the effect were produced in the first 10 seconds. The greatest external air velocity of the rifing mercury is 1° in  $3\frac{1}{2}$  feconds. After the much less than thermometer had attained its utmost height, it began to fall quent fall. again at the rate of 1 of a degree in a minute.

#### EXPERIMENT 6.

Took the fame thermometer and heated it to 50° above the Exp. 6. By temperature of the air, then let it be cooled by the medium of heating the air, and it began to fall at the rate of 1° in 3½ feconds.

The two last experiments feem to prove that when air is let the air, it was in to the receiver in the ordinary way, an increase of temperaelevation of 50°. ture of 50° is produced in the medium within the receiver for 3½ was required to feconds. This high temperature is reduced in a few seconds by produce as rapid an effect as in the receiver and furrounding bodies, to their own temperature. the former ex-

letting it cool in periment. Such therefore

#### EXPERIMENT 7.

On condensed Air.

Took a large spherical glass receiver, the capacity of which Exp. 7. Therwas fomething more than twice that of the former (above one mometer in congallon), and fuspended a thermometer in the centre of it, of a fuddenly when larger bulb than that before used; the receiver had a brass cap the air was let and ftop-cock adapted to it: then doubled the denfity of the in. air within it by a condenser. The thermometer rose 20 or more, Let out the air fuddenly and the thermometer imme-

was the real rife.

M 2

diately

diately funk each time from 30 to 30.5; at the same time an exceedingly denfe mist was produced in the receiver, which foon fubfided as well to see you have my those wall will

Suspecting that aqueous vapour, which always exists in the atmosphere, and is liable to assume the liquid or aërial form according to circumstances, might be the principal agent in the production of heat and cold by condensation and rarefaction, I thought that an increase of it might produce a greater effect, and that cold air, which contains less vapour, might have a less effect. The reverse however was the fact, as appears by the following,

#### EXPERIMENTS 8 and 9.

Exp. 8 and 9. Thefe effects did not arise from the water fuspended in the

was less when the vapours were most abun-

In a cold morning last winter when the air was clear and the thermometer without flood at 20°, I took the receiver and condenser into the open air, and let them stand for 15 minutes to acquire its temperature; then repeatedly condenfed the air to a double denfity, and fuddenly liberated it again. medium of five trials the mercury fell 3°.3 on opening the. For the change cock. The vapour precipitated was whiter than usual and not nearly fo denfe.

Again, took the receiver and condenfer into a dyer's flove where the temperature was about 100°, and the air abounded with vapour in a transparent state: after some time, condensed the air and liberated it as before, when on a medium of five trials the mercury funk only 30, and a very copious mist was precipitated, so dense that one could but just distinguish the degree of the thermometer through it.

These experiments shew that the greater the quantity of vapour condensed the less is the change of temperature; and that confequently, if air was entirely free from vapour, the change of temperature would be a maximum. Indeed this is clearly confistent with the known law, that when vapour is condenfed, heat is given out. Any process to cool the air must be retarded by the condensation of part of the vapour it contains. Suppose for instance that a portion of the atmosphere contained to of its weight of aqueous vapour, and that 3 of this vapour were condensed by 50° of cold; that is, 700 of the whole elastic mass was converted into water; then the heat given out would be fufficient to raise the temperature of the remaining mass of air and vapour 6 or 8°, which sufficiently

This might be inferred from the known theery.

ciently accounts for the small difference observed in the results upon warm vapoury air and cold dry air. Hence vapour, fat from producing the change of temperature in question, tends to diminish the effect.

If any doubt remained with me respecting the real change of temperature that takes place in the operations related above, it was completely removed by the refults of the two following experiments.

#### EXPERIMENT 10.

Inclosed a small graduated glass tube of 1 of an inch inter- A small manonal diameter, and 10 inches long, with a fhort column of meter was inmercury in it, in the large receiver; the tube was fealed at one end and open at the other, fo that a portion of given capacity was confined by the mercurial column, which was near the open end of the tube, and fubject to rife or fall by any variation of elasticity of the air on either fide, being a proper manometer: then doubled the denfity of the air in the re-the denfity of ceiver, and opening the stop-cock, the mercurial column soon the surrounding air was soubled; ran up to its sommer station, but instantly turning the cock then restored by again, the mercurial column returned or fell down gradually a momentary for 5 or 10 feconds, to the amount of nearly  $\frac{1}{10}$  of the whole from without. aërial column, and then became stationary. Again opening The manometer the cock, a quantity of air rushed out, and the mercury if the air in the refumed its original station. These effects were always the receiver had fame, on a repetition of the experiment.

#### EXPERIMENT 11.

Let the mercurial column of the manometer down by a Exp. 11. The wire to 4 of the length of the tube from the fealed end; then manometer as before, but in exhausted 3/4 of the air from the receiver, which was seen by rarefied air: the mercury rising to the top of the tube; and upon opening the cock the mercury fell to its former station, but then suddenly turning the cock, the mercury gradually rose for the The air let in fpace of 5 or 10 feconds to more than To of its original height showed figns of above the flationary point, and remained there till the cock was opened: after which it refumed its proper station.

The phenomena in the two last experiments can be ex-Fuller explanaplained only on the following principle:—The air in the re- tion. The air in the manomeceiver and in the manometer is subject to a like degree of ter recovers the rarefaction and condensation in those experiments, or nearly common temperature some for

gradually abforbed heat and become more

receiver: whence in the former case the gradual heating of the air in the receiver renders in the other cafe.

than that in the fo. When the equilibrium of heat in the air is disturbed by the operations of condensation and rarefaction, it is restored in the manometer instautly by reason of the contiguity of the glass to the air; but in the large receiver it requires a sensible time of 10 feconds or more to restore the equilibrium throughout its pressure from the whole internal capacity. It is this restoration that inon the mercury; creafes or diminishes the elasticity of the air confined in the and the contrary receiver, and thereby causes the retrogradation of the mercurial column. Now I have found by former experiments, that a change of 50° in temperature effects a change of to nearly, in the capacity or bulk of air. It follows therefore that in the case of restoring the equilibrium in condensed air, about 500 of cold is propuced; and in letting in air to an exhausted receiver fomething more than 50° of heat is produced. fmall difference feems to arife from this, that the condensation of vapour in the former case diminishes the effect, and in the latter, if any there be, increases the effect, that would arise from operating upon purely dry air. The experiments and observations hitherto related go prin-

The theory of thefe effects refts on the difcity of air for heat; viz. less the greater the denfily.

cipally to ascertain facts without any reference to the theory ference of capa- of them: This however may be given in a few words, and is the fame that is afcribed to Mr. Lambert by Messrs. Saussure and Pictet and by them adopted. He conceives that a vacuum has its proper capacity for heat, the same as air, or any other fubstance; and that the capacity of a vacuum for heat is greater than that of an equal volume of atmospherical air; also that the denser air is, the less is its capacity for heat: upon these principles the phenomena are easily referable to that class of chemical facts where heat and cold are generated by the mixture of two different bodies. If this theory be right, and I think there is little doubt of it, we may hence be led into a train of experiments, by which the absolute capacity of the capacities of a vacuum for heat may be determined; and likewise the capacities of the different gases for heat, by a method wholly new: but this must be left to future investigation.

Whence new experiments may elastic fluids.

### become VI.

An Account of a Method of copying Paintings upon Glass, and of making Profiles by the Agency of Light upon Nitrate of Silver. Invented by T. WEDGWOOD, Efq. With Observations by H. DAVY. From the Journals of the Royal Institution, I. 170.

WHITE paper, or white leather, moistened with solution White paper, of nitrate of filver, undergoes no change when kept in a dark act, wetted with nitrate of filver, place; but, on being exposed to the day-light it speedily becomes black by changes colour, and, after passing through different shades of light: grey and brown, becomes at length nearly black.

The alterations of colour take place more speedily in pro- with different portion as the light is more intense. In the direct beams of the speed as the light is stronger; e.g. fun, two or three minutes are sufficient to produce the full 2 minutes in effect. In the shade several hours are required, and light trans-sunshine. mitted through different coloured glasses, acts upon it with different degrees of intenfity. Thus it is found that red rays, Coloured rays or the common fun-beams passed through red glass, have very act with different intensities. little action upon it: yellow and green are more efficacious; but blue and violet light produce the most decided and powerful effects \*.

The

\* The facts above mentioned are analogous to those observed long ago by Scheele, and confirmed by Senebier. Scheele found, that in the prismatic spectrum, the effect produced by the red rays upon muriate of filver was very faint, and scarcely to be perceived; whilst it was speedily blackened by the violet rays. Senebier states, that the time required to darken muriate of filver by the red rays, is 20 minutes, by the orange 12, by the yellow 5 minutes and 30 feconds, by the green 37 feconds, by the blue 29 feconds, and by the violet only 15 seconds. - Senebier sur la Lumière, Vol. III. p. 199.

Some new experiments have been lately made in relation to this subject, in consequence of the discoveries of Dr. Herschel concerning the invisible heat-making rays existing in the solar beams, by Messrs. Ritter and Bockmann in Germany, and Dr. Wollaston in England.

It has been ascertained by experiments upon the prismatic spectrum, that no effects are produced upon the muriate of filver by the invifible

Hence the light that passes through a paintcause various tinges on the paper, &c. and afford a copy;

The confideration of these facts enables us readily to underfland the method by which the outlines and shades of paintings ing on glass will on glass may be copied, or profiles of figures procured, by the agency of light. When a white furface, covered with folution of nitrate of filver, is placed behind a painting on glass exposed to the solar light; the rays transmitted through the differently painted furfaces produce diffinct tints of brown or black, fenfibly differing in intenfity according to the fluxdes of the picture, and where the light is unaltered, the colour of the nitrate becomes deepest.

or profiles may be taken from fradows.

When the shadow of any figure is thrown upon the prepared furface, the part concealed by it remains white, and the other parts speedily become dark.

For copying paintings on glass, the solution should be applied on leather; and, in this case, it is more readily acted upon than when paper is used.

The tinge is permanent.

After the colour has been once fixed upon the leather or paper, it cannot be removed by the application of water, or water and foap, and it is in a high degree permanent.

The copies thus made are changeable by daylight;

The copy of a painting, or the profile, immediately after being taken, must be kept in an obscure place. It may indeed be examined in the shade, but, in this case, the exposure should be only for a few minutes; by the light of candles or lamps, as commonly employed, it is not fenfibly affected.

and this imperfection has not yet been remedied.

No attempts that have been made to prevent the uncoloured parts of the copy or profile from being acted upon by light, have as yet been successful. They have been covered with a thin coating of fine varnish, but this has not destroyed their fusceptibility of becoming coloured; and even after repeated washings, sufficient of the active part of the saline matter will still adhere to the white parts of the leather or paper, to cause them to become dark when exposed to the rays of the fun.

Copies of tex-

Besides the applications of this method of copying that tures thus made. have been just mentioned, there are many others.

> invisible heat-making rays which exist on the red side, and which are least refrangible, though it is powerfully and distinctly affected in a space beyond the violet rays out of the visible boundary. See Annalen der Physik, siebenter Band, 527 .- D.

will be useful for making delineations of all such objects as are possessed of a texture partly opaque and partly transparent. The woody sibres of leaves, and the wings of infects, may be pretty accurately represented by means of it, and in this case it is only necessary to cause the direct solar light to pass through them, and to receive the shadows upon prepared leather.

When the folar rays are passed through a print and thrown Prints not very upon prepared paper, the unshaded parts are slowly copied; distinct. but the lights transmitted by the shaded parts, are seldom so definite as to form a distinct resemblance of them by producing different intensities of colour.

The images formed by means of a camera obscura, have The images of been found to be too faint to produce, in any moderate time, the camera are an effect upon the nitrate of filver. To copy these images process, was the first object of Mr. Wedgwood, in his researches on the subject, and for this purpose he first used the nitrate of filver, which was mentioned to him by a friend, as a substance very sensible to the influence of light; but all his numerous experiments as to their primary end proved unsuccessful.

In following these processes I have found, that the images of Those from the small objects, produced by means of the solar microscope, may are better. be copied without difficulty on prepared paper. This will probably be a deful application of the method; that it may be employed successfully however, it is necessary that the paper be placed at but a small distance from the lens.

With regard to the preparation of the folution, I have To make the found the best proportions those of 1 part of nitrate to about 10 of water. In this case, the quantity of the salt applied to the leather or paper, will be sufficient to enable it to become tinged, without affecting its composition, or injuring its texture.

In comparing the effects produced by light upon muriate of The muriate is filver, with those produced upon the nitrate, it seemed evident, better than the that the muriate was the most susceptible, and both were more licacy, readily acted upon when moist than when dry, a fact long ago known. Even in the twilight, the colour of moist muriate of silver spread upon paper, slowly changed from white to faint violet; though under similar circumstances, no immediate alteration was produced upon the nitrate.

The

but is less soluble. The nitrate, however, from its folubility in water, possesses an advantage over the muriate: though leather or paper may, without much difficulty, be impregnated with this last substance, either by diffusing it through water, and applying it in this form, or by immersing paper moistened with the solution of the nitrate in very diluted muriatic acid.

Caution.

To those persons not acquainted with the properties of the salts containing oxide of filver, it may be useful to state, that they produce a stain of some permanence, even when momentarily applied to the skin, and in employing them for most-tening paper or leather, it is necessary to use a pencil of hair, or a brush.

Something like a dying process may perhaps be effected here.

From the impossibility of removing, by washing, the colouring matter of the salts from the parts of the surface of the copy which have not been exposed to light, it is probable that, both in the case of the nitrate and muriate of silver, a portion of the metallic oxide abandons its acid to enter into union with the animal or vegetable substance, so as to form with it an insoluble compound. And, supposing that this happens, it is not improbable but that substances may be found capable of destroying this compound, either by simple or complicated affinities. Some experiments on this subject have been imagined, and an account of the results of them may possibly appear in a future number of the Journals. Nothing but a method of preventing the unshaded parts of the delineation from being coloured by exposure to the day is wanting, to render the process as useful as it is elegant.

## VII.

Observations and Experiments upon Oxigenized \* and Hyperoxigen-. ized Muriatic Acid; and upon some Combinations of the Muriatic Acid in its Three States. By RICHARD CHENEVIX. Efy. F. R. S. M. R. I. A. From the Philosophical Transactions for 1802.

WHEN Mr. Berthollet made known the combination of Inference of what was then called oxigenated muriatic acid with potath, the fo called oxhe gave as his opinion, that the proportion of oxigen, rela- igenated muriate tively to the quantity of acid, was greater in the falt than in of potash conuncombined oxigenized muriatic acid. This conjecture was gen than is in fairly founded upon the observation, that, in his mode of pre-the uncombined paring this falt, a large portion of common muriate was formed acid. in the liquor, along with the hyperoxigenized muriate. memoir which he published in the year 1788, is the last with which I am acquainted, upon this subject. It does not contain any thing that, confidering the accuracy which is now required in experiments, amounts to a demonstration of the relative proportions of oxigen, in oxigenized and hyperoxigenized muriatic acids. Unfortunately, this chemist has not purfued his refearches any farther; although, from his own words, we had every reason to hope that they would have been continued.

In the Système des Connoissances Chimiques of Mr. Fourcroy, Fourcroy's acwe find a fummary of the experiments that had preceded the count. impression of his work, together with the following sentence: "Tous les muriates suroxigénés sont décomposés par les " acides, fouvent avec une violente décrépitation, avec un " dégagement de vapeur jaune verdâtre, et une odeur très-" forte. Cette vapeur est de véritable acide muriatique sur-" oxigéné. Elle est lourde, tombe en goutellettes d'un jaune " vert, et forme des stries comme de l'huile, sur les corps auxquels elle adhère." This affertion carries no confirmation

\* I have preferred this word to oxigenated, because ate is the appropriate termination of certain falts formed by acids in ic. further remarks upon this subject will be made in a work now in the press, intitled, Remarks upon Chemical Nomenclature.

along

hyperox. mur. acid not yet proved.

along with it; and does not amount so near to proof as the The existence of position of the former chemist: so that, in fact, the existence of hyperoxigenized muriatic acid, and of its combination with potash, rests, at present upon the conjecture of Mr Berthollet; a conjecture however which, as well as his whole differtation upon the subject, bears all the marks of genius which so strongly characterife every production of that fagacious philosopher. Some notice has been taken of other faline combinations, formed by caufing a current of oxigenized muriatic acid to pass through solutions of the alkalis, or earths, or by otherwise combining them. Meff. D'Olfus, Gadolin, Van-Mons, Lavoisier, and others, have slightly mentioned some of these combinations. But, with the exception of Mr. Berthollet, I know of no chemist who has approached so near to the real state of the combination of muriatic acid and oxigen with potash, as Mr. Hoyle, of Manchester. The true nature of this falt, however, is one of those things which many persons have credited without proof; and which many others have been on the eve of discovering.

New obs. and expts. in proof of that state.

I shall now proceed to lay before the Society, an account of the observations and experiments which have led me to conclude, that muriatic acid does exist in the form of oxigenized and hyperoxigenized muriatic acid, as announced in the title of the present communication; and that, in either state, it is capable of entering into faline combinations.

Order of narrative.

With this view, I shall describe,

1st, The means by which I think I have succeeded, in aftertaining the constituent parts, as well as the proportions, in oxigenized and hyperoxigenized muriatic acid.

2dly. I shall mention some of the combinations of the muriatic acid, in its three states.

In treating upon the first of these objects, I must in some measure anticipate the second; and must suppose some things known, which are hereafter to be described. This inconvenience is inevitable; as the natural order of things leads me to treat of the acid, before I confider the bodies into the composition of which it enters.

Hyperox. mur. of potash over a lamp decrepitated, melted, remained in fufion, and loft

I exposed to the heat of a lamp, 100 grains of hyperoxigenized muriate of potash. It decrepitated gently, and in a short time melted. After remaining in fusion nearly an hour, I allowed it to cool; it crystallized as formerly, and had lost

2,5 per

2,5 per cent. I increased the heat to redness, in a surnace. 0,025. Red The salt boiled with a violent effervescence, and rapid disheat caused engagement of gaseous sluid, together with a thin white loss of 0,5. vapour, and then sunk suddenly into a white spongy mass.

The loss of weight usually varied from 42 to 48 or 50 per cent.

I put 100 grains into a coated glass retort, luted to a small Distillation of hyperox. mur. and perfectly dry receiver, having a tube communicating with of potafh, a glass bell in the pneumatic tub. The fire had not been lighted very long, when a flight dew began to line the infide of the receiver; and, as foon as the retort was nearly red-hot, afforded oxigen; a difengagement of gas, fo fudden as almost to be explosive, took place. A quantity of thin white vapour arofe, which afterwards was deposited, in the form of a white sublimate, in the receiver and the tube. When the extrication of gas had ceased, the apparatus was allowed to cool. The gas, with the usual corrections of temperature and pressure, measured 112,5 cubic inches, = 38,3 grains. The 2,5 mentioned above, as and left muriate. the loss of this falt at a low heat, were water. 53,5 remained in the retort; and the white sublimate in the tube and receiver amounted to 5. Sale and a constant, but, and a constant

The products of this operation were therefore,

Water	Lan	12	2;5
Oxigen	19		38,3
Salt in the tube and receiver -	. /	- 1	5
Salt remaining in the retort			53,5
A STATE OF THE STA	,		
Hart to a great remaining the		417	99,3.

To find the proportions of oxigen and muriatic acid, in hyperoxigenized muriatic acid, it now only remains to determine the fum of the quantities of muriatic acid, contained in the The refidual 53,5 of the retort, and the 5 of the tube and receiver. The muriate was employed to preciple to precipitate corresponding to pitate nitrate of 18,21; and the 5, a precipitate corresponding to 1,76; in all, filver, and its 20 of the muriatic acid. Therefore, 38,3 of oxigen, and 20 of acid inferred muriatic acid, combine to form 58,3 of hyperoxigenized murfrom the precipitate acid; or, 100 of hyperoxigenized muriatic acid contain, pitate, within a fraction.

Oxigen	-	. ,-	- 65
Muriatic acid		p erman	- 35
n Changian and Lon		12 8 11 17 17	
Dailes emg.			100.

276 747

The hyperox. muriate therefore contained nearly I part common acid, 2 parts potafh, and 2 parts oxigen.

And the elements of hyperoxigenized muriate of potash, should be thus stated:

Oxigen - Muriatic acid	38,3   hyperox 20   muriati	igenized }	58,3	
10.36	Potash	11 1/15	39,2	
	Water	-	- 2,5.	
	0, 8.4	1 1773 0	100,0.	

Muriates lose a little acid by ignition.

It may be observed, that the 53,5 of the retort did not yield the same proportion of acid as the 5 of the tube and receiver. The fact is, that all muriates lofe a little of their acid at a red heat, as I shall prefently mention more particularly; and the fmall lofs was, in all probability, owing to a portion of acid difengaged by the heat to which the falt was necessarily exposed during the operation. Having thus afcertained the proportion of oxigen in hyper-

To determine the oxigen added to convert muriatic into oxyg mur. acid.

Ox. mur. acid

oxigenized muriatic acid, by means of its combination with potash, a ready method occurred to arrive at the knowledge of that contained in oxigenized muriatic acid. For this purpose, I disposed in the following manner, a Woulse's apparatus, confifting of three bottles, and connected with the pneumatic tub. In the first bottle, I put a solution of potash \*, in about six parts of water. In the fecond, a folution of the fame; but so dilute, as that no part of the falt, which might be formed, should crystallize during the operation. About twenty parts of water was the proportion there employed. In the third bottle, I put common carbonate of potash. Through this apparatus, I fent a current of oxigenized muriatic acid, difengaged by fulphuric acid, from a mixture of muriate of foda and black oxide of manganese, in the well known manner. Crystals of hyperoxigenized muriate of potash were formed in the liquor of the first bottle; and, as long as they remained, I was certain, from previous experiment, that no fulphuric or muriatic acid could pass into the second bottle. The current was continued, till the liquor of that bottle contained an excess of acid. The car-The fecond bot-bonate of potash, in the third bottle, absorbed the superabundant vapours; and the pneumatic apparatus was ready to

gas was paffed in fucceffion. through ftrong folution of potash, weak solu. tion of potash, and folution of carbonate of potafh. The first bottle exhibited cryftals and detained all the fulph. or mur. acid. tle received only hyperox. mur. acid, and contained its diffolved combination with potash.

\* Whenever potasi, soda, barytes, an acid, an alkali, water, or the names of other substances are used without an epithet, they are meant to denote them in that state which is commonly called pure.

collect any gafes that might be evolved. By these means, I

obtained

obtained, in the second bottle, a solution of whatever substance The third bottle might refult from the action of potath upon hyperoxigenized abforbed the fumuriatic acid.

perabundant vapours.

I took a portion of this liquor, which I shall call entire The liquor of liquor \*, and diffilled it to dryness in a glass retort, taking the second bottle gave no gas by care to screen it from light. A tube from the receiver commu- diffillation, but nicated with the pneumatic tub. My object was to afcertain, left the dry cryftallized falt. whether the change observed by Mr. Berthollet, in the distribution of the elements of oxigenized muriatic acid, to form, with potash, a simple and a hyperoxigenized muriate, really took place among those elements themselves, independently of any absorption of oxigen from the atmosphere, or extrication of it from the falt. Nothing but some water, and a few inches of the dilated air of the vessels, passed into the receiver and the pneumatic apparatus; and I found, in the retort, a faline mass +, perfectly dry and crystallized. Hence it is evident, that the same quantity of oxigen as that formerly contained in the oxigenized muriatic acid, which had been united to the alkali, to form the total mass of salt, was now condensed, in that part which had become hyperoxigenized muriate.

To ascertain this quality, I dissolved 100 grains of the entire 100 grains of falt in water, and precipitated by nitrate of filver. I thus ob-this falt precipitated as much tained a quantity of muriate of filver, which, by proportions filver as 84 previously determined, I knew to correspond to 84 of muriate grains of muriof potash: therefore, 16 were hyperoxigenized muriate of would have done. potafii t. But, according to the proportions established above Therefore 16 in hyperoxigenized muriate of potash, 16 of this salt contain 6 Parts were hyof oxigen, with 3,20 of acid, the remainder being alkali and muriate. water; and, by preliminary experiments, I found that 84 of muriate of potath contained 27,88 of muriatic acid. Therefore 27,88+3,20=31,08 of muriatic acid, with 6 of oxigen, or, to reduce it to the quintal,

I am well aware that, upon philosophical principles, this appellation is objectionable; but, for the fake of brevity, I have used it as a temporary name for a substance which has only a relative existence among chemical bodies.

+ This falt I shall call entire falt.

Line

I must observe here, that hyperoxigenized muriate of potass does not, like simple muriate, decompose the falts of silver. This shall be further animadverted upon, and proved, in its proper place.

Muriatic acid + - 84
Oxigen - - 16

100, are the proportions

which combine to form oxigenized muriatic acid.

Confirmation by diffilling off the oxigen.

To corroborate this evidence, I diffilled 100 grains of the intire falt mentioned above; and obtained nearly 16,5 cubic inches of oxigen gas; which as accurately corresponds with the trials by nitrate of filver, as can be expected in experiments of this nature.

Comparison with the experiments of Berthollet;

Mr. Berthollet, in his memoir upon oxigenized muriatic acid, gives, if I understand him rightly, the following statement of the proportions, and of the means by which he obtained his refults. He exposed to the light of the fun, 50 cubic inches of water, faturated with oxigenized muriatic acid; and collected in the pneumatic tub, 15 cubic inches of oxigen gas. I here neglect fractions; because our results appear, at first fight, to differ so widely as not to require great accuracy in giving their comparative statement. He then precipitated, by nitrate of filver, the 50 cubic inches of liquor, which had become fimple muriatic acid, and obtained 383 grains of muriate of filver. But, by experiments, I found that 383 grains of muriate of filver contain 65 of muriatic acid. Therefore, 65 of muriatic acid combine with 15 cubic inches \* (= 8 grains) of oxigen, to form 73 of oxigenized muriatic acid. But 73:8::100:11, or nearly. For this difference, however, it may be easy to account. Perhaps Mr. Berthollet's 50 cubic inches of oxigenized muriatic acid, contained originally a little simple muriatic acid; and he fays besides. that he suspects all the oxigen was not disengaged. This indeed is most probable; and I am happy that I can reconcile the proportions which I have found, to the opinion of fo skilful a chemist.

and Cruick thank. Mr. Cruickhank likewife, in his additional Observations upon Hydrocarbonates, has stated that 2,3 parts of oxigenized muriatic acid contain 1 of oxigen, or about 43,5 per cent. But this able chemist, to whom we are indebted for the discovery of the gaseous oxide of carbon, procured his oxigenized muriatic

<sup>\*</sup> Mr. Berthollet's proportions are in the old French weights and measures.

acid by a peculiar method, which I shall notice in speaking of the action of acids upon hyperoxigenized muriate of potash. The fubstance he obtained was, in fact, not oxigenized muriatic acid gas, but a mixture of that gas with hyperoxigenized muriatic acid. I have not the smallest doubt of the accuracy of his flatement; but, being the proportion of a mixture, it in no way contradicts either of those I have determined in this Paper.

Before I dismiss this part of the subject, I wish to anticipate There was no an objection, founded upon an observation of Mr. Berthollet, effervescence or loss of oxigen in which may be made to the above experiments. He fays, that the experiments. when the alkaline folution is very concentrate, an effervefcence takes place durin thewhole of the faturation, and for some days after; and this effervescence he attributes to the escape of oxigen. But I have already faid, that no oxigen gas was disengaged in any part of my process; and no effervescence took place in any of the bottles, except the third; fo that no superabundance of oxigen could have passed from one It is probable the into the other, nor could any diminution of the total quantity effervescence of Berthollet was have been produced. By repeating the experiments, some-owing to carbotimes with a folution of alkali, and fometimes with water alone, nic acid. in the first bottle, I obtained the liquor in the second bottle uniform in all cases. Indeed, as potash prepared in Mr. Berthollet's manner, was not in fuch general use at the time he performed his experiments as at prefent, I suspect that a great part of this effervescence was owing to a disengagement of carbonic acid from the alkali.

Having thus proved the difference between the states of these two acids, I shall now proceed to the combination of each with falifiable bafes.

# OXIGENIZED MURIATES.

As many properties of the entire liquor, before it had been Whether the lievaporated to dryness, had led me to imagine that the acid was quor before eva-poration did con-united with the alkali, and remained in combination with it, tain the two falts in the state of oxigenized muriatic acid, till the moment of or only one oxicrystallization, I think it necessary to state at length the ap- fore the instant pearances which induced me to draw that conclusion, and the of crystallization. experiments which afterwards convinced me that it was erroneous.

Sulphuric or acetic acid difengaged oxyg. m. acid from the muriatic acid.

A few drops of fulphuric acid, poured into fome of the intire liquor, caufed an effervescence, and a smell of oxigenized

liquor; but not from the falt.

Very strong acetic acid produced the same effect.

By other experiments, I had afcertained that acetic acid could not decompose any part of the intire salt; and hence I concluded, that in the intire liquor, before evaporation, some of the falt remained in the flate of oxigenized muriate, the acid of which was expelled by the fulphuric or acetic acid; and, that it was not till the moment of crystallization that the elements of the falt underwent a total resolution into muriate, and hyperoxigenized muriate of potash. However, a small quantity of any of the very foluble neutral falts, fuch as nitrate or muriate of ammonia, or even a little alcohol, produced the fame effects; and I was then convinced, that the effervescence was owing to fome uncombined oxigenized muriatic acid gas remaining in the liquor; and which was difengaged, in proportion as the water was taken from it, by the superior affinity of the falt, or the alcohol. I had used.

This however arises from some disengaged excess of oxyg. mur. acid; for neutral falts could extricate it.

Acetic and acedecompose hyperox. m. of potash:

but ox. m. acid decomposes acetite of potash.

In fuch difengagements the alkali becomes had at first been free.

By some previous experiments I had ascertained, as I have tous acids do not just mentioned, that acetic or acetous acids do not decompose hyperoxigenized muriate of potash. I fent a current of oxigenized muriatic acid through a folution of acetite of potafit; and, upon evaporation, I found that the acetous acid had been difengaged, and that muriate, with hyperoxigenized muriate, of potash had been formed. But, from some trials, which I shall presently relate, I was induced to believe, that oxigenized muriatic acid attracts the falifiable bases with a much weaker affinity than acetous acid. It is well known that the contact of oxigenized muriatic acid with an alkali, is fufficient to produce combined as if it a combination of that acid with the alkali; and, from the lastmentioned experiments it appears, that it is not abfolutely neceffary that the alkali should be in a free state. If it be combined with an acid weaker than hyperoxigenized muriatic acid, the original acid will be expelled; and muriate and hyperoxigenized muriate will be formed, as if the alkali had been free.

As a further proof, that the change in the distribution of oxigenized muriate of potalh takes place at the moment of contact of the acid and the alkali, and confequently long before the crystallization, I mention the following experiments.

[ pre-

I precipitated, by nitrate of filver, 400 grains of the entire The first liquor liquor, previously to its being evaporated; and obtained 71 was precipitated by nitrate of grains of muriate of filver. fliver :

I evaporated to dryness, 400 grains of the same liquor, redis- and afforded the folved the refiduum, and, by dropping in nitrate of filver, ob-folution of the tained 70 grains of muriate. The difference of one grain, in intire falt. these experiments, does not amount to 0,2 of a grain of muriate

of filver; and ought not to be regarded.

From these experiments it is past all doubt, that the original Hence the first intire liquor did not contain oxigenized muriate of potash. liquor did not contain oxigenized muriate of potash. For, if such a combination had existed in it, I should have of potash; but obtained a finaller portion of muriate of filver in the first than the two diffines in the fecond cafe, on account of the total feparation into muriate and hyperoxigenized muriate having not yet taken place.

We are not however to conclude, from these experiments, Yet it is possible that there are no fuch things as oxigenized muriates. Although to form oxigthey cannot be exhibited in a palpable state, it is easy to demonstrate that they do really exist. I shall prove, in the pro-for example, of per place, that hyperoxigenized muriate of ammonia is not ammonia. an incompatible combination; and must, for the present, asfume the datum, in order that I may demonstrate the necessary existence of oxigenized muriates. Therefore: If muriatic acid, or if hyperoxigenized muriatic acid, be brought in contact with ammonia, the refult will be muriate, or hyperoxigenized muriate, of ammonia. But, if the acid, difengaged by For the oxigenfulphuric acid, from a mixture of black oxide of manganese ized acid can deand muriate of foda, be fent through ammonia, both are decom- nia, and the hyposed. Hence it is evident, that the acid combines with the perox. m. acid alkalis, in the state of oxigenized muriatic acid; and that the this last does, in feparation into muriate and hyperoxigenized muriate, is pro-fact, form a falt duced by a subsequent action, among the elements of oxigen- with it in the experiment. ized muriate of potash.

Upon the whole, it appears to me fair to conclude,

1st, That the falts of this genus do really exist, previous to the formation of hyperoxigenized muriate of potash.

2d, That the affinity exercised by hyperoxigenized muriatic Consequently acid for ammonia, and (by very strong analogy) for the other there must have been an oxig. bases, is much greater than that of oxigenized muriatic acid. muriate first, and For, hyperoxigenized muriatic acid, as shall presently be shewn, this was immediately, by having a much more powerful action upon all combustible stronger affinity,

bodies, fucceeded by the

other falt, instead of decomposition ensuing.

bodies, whether fimple or compound, than oxigenized muriatic acid, it would be natural to suppose that the former acid would act more powerfully upon the inflammable element of ammonia. But oxigenized muriatic acid combines with the hidrogen of that alkali; which, however, is not decomposed by hyperoxigenized muriatic acid; yet the affinity of hyperoxigenized muriatic acid for ammonia, is the only cause that determines the union of the acid and the alkali, without decomposition. these affinities shall be more fully treated of, in speaking of hyperoxigenized muriate of ammonia.

# ALKALINE AND EARTHY HYPEROXIGENIZED MURIATES.

### Generic Characters.

Hyperoxigenized muriates are formed by passing a current

Alk. and earthy hyperox. muriates.

of oxigenized muriatic acid through the basis, disfolved or Generic charac- suspended in water, as in the formation of the last-mentioned genus. The first formation is owing to the separation of the elements of an oxigenized muriate, into hyperoxigenized muriate and simple muriate; from which latter, they may be feparated by chrystallization, or by another process, which I shall mention, in treating of the earthy hyperoxigenized muriates. By fimple trituration, they scintillate, with noise. They are decomposed by a low red heat; and give out a confiderable quantity of oxigen, as they become simple muriates. They cannot be brought down, by any means that I have tried, to that diminished state of oxigenizement, which would constitute oxigenized muriates. They inflame all combustible bodies with violence, as is well known. They are foluble in water; many of them, in alcohol; and fome are deliquescent. The acid is expelled, with particular phenomena, by fulphuric, nitric, and muriatic acids, without heat; and, a little below a boiling heat, by phosphoric, oxalic, tartareous, citric, and arsenic acids: but they are not acted upon by benzoic, acetic, acetous, boracic, pruffic, or carbonic acids. Those vegetable acids which are powerful enough to decompose them, give out, towards the end, a gas of a peculiar nature, which has not to much finell as oxigenized muriatic acid gas, but which affects the eyes in an extraordinary manner, and promotes an

uncommon

uncommon and rather painful fecretion of tears. I have not yet examined this gas, as there was invariably an inflammation of the mixture, with explosion and rupture of the vessels, almost as soon as it began to be evolved. When pure, the hyperoxigenized muriates do not precipitate any of the metallic falts, although I believe they decompose some. The order in which the bases seem to be attracted by the acid, is, potash, soda, barytes, strontia, lime, ammonia, magnesia, alumina, filica. The other earths I have not tried, and but few of the metallic oxides.

# First Species. Hyperoxigenized Muriate of Potash.

This falt is the best known of all the saline combinations of Hyperox. mur. this acid. It has been erroneously considered as simply oxigenized, for its acid is really hyperoxigenized. It is foluble in about fixteen parts of cold water, but in much less of warm; and is easily separated, by crystallization, from muriate of potash. Alcohol can dissolve a small portion of it. It feems ca-Probably capable pable of existing in more states than one; for, in passing a cur-various states. rent of oxigenized muriatic acid, very flowly, and in the dark, through a folution of potash, till saturated, I have obtained flexible and shining needle-like crystals. This leads me to suspect, either a hyperoxigenized muriate of potash with excess of acid, or that acid with a superaddition of oxigen. It would be superfluous to enter into a minute description of a substance so well known as hyperoxigenized muriate of potasi; but, it being the fubstance whence I have chiefly attempted to difengage the acid, I shall enter into a particular detail of the action of the more powerful acids upon this falt.

If concentrate fulphuric acid be poured upon hyperoxigen-Attempt to deized muriate of potasis, a violent decrepitation, sometimes but compose it by sulph. acid addrarely accompanied by a flash, takes place. A thick heavy ed to the falt. vapour, of a greenish yellow colour, which rifes with difficulty Peculiar vapour. to the top of the vessel, if it be deep, is disengaged. The finell is not altogether unlike nitrous gas; but it is peculiarly fetid, and may be compared to that which is emitted by brickkilns, mixed with that of nitrous gas. It differs much from oxigenized muriatic acid gas; the latter being pungent and penetrating, the other heavy and oppressive; and it does not produce, at least in so great a degree, the catarrhal symptoms caused by the other. At the bottom of this vapour is a bright

engaged is not pure.

orange-coloured liquor, which has the fame fmell as the va-The hyperox.

m. acid thus difpour. This is the acid contained in the falt; and I have proved it to be hyperoxigenized muriatic acid. But, although the falt from which the acid is difengaged be pure, the acid itself is never fo; because the very act of disengaging it effects its decomposition, and some of it is converted into oxigenized muriatic acid. The colour of litmus paper, on this account, is generally destroyed by the liquor. I fay on this account, because I have some reason to believe, from having observed this not to be uniformly the cafe, that hyperoxigenized muriatic acid reddens the vegetable blues. However, it must be considered, that the fulphuric acid used to difengage the hyperoxigenized muriatic acid is still present; and we can draw no certain conclusion, until we have obtained this acid free from all other fubftances. If to this mixture of hyperoxigenized violent and dangerous explosion, muriate of potash and sulphuric acid, heat be applied, an exceedingly violent explosion, with a white and vivid flash, takes place, before the liquor has attained the temperature of 125 of Fahrenheit. In order to obtain this acid, I attempted to diffil 500 grains, in a glass retort, in a water bath, with every precaution against such accidents as I could not but in some measure expect; when, almost as soon as I had kindled the fire, I faw, in the bottom of the retort, an extremely white, vivid, and rapid flash, which was immediately followed by a

loud report. The retort was reduced almost to powder; so

that fearcely any fragments of it could be found in the labora-

The windows, and feveral glass vessels, were broken. I happened to be holding the neck of the retort, at the moment of the explosion, yet received no injury, except a slight contufion in the hand. But Dr. Vandier, a French gentleman of confiderable chemical and medical talents, to whom I am indebted for much able affiftance in my laboratory, was wounded in feveral places; particularly, the tunica conjunctiva of the eye was so lacerated, that a piece of it hung down, and, by getting under the inferior eyelid, caufed the most painful irritation, and endangered his fight. One of the frontal arteries alfo was divided. I relate these circumstances thus fully, as

Narrative of an accident of this nature.

Heat produces

plied.

the most effectual means of putting upon their guard those who Dilute acid ap- would repeat the experiment. If the fulphuric acid be dilute, heat may be applied with more fafety; and the phenomena are different. The hyperoxigenized muriatic acid is difengaged

gaged from the basis; but, as the heat requisite to distil the acid is more than sufficient to decompose it, oxigenized muriatic acid comes over with it; and oxigen gas is collected in the pneumatic tub. If the distillation be continued, the same danger arises as in the former case, because the sulphuric acid becomes concentrated; and it would seem, that its action upon the salt is slight and partial at a low temperature, but violent and instantaneous when heated and concentrate. I could not, therefore hope, by these means, to obtain the acid disengaged and pure.

If the manner of bringing the fulphuric acid and the falt into The falt added contact be reverfed, and the falt be dropped into the acid, the to the acid, yellow vapours and the orange-coloured liquor are produced, but generally without decrepitation. If they be allowed to remain some days in contact, the vapours continue, and oxigen gas is constantly disengaged, even in the common light of the day, and at the temperature of the atmosphere. By cooling the first receiver with ice, I thought that I had once obtained this acid crystallized in the form of four-sided pyramids, of an orange colour. But, though I really believe this to have been the case, I do not positively affirm it.

Nitric acid produces nearly the same phenomena; but the Nitric acid apfmell and other properties are rather less distinct and marked, plied.

than with fulphuric acid.

Muriatic acid decomposes this salt, and unites to its basis; Muriatic acid. but neither the yellow vapours, nor the orange-coloured liquor, are produced. The circumstances which attend the contact of the acid and the salt, are as follows. If no more muriatic acid be present than is merely necessary to decompose the salt, I do not doubt that hyperoxigenized muriatic acid will be driven off, as little decomposed as with the other acids, supposing the action to be instantaneous; but, during the contact of these two bodies, the acid expelled must meet muriatic acid not yet combined, and, uniting with it, always forms a portion of oxigenized muriatic acid. The quantity of the last acid must vary, according to the quantity of muriatic acid employed, and not combined with the alkali. It was by this method that Mr. Cruickshank obtained the muriatic acid gas, which he stated to contain 43,5 per cent, of oxigen.

Phosphoric and arfenic acids do not act upon this falt, till Phosph. and aceheated with it; and then much oxigen gas is evolved. These, gage oxigen by therefore, heat.

therefore, afford no better method of difengaging hyperoxigenized muriatic acid without decomposition.

Oxalic, tartare-1 ous, and citric acids.

Oxalic, tartareous, and citric acids, act as I before mentioned; and the hyperoxigenized muriatic acid holds its place, in the order of affinities for potash, immediately before the benzoic.

Amuling combuffions by this falt and strong acids.

I shall not stop to detail a number of amusing phenomena that may be produced, by projecting into the stronger acids, mixtures of combustible bodies, whether metallic or not, and hyperoxigenized muriate of potash. The cause of them is well understood, and the theory points them out: they are, therefore, no longer objects of philosophical admiration. But I must mention one experiment, which, had it fucceeded, I should have thought important. I projected various mixtures of very minutely pulverized diamond and this falt, into the different acids; but found the diamond undiminished, by every attempt to combine it with oxigen in the humid way \*.

The diamond does not burn in this humid way.

The h. ox. muriates contain much caloric.

Another, but imponderable, part of this falt, as indeed of all hyperoxigenized muriates, feems to be an extraordinary quantity of caloric. For, during their formation, scarcely any heat is difengaged, as by other acids; and, very little heat applied to the falts, gives the gaseous form to their oxigen.

Erroneous opi- .

An opinion has prevailed among fome ingenious chemists, nion, that ni-trous gas may be that, from a mixture of this falt with fulphuric acid, nitrous gas formed from this is disengaged, and sulphate of lime formed in the retort. But this is a mistake, arising, on the one hand, from the smell and vapour of the hyperoxigenized muriatic acid, and, on the other, from fulphate of lead, which the common fulphuric acid of this country frequently contains in folution, and which is precipitated from it by water. Before we affert a fact, we should be well affured of the pureness of our chemical agents. This

> \* I must confess, that the vivid stasses of light, emitted from the mixture of this falt and combustible bodies thrown into an acid, appear to me, in some measure, to prove the modification proposed by Leonhardi, Richter, Gren, &c. to that part of the Lavoisierian theory which regards the emission of light during combustion. Another testimony in favour of their modification, may be drawn from the vegetable kingdom. All plants growing in places deprived of light, are merely mucilaginous. But the mucilage of these plants burns without the emission of light. Light, therefore, appears not to be disengaged from oxigen; else, why not by this mucilage as well as by other combustible bodies?

supposed conversion of muriatic or hyperoxigenized muriatic acid into nitrous gas, will not pass for a decomposition, or a transmutation, of that refractory radical; and the idea of the change of potash into lime, is as erroneous as some other late affertions respecting the decomposition of the alkalis.

The proportions of this falt are, as I before stated,

Hypero	viceni	zed m	uriatic	acid	_	58.3
Potash	go	zea n	-	-	-	39,2
Water	-	-	•		-	2,5
4	250	_ =				100,0.

Component parts of hyp. ox. mur. of potash.

Second Species. Hyperoxigenized Muriate of Soda.

This falt is prepared in the fame manner, and with the fame Hyper ox. mur. phenomena, as the former. It is extremely difficult to obtain of foda. Very difficult to it pure, as it has nearly the same degree of solubility in water be separated, beas muriate of foda. It is foluble in three parts of cold, and cause of the little less of warm water; and is slightly deliquescent. It is soluble limiting. in alcohol; but this property alone is not fufficient to enable us to obtain it free from the muriate of foda, formed along with it in the entire liquor; as the latter falt, contrary to the affertions of all authors, is foluble in alcohol, and feems to be much more fo, when accompanied by the hyperoxigenized muriate. "It Obtained by was by taking a large quantity of the entire falt, formed by crystallizing in fending a current of oxigenized muriatic acid gas through a folution of carbonate of foda, and repeatedly crystallizing in alcohol, that, with great difficulty, I obtained a little pure hyperoxigenized muriate of foda. It crystallizes in cubes, or in rhomboids little different from cubes. It produces a fensation of cold in the mouth; and its taste is easily distinguished from muriate of foda. It is decomposed by heat, by combustible bodies, and by acids, in the same manner as the former species; and the acid holds its place for foda, as for potash, immediately before the benzoic. The basis is separated by potash only. This falt is composed of,

Hypero	xigeniz	ed muria	tic acid	' II . I	66,2	Component
Soda	CT OF		- 11	-	29,6	parts.
Water		1-1-	H		4,2	
1,000		- 12				
73751		. 13			100,0.	

(The Remainder in our next.)

### VIII.

Description of the Indian Hand Mill. From a Correspondent.

General account THE Indian hand mill, like most other Indian inventions, is characterifed by confiderable fimplicity and effect. cheapness and its general utility are such, that the meanest hut in India is never without one. The whole of the grain used by the natives of India is ground in these mills, and chiefly by the women, who appear to execute the task with astonishing eafe. A woman will continue grinding with this mill for feveral hours, and in this time she will reduce a very confiderable quantity of grain into flour.

Description.

The mill is composed of two circular stones, an upper and an under one, each about two feet in diameter, and  $2\frac{1}{3}$  inches in thickness. The quality of the stone nearly resembles that of our common mill stones.

Figure and fitftone.

The upper face of the under stone is dressed perfectly flat. ting of the under A small hole, of nearly an inch in diameter, is sunk in the center of this stone: this hole is fitted with a plug of hard wood, firmly driven into it, which contains a finall feel gudgeon or pivot. The furface of this wooden plug is flightly elevated above the face of the stone, and the gudgeon projects about half an inch beyond it. This gudgeon fits into a hole in the middle of a cross bar, or flat piece of iron, eight or nine inches in length, about \$\frac{1}{4}\$ of an inch in breadth, and \$\frac{1}{4}\$ of an inch in thickness: the ends of this cross bar are let into notches in the upper stone, and it thus forms a center on which the upper stone may revolve.

The upper stone.

The lower face of the upper stone is dressed perfectly flat, to correspond with the upper face of the under stone. This from is 21 inches in thickness at the edges, but it has a proiection of nine inches diameter in the middle, at which place it is five inches in thickness. The stone is pierced in the middle by a hole of fix inches in diameter, which ferves as a hopper to contain the grain. In the upper face of this stone a hole is funk near the circumference, in which hole a wooden handle, about fix inches in length, is firmly driven. means of this handle the stone may be made to revolve on its gudgeon with confiderable facility.

The

The projection of the wooden plug in the under stone in How the fine-which the gudgeon is inserted is so regulated as to keep the ness of the flour stones at the proper distance for forming the finest flour: if a coarser sort is wanted, it is readily obtained by placing one or more slips of paper or card in the notches of the upper stone, between the cross bar and the stone.

The under stone is usually placed on the earthen floor of the Local situation hut, which is previously beaten and swept, or otherwise it is or positional placed on a cloth spread on the floor. As the grain is reduced to flour it escapes, together with the bran, at the circumference of the stones, and is collected on the floor or on the cloth.

I would fuggest as an improvement of this simple machine, Improvements the forming the cross bar with three claws, which would give suggested. three points of bearing instead of two, and consequently render the upper stone more steady. I would also suggest, that the block of wood which contains the gudgeon, should pass completely through the under stone; thus it might easily be adapted to the requisite height by a few blows of the mallet. Should it be thought necessary, the stones might be enclosed in a drum, and sitted on a stand.

Explanation of the sketch, Plate X.

Reference to the

Fig. 1. Reprefents the lower face of the upper stone.

Fig. 2. Represents the upper face of the same.

Fig. 3. Is an elevation of the mill.

Fig. 4. Is a fection of the mill.

a, Is the crofs bar, which is represented with three claws.

b, Is the handle by which the upper stone is turned.

c, Is the steel gudgeon on which the upper stone revolves by means of the cross bar.

d, Is the fmall block of hard wood in which the gudgeon is fixed.

\* It is likely that this steadiness would do harm. The totter in all probability answers the same purpose, of pounding, as the vertical dancing motion of the upper mill stone upon the support of its pivot, of which Desaguliers has given a very rational account. Lectures II. 428.

#### IX.

An Effect hitherto unpublished, by which the two Electricities are In a Letter from Mr. JOHN CUTHBERTSON. distinguished.

To Mr. NICHOLSON.

DEAR SIR,

Letter.

N reading your note respecting the instruments by which the two kinds of electricity are diffinguished, or its direction ascertained, it brought to my mind an experiment which I made with that view in the year 1777 in Holland; I believe it is not known here, so perhaps you may think it worthy of a place in your valuable Journal.

I am.

Dear SIR,

your's respectfully,

I. CUTHBERTSON.

Poland Street, Soho, Oct. 19, 1802.

Experiment. A candle between two balls in the electricity has its flame much agitated, and heats the negative ball only.

INSULATE two wires furnished at each end with a metal ball of about \( \frac{3}{4} \) of an inch diameter, connect one with the posiopposite states of tive conductor, and the other with the negative conductor of an electrical machine; fet them so that their balls may be at about four inches distance, place between them a common fized lighted candle, with the center of its flame nearly upon a level with the centers, and at an equal diffance from each; if the machine be put in motion the flame will waver very much, and feems rather more to incline to the negative ball than to the positive one, but is very equivocal: continue turning (if the machine be a plate of two feet diameter) about 50 revolutions, then the negative ball will begin to grow warm, and the positive ball remain cold; if the revolutions be multiplied to 200, the negative ball will be too hot to be touched, and the positive remain as cold as at the beginning.

# X.

Palpable Mathematics; or Methods of writing and calculating for the Use of the Blind. By J. B. BERARD.\*

CITIZEN Haüy having compared the methods of writing Citizen Hauy's and calculating of several celebrated blind men, has digested instructions to into a body of doctrine the best productions of experience in writing, sec. this art. He has composed a system of instruction, and has described the instruments proper to be used in this operation: and he truly deserves well of humanity, who increases the means of communication between society, and a class of men who seem to have been excluded from it, and to have been condemned to lead an unhappy life, useless to others, and without enjoyment to themselves.

Having been myself deprived of fight fince the age of twenty-The author is three, meditation and experience have given me the unhappy blindaright of treating upon the art concerning which he has written.

I here give some observations which may not be entirely use-

less to fociety.

# ARITHMETIC.

The method by which the celebrated Saunderson calculated Professor Saunis well known; however ingenious it may be, it does not derson's method of computing seem to me to be the best; I observe two principal defects in it.

He used small square pieces with nine holes in each: the by squares to dethe position of a pin in one of these holes denoted one of the note the digits, nine sigures. It is therefore necessary to have as many of these squares as the operation requires of sigures or characters: and we must employ an extended space for a very simple calculation.

In the fecond place, this method of calculating being en-required a great tirely different from that of persons who can see, they cannot space, and is not understand the particulars of it; this is a considerable inconvenience, because it destroys all connection between the blind and those who operate by sight.

The method which appears to me the best, consists in em-It is better to ploying characters representing figures, and arranging them compose as printers do, nearly as they are composed in printing.

Melanges Physico Mathematiques.

#### ALGEBRA.

and in algebra likewife.

The best method of writing algebra appears to me, like that of arithmetic, to confift in composing in the manner of printers, with this difference, that the characters are a little larger, and must be read from left to right. This process has the important advantage of enabling readers by fight to perufe and examine the calculations.

### GEOGRAPHY:

There is but one method of using geographical maps and Easy method of forming tangible the globes: it is simple and easy. A small string or thread is maps; passed on all the windings or outlines of the map. The rivers and mountains are diftinguished by different threads. Grains or prominences of different fizes and shapes are pasted on the and multiplying towns and other remarkable places, &c. With one of these copies. anaps others may be procured by the prefs, in which the out-

lines will be raifed into relief.

## MUSIC AND BOOKS

Music books.

Citizen Hauy has composed music books for the use of the establishment of the Quinze Vingts at Paris, the characters of which project, or are emboffed: this method is without doubt useful, but it is scarcely applicable but to great foundations.

#### WRITING.

Writing on pathe protuberance.

Citizen Hauy's method confifts in using an iron pen, the per with a blunt point of which is not split; by writing without ink and prefitile is legible by fing on a strong paper, the blind man produces a character in relief, which he can immediately read by passing his singers over the projecting characters on the opposite side of the paper, in the contrary direction. This relief is sufficient, provided a foft furface be placed under the paper, fuch as leather, blotting paper, &c.

Frame for writing straight.

In order to write in a straight line, a frame is put on the paper, having a number of brass wires across, parallel to the direction of the writing, at the distance of about three quarters of an inch.

His method of writing in relief has this advantage, that the blind can read what they have written, after a little practice; but

but it has the inconvenience of preventing persons that can see from reading it, unless they have also practifed the same method \*.

A black writing may likewise be obtained with this iron pen, How to make by placing on the writing paper another sheet done over with the writing wift-hogs-lard and ivory black, covered again by another, through which the letters are traced. The inconvenience of this method is, that the blind cannot read what they have written.

Instead of the frame with parallel threads, I employ a board

which has feveral conveniences.

This board is about 10 inches in breadth, and 16 high; on A more convethe left fide, and near half an inch from the edge, there is a nient frame for small dove-tailed groove from top to bottom. A rule three quarters of an inch wide, and  $4\frac{1}{2}$  long, slides in this groove by a piece so formed as to sit the groove. Another rule sive eighths of an inch wide, is fixed perpendicularly to the first rule in the form of a T, and serves very conveniently to direct the little singer of the hand that writes. Lastly, the left side of the board is divided into notches almost half an inch distant from each other: this being done, the following is the method of writing:

The paper is placed under the long rule, and is fastened by Method of writtwo screws easy to be imagined, which press it against the ing with ink. board: the writing is performed by keeping the hand lightly over the rule, the little finger being behind. The rule is shifted from line to line very readily by the left hand, the thumb being applied to feel the notches. At the finish of every line ink is dipped or taken (in a filver pen formed in the usual method) out of a broad slat vessel constantly containing ink to the depth of between one eighth and one quarter of an inch.

By this process, the lines may be written as near each other as may be required. The rule directs the hand better than a wire, which is too flexible, and not broad enough. When, after discontinuing writing the writer is desirous of proceeding, the rule indicates the point to begin from, &c.

\* If the paper be rubbed with pumice, or have some earthy powder in the size, and a blunt point of hard pewter be used, the letters will be neat and visible...-N.

### INK IN RELIEF.

Defideratum; a It would be a very useful invention, if an ink could be visible and tangible ink.

formed which in coming from the pen should leave a mark that would project. This ink would have the double advantage of enabling the blind and those who see to read at the same time writing, the figures of arithmetic, the characters of algebra, and the notes of music. I have not yet thought of any satisfactory expedient for this purpose, and shall not therefore enter into any further detail.

### GEOMETRY.

Saunderson's method of forming geometrical figures, placed pins in holes made in a board, and passed threads from one pin sing geometrical figures with pins to another. My method appears to be preserable in every reand threads. A better method; by wires on a flat cushion and thickness three quarters of an inch. I have covered the two faces with woollen cloth rather open and fost, and have filled the intermediate space with other loose and soft woollen cloth, so as to form a flat cushion three quarters of an inch thick.

I have very small iron wires, bended at right angles at the two ends, which are pointed and polished. These iron wires are easily fixed in the cushion, and form lines in relief, which nothing can derange. I have an assortment of all lengths, as well as circles, semi-circles, parabolas, ellipses, &c. Lastly, I have letters in relief, with a point in order to fix them on the cushion.

This method is very ready and comprehensive. By means of this little apparatus, there is no figure in geometry, however complicated, but may very fpeedily be represented in relief.

We might likewife, and with advantage, use this cushion for making figures with pins and threads; but this method, though better than that of Saunderson, is not so good as the iron wires here described. Experience is decisive in this respect.

and is applicable to mufic, &c.

It is easy to perceive that the above method is likewise the best for the notes of music; but the limits I have adopted will not allow me to enter into further details.

### XI.

Observations on the real Nature of the Precipitates formed by the Prushates in Acid Solutions of Barites, and the Affinities of the Prufic Acid \*. By CITIZEN GUYTON,

BERGMAN announced, that the acid folutions of barites Precipitation of are precipitated by the faturated Prussian alkali. This fact, fian alkali conconfirmed by the observations of a great number of chemists, sidered as an inappeared to many as a new indication of its metallic nature, dication of its metallic nature, metallic nature. already suspected on account of its great weight. Lavoisier in particular thought it very probable, that the difficulty of reducing it to the metallic state proceeded from its having more affinity with oxigen than carbon +.

The little fuccess of the attempts to effect this reduction This notion having in some measure caused the notion of this composition abandoned, to be abandoned, the attention was again fixed by the phenomenon of the precipitation of an earth by a re-agent, which according to the generally received opinion, afforded a characteristic indication of the presence of a metal. It was then suspected to be owing to some accidental cause. M. Meyer and the precipiof Stetin, announced in 1786, in Crell's Annals, that the tation by pure pruffiate denied. profliate of potath, if very pure and carefully prepared, did not precipitate barites from its folutions. Kirwan and Klaproth adopted this opinion from their own experiments. Most chemists however continued to believe that the precipitation

Some time ago being defirous to try a pruffiate of lime re- Carbonate of cently prepared with lime of marble, and very clear, I poured potash precipitates prushate of into it a folution of carbonate of potash. The liquor instantly lime. became milky and opake. As it is certain that no metallic substance was present, I could only attribute this decomposi-

took place, whatever might be the means of purification of the proof liquor. Pelletier, amongst others, supposed it one of the most essential distinctive characters of barites and stron-

- \* Read at the fitting of the class of Philosophical and Mathematical Sciences of the Institute, the 2d Thermidor in the year 70, and inferted in the Annales de Chimie, XLIII. 185.
  - + Traité Elementaire, &c. Vol. I. p. 174.
  - # Mem. de Chimie, Vol. II. p. 454.

tian t.

Vol. III .- NOVEMBER, 1802.

Q

tion

by double affinity.

tion to a change of base of the two salts by double assinity. then purposed to examine whether this might not be absolutely the same phenomenon which had given rife to the suspicion of the metallic nature of barites.

Mr. Henry's experiments from the Philof. Journal.

I was anticipated in these researches by Mr. W. Henry. He has fucceeded in placing this point of theory in the clearest light: I shall repeat in a few words the experiments that led him to it, and which he published in Mr. Nicholfon's Journal \*; after which I shall make some reflections upon one of the confequences he deduced from them, which appears to me to require amendment.

Pruffiate of barites formed,

Mr. Henry's first object was the direct composition of prusfiate of barites, in order to obtain a perfectly pure pruffiate of potash: the following is his process, which he affirms that he carried into effect, and was fuggefted to him by observing the decomposition of the sulphate and carbonate of potash by the pruffiate of barites.

. He calcines the carbonate of barites; he dissolves this earth

in boiling water, and adds Prussian blue till it is no longer dif-

coloured. The filtered liquor fometimes becomes turbid by

by calcining the carbonate of barites; diffolying the earth in Pruffian blue ger discoloured by the earth feizing the p. acid. The crystallized prusifiate of ba-

water, and add cooling, and deposes a little oxide of iron. He filters it again, till it is no lon- and at the end of feveral hours small yellowish crystals are formed, which are the pruffiate of barites. He obtains an. additional quantity by evaporation. This pruffiate of barites in powder is thrown into a hot folution of carbonate of potath; until it no longer restores the, colour of blue paper. The author observes, that it is better to rites is decompofed by carboemploy more pruffiate than is necessary for the decomposition of the carbonate. He digefts the mixture for about an hour; then filters the liquor, evaporates it flowly, and obtains by

nate of potash, and thus gives fine crystals of pruffiate of potafh.

this method very fine crystals of prussiate of potash. These crystals nevertheless sometimes contain as much as by acetous acid. 0.24 of oxide of iron; the greatest part of this is separated by digesting the liquor before filtrated, with a little acetic acid, which has the advantage of forming with the potath an una crystallizable salt, and consequently cannot mix with the crystals of the prussiate. Read or the Little - I Tile clair of

Last purification,

The precipitated carbonate of barites may again be used in the fame operation. I mail at a land of mi berreini ben or † Tanifé I'll mentaire, du. V l. I. p.

Let us now observe the use Mr. Henry made of the prussiate of potash thus prepared to resolve the controverted question.

He poured some of it into a solution of muriate of barites. The pure prus-No precipitation followed; the mixture was not in the least did not precipidisturbed, and he was inclined to think with Meyer and Klap- tate the muriate roth that a pure pruffiate does not precipitate barites.

But having observed the mixture half an hour afterwards, but it did in half he perceived small crystals forming on the sides of the vessel; an hour. and at the end of fome hours they were confiderably aug-

mented.

These crystals when examined presented the following chatacters:

1. They are very sparingly soluble in cold water; one Character of the ounce is required to dissolve a quarter of a grain \*. precipitated cry-

2. Warm water dissolves them rather quicker, though in a

very small proportion.

- 3. These solutions afford with sulphuric acid and sulphate of They were unpotash a precipitate of sulphate of barites; and Prussian blue doubtedly pruswith fulphate of iron.
- 4. These crystals are completely dissolved in muriatic acid diluted with water, and this folution afforded unequivocal figns of the presence of prussiate of barites.
- 5. These crystals heated to redness in a filver crucible became black, and loft their form; muriatic acid poured on the coaly refidue produced an effervescence, and muriate of ba-Ates was formed.

Mr. Henry draws feveral consequences from these facts. Inductions from The first, that he was deceived, as well as those who an-these facts. nounced, that pure barites does not decompose prussiate of potafh.

The second, that barites ought to be placed before potash in the column of affinities of pruffic acid.

The third, that the precipitation effected by pruffiate of potash' in a solution of muriate of barites, is the product of a double affinity.

The fourth, that barites differs in this respect from the other earths, and approaches the metals.

Of these four propositions, the three first seem to me to be open to no objection; for though Mr. Henry does not relate

\* As the English ounce is here spoken of, which contains 480 grains, cold water only diffolves about 0,0005 of its weight .--- G.

any direct experiment on this fubject, it is known, and I have myfelf observed it, that the aqueous solution of barites poured into a folution of alcaline pruffiate, decomposes it by fimple affinity.

Objection to the inference that barites approaches to the metals :

It is not the same with the last conclusion of Mr. Henry; for though these acid folutions of barites are decomposed by the alcaline pruffiates, precifely like the metallic folutions by the way of double affinity, or the concurrence of divellent forces which determine the exchange of bases, it in no respect follows that barites is on that account removed from the because the pre- earths, or approached to the metals. If that were the case, the fame must be afferted of all substances, which present the fame phenomenon, and it would no doubt be a matter of furprize to find this reasoning lead us to assimilate to the metals lime, strontian, magnesia, potash, soda, and even ammonia, fince the Pruffic acid can take all these bases from their solvents when the fum of the divellent affinities is in its favour.

cipitation is by double affinity, and happens with other earths and alkalies,

nàmely potash;

I have proved this in the case of potash by the experiment already related, where it is feen that it takes the Pruffic acid from the lime, and yields to it the carbonic acid.

foda;

This effect is still speedier with carbonate of foda, and prusfiate of lime.

Arontian;

The folution of nitrate of strontian, is likewise decomposed. by the pruffiate of lime.

lime;

If a few drops of pruffiate of lime be poured into a folution of sulphate of ammonia in five or fix minutes, a whitish frothy film of the thickness of about a quarter of an inch thick swims at the furface, and is at last deposed in small slakes.

magnefia;

Lastly, sulphate of magnesia mixed with prussiate of lime becomes clouded, and a precipitate is formed.

and because the is not the only distinctive cha-

Let us therefore return to the true cause of these phenoorder of affinity mena, the action of the double affinities in which we still trace the fimple elective attractions, the order of which is toracter of bodies. tally independent of that feries, or arrangement in which we place such substances as have several other properties in common. Therefore the station which barites holds before potash in the columns of the attractions of prussic acid, ought no more to be applied in its classification, than the position of lime before the fixed alcalis in the columns of the oxalic, tartarous, and boracic acids, &c. With much more foundation it has been proposed to rank barites with the alcalis, whose chemical action it in fact performs in many circumstances, but from which it however differs, as I have elsewhere observed \*, in some characters, which may be considered as limits, and particularly by its infolubility in alcohol.

#### XII.

A Review of some Experiments, which have been supposed to disprove the Materiality of Heat. By WILLIAM HENRY. From the Memoirs of the Literary and Philosophical Society of Manchester, Vol. V. Part II.

HE following remarks, on the subject of heat, were written Preface. foon after the publication of Count Rumford's Inquiry concerning the Source of the Heat evolved by Friction; and of the interesting essays of Mr. Davy, which appeared in Dr. Beddoes's West Country Contributions. They were transmitted to Dr. Beddoes, for publication, about the close of the year 1799; but circumstances, with which I am unacquainted, have, I believe, induced the Doctor to decline the continuation of his periodical work. These circumstances I deem it necesfary to flate; because, had the essay been written nearer the period of its publication, it would probably have assumed a very different form. At present, I have not leisure to review the subject, or to attempt any material alteration; and still less, to examine whether I have been anticipated by any of the authors whose essays have been published during the two last years.

A Review of some Experiments, which have been supposed to disprove the Materiality of Heat.

It has long been a question among philosophers, whether question whethe fenfation of heat, and the class of phenomena arising from ther heat be matthe same cause, be produced by a peculiar kind of matter, or by motion of the particles of bodies in general. The former of these opinions, though far from being universally admitted, is now most generally received; and the peculiar body, to which the phenomena of heat are referred, has been demonstrated by M. Lavoisier, caloric. Against the doctrine of the French

<sup>·</sup> Annales de Chimie, Vol. XXXI. p. 267.

school some forcible arguments have lately been advanced by Count Rumford and by Mr. Davy, both of whom have adopted that theory respecting heat, which assigns, as its cause, a motion among the particles of bodies.

The method of reasoning, employed by Mr. Davy, in

Mr. Davy's opfurdum.

matter of heat is proving the immateriality of the cause of heat, is the reductio by reductio ad ab- ad abfurdum, i. e. the oppugned theory is assumed as true, together with its applications; and facts are adduced, directly contradictory of the assumed principles. I shall take the liberty of offering a statement of the argument, rather different from that of Mr. Davy; though I trust without misrepresentation, or any material omission.

Another statement: Affuming the existence of caloric, &c.

Let heat be confidered as matter; and let it be granted, that the temperature of bodies depends on the presence of uncombined caloric. Now, if the temperature of a body be increafed, the free caloric, occasioning that elevation, must proceed from one of two fources; either, 1stly, It may be communicated by furrounding fubftances; or, 2dly, It may proceed from an internal fource, i. e. from a difengagement of what before existed in the body, latent or combined. But the temperature of bodies is uniformly increased by friction and percuffion, and, necessarily, in one of the foregoing modes.

in vacuo,

Heat by friction I. Mr. Davy found, by experiment, that a thin metallic plate was heated by friction in the exhausted receiver of an airpump, even when the apparatus was infulated from bodies capable of supplying caloric, by being placed on ice. This experiment he confiders as demonstrating, that the evolved caloric could not be communicated by furrounding bodies.

does not exclude caloric,

To the inference deduced from this experiment it may be objected, that the mode of infulation was by no means perfect. Admitting the vacuum produced by the air-pump to have been complete, still the supply of caloric could not thus be intirely cut off; fince it has been shewn by Count Rumford, that caloric passes even through a torricellian vacuum. If, therefore, friction produce in bodies some change, which enables them to attract caloric from furrounding fubftances, this attraction may be equally efficient in an exhausted receiver, as in one containing an atmosphere of mean density. It would be an interesting subject of experiment; to determine the influence of atmospheres of various densities, as conductors of culoric; for, fince effects are proportionate to their causes, and it is afcertained

becaufe a vacuum conducts.

tained that common air conducts caloric better than it is conveyed through a vacuum, as 1000 is to 702, it may be expected that the ratio will hold in all intermediate degrees;

In Count Rumford's mafterly experiment, the metal, full-In Count Rummitted to friction, was encompassed by water; and air was ford's experiments (Philof. carefully excluded from the surfaces in motion. Yet the water Jour. 4to, II. became hot, and was kept boiling a confiderable time. In this might be fupcase, the only obvious source of caloric, from without, was posed to be abthrough the borer employed in producing the friction; if it be forbed and emittrue, as the Count has observed, that the water could not, at time. the fame inflant; be in the act of giving out and receiving heat. The same objection to the communication of heat, from an external fource, exists also in thus explaining Mr. Davy's experiment: but I cannot admit that the argument is demonstrative. in proving the evolved caloric not to be derived from external substances; for no absurdity is implied in supposing, that a body may be receiving caloric in one state, and giving it out in another. We have an example of the fimultaneous admiffion and extrication of a fubtle fluid, the materiality of which is admitted by Mr. Davy, in an excited electric, which, at the Instance in elecvery fame inflant, receives the electric fluid from without, and tricity. transfers it to the neighbouring conductors. In an ignited body, allo, the two processes of absorption and irradiation of light, are, perhaps, taking place at the fame moment. Mi Another cause of the increase of temperature in bodies, In bodies which

is the liberation of their combined caloric; and, if this be a mit heat by fource of temperature, the absolute quantity of caloric in a body not have their must be diminished by friction. That no such diminution really capacity diminished; takes place, we have the evidence of two experiments—the one of Mr. Davy, the other of Count Rumford. Mr. Davy, by rubbing together two pieces of ice, converted them into water. Now water, ex hypothesi, contains more caloric than the ice from which it was formed; and, on the fame hypothefis, the absolute quantity of caloric in ice is diminished by friction and liquefaction, which is abfurd. Count Rumford alfoafcertained, that the specific heat of iron was not diminished when converted by a borer into turnings, and confequently when it had been the fource of much temperature. In explaint may be sup-

nation of these facts, we may be allowed to assume the communication of these facts, we may be allowed to assume the communication of these facts, we may be allowed to assume the communication of these facts, we may be allowed to assume the communication of these facts, we may be allowed to assume the communication of these facts, we may be allowed to assume the communication of these facts, we may be allowed to assume the communication of these facts. nication of caloric from furrounding bodies; till this communi-out;

cation has been demonstrated to be impossible. But even were

the

and the experiments for decidquestionable.

the impossibility established, it would yet remain to be proved, that the evolved caloric does not proceed from an internal fource; and this can only be done by an accurate comparison of the quantity of caloric in bodies, before and after friction. Now, in instituting this comparison, it is implied, that we ments for decid-ing capacities are possess means of determining the absolute quantity of caloric in bodies, and that we can compare quantities of caloric with as much certainty as we can obtain from an appreciation by weight or by measure. Such perfection however does not, I apprehend, belong to the prefent state of our knowledge respecting heat; for I have always been distrustful of that part of the doctrine, which assigns the ratio of heat latent in bodies. The grounds of this diftrust I shall state pretty fully; for, if it can be proved that we have no accurate conceptions of quantity, as appertaining to heat, all arguments against its materiality, derived from supposed determinations of its quantity, must be inconclusive.

It is afferted that our only clear conceptions of quantity are from magnitude er weight;

the gravity of bodies. In the inftance of caloric, both thefe modes of mensuration fail us. We cannot estimate the bulk of a fubstance which eludes our grasp and our vision; nor have we yet fucceeded in comparing its gravity with that of the groffer kinds of matter, which it surpasses in tenuity beyond all neither of which comparison. Our notions of the quantity of caloric are derived, ean be exhibited not from fuch simple judgments, but from complicated processes of reasoning, in the steps of which, errors, fatal to the whole, may perhaps fometime appear.

The only clear conceptions which the mind has of quantity,

are derived either from a comparison of the magnitude, or of

Caloric is peculiar in its incre-Cons.

in caloric.

Whatever be the nature of caloric, whether it be a body fui ments being de-generis, or a quality of other bodies, its effects are peculiar and moted by expan-appropriate; and, like all other facts, bear a proportion to the energy of their cause. Expansion, for example, it is proved by experiment, keeps pace with the actual increments of heat; and on this principle is founded the thermometer, the great. agent in the acquirement of all our ideas respecting heat, both absolute and relative. The competency of this instrument, however, to afford information of the quantity of caloric, is limited by the following circumstances:

The thermomeser indicates no ties,

1/2, The mercury of the thermometer indicates only the per indicates no absolute quantity of heat which it has itself acquired, and by no means that contained in furrounding bodies. 2dly, The scale of ex-

pansion

pansion is wholly arbitrary, commencing far from the absolute privation of heat, and falling far short of its maximum. 3dly, The caloric, latent in bodies, or chemically combined with them, has no effect on the thermometer. 4thly, The experi- and the laws dements of Dr. Crawford, though sufficient to shew that the velopes by Crawford apply expansion of the mercury of the thermometer bears a ratio to only so a small the actual increments of heat, in any temperature between the part of this scale. boiling and freezing points of water, by no means prove that this proportion holds univerfally.

Equal weights of heterogeneous bodies, it is prefumed, contain unequal quantities of caloric; and the ratio of thefe

quantities is approximated in the following manner.

Equal weights of the same body, at different temperatures, Common megive, on admixture, the arithmetical mean: but equal weights the ratios of caof different bodies, at different temperatures, afford a tempe-bric by the comrature which varies confiderably from the mean. Thus a pound + or - than of water at 100 degrees, and a pound at 200°, give the tem-the mean. perature of 150°; but a pound of water at 200°, and a pound of mercury at 100°, afford, not the mean, but a temperature confiderably higher. Hence it follows that a pound of mercury has not the power of fixing and retaining fo much caloric as a pound of water: and the fixation of more heat, by the water than by the mercury, is ascribed to the superior energy of a power inherent in both, and termed capacity for caloric.

From an extensive series of experiments Dr. Crawford in-Capacities perfers, that the capacities of bodies are permanent, fo long as manent by Crawford's hythey retain their form. Thus, the capacity of water has to pothefis. that of mercury the ratio of 28 to 1, at any temperature between 32° and 212°. The difference of capacities of bodies. it is inferred, therefore would continue the same, down to the absolute privation of temperature. Imagine, then, two bodies at this point of privation: they may still contain unequal quantities of combined caloric; for, when chemically combined, caloric does not produce temperature. On Dr. Craw- not applicable at ford's hypothesis, these comparative quantities of combined known temperacaloric, in the two bodies, may be learned by observing the tures, ratio of temperature, produced, by the addition to each, of fimilar quantities of heat. This supposition, however, is manifestly gratuitous; and the contrary might be maintained with equal or greater probability: for it may be supposed that, at this assumed negation of temperature, one body renders

not probable.

latent more caloric than another, because it actually contains less; as certain day falts attract more water, from the atmosphere than others containing much water of crystallization. The commonly employed model of afortaining the specific caloric of bodies, is founded, therefore, on an affumption. which is deficient in the character of a datum, and which itself requires proof, alternant van at the state of the period after after after after after a transfer at the state of the stat If these objections be valid, they will apply alfo to thew the

Theorem for finding the abso lute zero, fallacious.

The method.

fallacy of the theorem, for finding the abfoliate zero of bodies; By this term fome philosophers appear to understand the point of absolute privation of caloric, both free and gembined, "I apprehend, howevery that in thrich propriety it can only be mied to fignify the negation of uncombined calorie, or, as Dro Crawford expresses him felf, the point of absolute cold, "As applied; however, to water, it is evident that the whole quantity of heat is understood. In afcertaining the zero, fay thefe calculators; the capacity of ice to that of water is as 9 to 10. It is plain, therefore, that when water freezes; it must give out the of its whole heat, and this 10th part is found to answer to 1460 of Fahrenheit. Confequently its whole heat is 10 times 146, or 1460°; and bence the natural zero is 1460 - 32 or 1428°. Now of this estimate it is a datum, that the capacities of ice and water have precifely the above ratio. But if the general formula, for afcertaining the specific caloric of bodies; be

Arguments including thefe doctrines, therefore prove nothing either way.

The materiality of caloric may, I apprehend, be maintained, without admitting that we have made any steps towards determining its quantity in bodies; and the arguments of Count Rumford and Mr. Davy are not demonstrative, because they affume, that this part of the doctrine of caloric cannot be relinquiffied; without abandoning it in toto. I may be permitted. therefore, to state my reasons for believing caloric to be matter; which would liave been unnevellary; had the contrary been proved, with all the force of mathematical demonstration. in

founded on erroneous principles, it cannot ferve as the groundwork of any folid conclusions: had to a manage and tadt , and

That caloric is matter;

Avoiding all metaphyfical reasoning on the nature of matter, and affuming the generally received definition, as fufficiently characterizing it, I fliall examine how far this general character of matter applies to the individual-calorie. Caloric occupies space or is extended, because it enlarges the dimensions cupies space, and of other bodies; and, for the same reason, it is impenetrable, fince

because it expands bodies, and therefore ocis impenetrable:

fince if it could exist, at the same time, in the same place, with other bodies, their volume would never be enlarged by the addition of heat. Of form or figure, as only a mode of extenfion, it is unnecessary to prove that caloric is possessed; and indeed there is perhaps only one general quality of matter that will not be allowed it, viz. attraction. That caloric is in- It has not been fluenced by the attraction of gravitation, or by cobelive attractor of weight. tion, has never yet been proved. Yet the various experiments of Buffon, Whitehurft, Fordyce, Pictet, &c. cannot be alledged as proofs, that it is actually devoid of this property; fince they only decide, that the small quantities, which can be artificially collected, are not to be set in the balance against the grosser kinds of matter. One kind of attraction, that which has lately been termed chemical affinity, may, I think, after a full furvey of phenomena, be fairly predicated of caloric—and if its poletion of this quality be rendered probable, we hall thence derive a powerful argument in favour of its materiality.

That chemical affinity has a confiderable fhare in producing It appears to the phenomena of heat, appears probable from the following attraction; confiderations:

1. All the characters, diffinguishing caloric when separate, because its procease to be apparent, when it has contributed to a change of combination; form in other bodies; and the properties of the substances so changed are also materially altered. Now this is the only unequivocal mark of chemical union that we can apply in any instance; and chemical union implies the existence and efficiency of chemical affinity.

2. The relation of caloric to different substances, appears and it shews to observe that peculiar law, which, in other instances, is elective affinity; fingle: termed elective affinity. If a compound of two or more principles, a metallic oxide for infrance, be exposed in a high temperature, the caloric forms a perfect union with the one, but not with the other. In certain inflances, caloric is evolved when two substances, attracting each other more powerfully than they attract caloric, produce, on admixture, an elevation of temperature. In other infrances, caloric is absorbed when it is attracted by the new compound more strongly than by the feparate components. Such facts warrant the deduction, that caloric is subject to the laws of chemical affinity. But the precife order of its affinities remains to be decided by future experiments. 3. Caloric

and double or compound; 3. Caloric feems also, on some occasions, to bear a part in the operation of double elective affinities. In this way it produces decompositions, which, by single affinity, it is incapable of effecting. Thus a most intense fire does not expel intirely the carbonic acid from alkalis. But when the affinity of an acid for an alkali concurs with that of carbonic acid for caloric, a decomposition ensues.—Again, water may be submitted to the highest temperature, without imparting a gaseous form to the hidrogen which it contains; but the conspiring affinity of a metal for oxigen occasions the production of hidrogenous gas. On this principle, many chemical facts are resolved into the law of double affinity, which are, at present, explained by that of single elective attraction.

and by interme-

4. Caloric acts, fometimes, as an intermedium in combining bodies, which, without its aid, are not susceptible of combination. Thus carbon and oxigen do not evince any tendency to combination, at the ordinary temperature of the atmosphere; but caloric brings them into union, and constitutes, itself, part of the resulting compound. This, and a variety of other instances, have a striking resemblance to what is called intermediate assigning.

Capacity and affinity are not fynonimous terms.

In the theory of Dr. Crawford, no influence is allowed to chemical affinity over the phenomena of heat; and indeed that philosopher expresses a decided opinion, that elementary heat is not capable of uniting chemically with bodies. Hence it appears, that the difference between the terms affinity and capacity is not merely a verbal one; but that they are actually expressive of different powers or causes: and the question, therefore, which of these terms shall be adopted, in the description of sacts, is one involving the determination of causes.

On the word capacity as applied by Crawford,

The term capacity for heat is employed, by Dr. Crawford and others, to denote, in the abstract, that power, by which different kinds of matter acquire different quantities of caloric. But in the various applications, that are made of this theory, a more precise meaning is often affixed to it; and the term is applied, in much the same sense, which it has in common language. When thus understood, a difference of capacity necessarily implies a difference in the extent of the spaces, between the minute particles of bodies; and that these differences occasion the varieties, observed in the acquirement of

heat.

heat by different bodies. On this theory, there is no active it usually imprinciple or power inherent in bodies, and more active in plies no power but mere locafome than in others,—no tendency in the matter of heat to lity. attach itself, in preference to any one substance. The affigned cause of the phenomena of heat is not, I apprehend, adequate to produce the effects ascribed to it.

On the theory of capacities, a change of form is, in certain By this theory instances, antecedent to the absorption of caloric. Thus, the change of capacity may when ether is converted into gas, on removing the pressure precede the of the atmosphere, according to this hypothesis, the capacity transition of heat. of the ether is increased by its volatilization; and the change of form is prior to, and the cause of, the absorption of caloric. The order of events, then, in the volatilization of ether, is first an alteration of form; next a change of capacity; and lastly an absorption of caloric. On this hypothesis, ether may exist in the state of gas, without containing a greater absolute quantity of caloric, than in a liquid form. But such an interpretation of phenomena is directly contradictory to an established principle, admitted, even by those who prefer the doctrine of capacities, viz. that all bodies, during their conversion from a sluid to a vaporous state, absorb caloric. It is at variance, also, with observed facts: for if a thermometer be immerfed in a portion of ether, confined under the receiver of an air pump, the temperature of the ether will be found to fink gradually, during the exhaustion of the air; and the evaporation becomes proportionally flower, till, at last, it is scarcely perceptible. We may, therefore, infer, that at a certain point of diminished temperature, the volatilization of ether would entirely cease, if the supply of caloric, from surrounding bodies, could be completely intercepted. But on the theory of capacities, the evaporation should proceed as rapidly at the close, as at the commencement, of the process -or, in other words, evaporation should be wholly independent of temperature, which every one knows is contrary to fact.

It may be confidered therefore, as extremely probable, that Whence it is the tendency of ether to assume a gaseous form depends on its most probable that the effects chemical affinity for caloric. But, (it may be asked) how is arise from affithis affinity counteracted by an increased pressure, and aug-nity. mented by a diminished one?

A cir-

Mechanical i an argument against this docarine.

A circumstance, absolutely essential to the formation of preffure prevents gales, is, that free space shall be allowed for their expansion. gas; but is not Mechanical preffure acts as a counteracting force to this expanfion; and either prevents it completely, or partially, according to the degree of, its application. But from this fact, no argument can be drawn against the existence of chemical affinity, as an attribute of caloric. Two opposite forces in physics may be so balanced, that neither shall produce its abpropriate effect. Thus a body, impelled in contrary directions, may remain at reft, yet the operation of the oppofing forces, in this case, cannot be denied. Even in chemistry, we have unequivocal examples, in which the action of the affinities is suppressed by more powerful causes. Thus bodies, that have a firong chemical affinity, are kept perfectly diffinct; even when placed in contact, by the affinity of aggregation. The only inference, then, that can fairly be adduced from the effects of preffure, in preventing the formation of gales, is, that it is a power, fometimes superior, in energy, to that of chemical affinity.

Caloric having all the properties of matter, be taken to be matter.

Since, therefore, caloric is characterized by all the properties, except gravity, that enter into the definition of matter, but gravity may we may venture to confider it as a diffinct and peculiar body. Nor is its deficiency of gravity fufficient to exclude it from the class of material substances. Such nicety of arrangement might, with equal propriety, lead us to deny the materiality of light, the gravity of which has never yet been proved: for besides the experiments of Mr. Michell, which failed in ascertaining this property of light, we have feveral chemical facts tending to the same conclusion. Thus Mr. Cavendish, after firing a mixture of hidrogenous and oxigenous gales, in a cloid vessel, a process during which much light is always emitted, found not the smallest diminution of weight.

Phenomena of heat differ from those of motion.

To have completed this defence of the material nature of heat, it would have been proper to have pointed out the circumstances, in which the phenomena of heat differ from the known and acknowledged phenomena of motion. At prefent, however, I have not leifure to purfue the fubject at much length; and, though several points of disagreement would doubtless be found, I shall mention only one of the most marked and decifive.

Motion

Motion is an attribute of matter, independently of which Heat can pais it cannot possibly subsist. If therefore, the phenomena of heat cuum; but mocan be fliewn to take place, where matter is not prefent, we tion furely canshall derive, from the fact, a conclusive argument against that not be transtheory of heat, which affigns motion as its cause. Now, in there is no matthe experiment of Count Rumford, before alluded to, heat ter. Hence the heat is matter. passed through a torricellian vacuum, in which, it need hardly be observed, nothing could be present to transport or propagate motion This experiment, in my opinion, decidedly proves, that heat can subfift independently of other matter, and consequently of motion—in other words that heat is a diftinct and peculiar body. alcida 10

Enquiries concerning the Dilutation of the Gafes and Vapors. Read to the National Institute of France, By CIT. GAY at Luss Action of the

ART, In The Object of this Memoir.

NATURAL philosophers have been long engaged in re-The refearches searches on the dilatation of the gases; but their labours pre-on the gases are fent fuch variations in their refults, that fo far from having fixed the general opinion, they have on the contrary left us

great reason to wish for more accurate investigations.

ni dille I bennell in lu lu,

The attention of philosophers has not been equally directed The expansions to the dilatation of vapors. Though the prodigious effects of of vapor have the vapor of water has long been known and applied with the mined. happiest effect, Ziegler and Bettancourt are the only persons

who within my knowledge have attempted to measure them. Their experiments however are not adequate to shew the true dilatation of the vapor; for as there was always water in their apparatus, every new degree of heat not only produced a dilatation of the wapor formed by the preceding degrees, but also increased the volume by the formation of new vapors: two causes which evidently tend to raise the mercury in their int as Alich of a some a avenum in his boiles, the resonante

\* From the Annales de Chimie, XLIII. 187.

on this experiments and refults. See the Hydraulier

The apparatus of Bettancourt confifts in a boiler of copper closed by a cover of the same metal, through which pass three tubes.

The thermometer does not indicate the real quantities of heat.

The thermometer, as it is at prefent conftructed, cannot be applied to point out the exact proportions of heat, because we are not yet acquainted with the relation between its degrees and the quantities of heat. It is indeed generally thought that equal divisions of its scale represent equal tensions of caloric; but this opinion is not sounded on any well decided fact.

Numerous demands in philofophy and the arts for the knowledge of the expansion of gas and vapor.

We must admit therefore, that we are still far from possesfing positive knowledge respecting the dilatation of the gases and vapors, and the correspondent march of the thermometer: nevertheless, we have daily occasion in natural philosophy and chemistry to reduce a given volume of gas from one temperature to another, to measure the heat disengaged or absorbed in the changes of conftitution of bodies; or that difengaged or absorbed by the same body in passing from one temperature to another; and in the arts to calculate the effects of fire engines, or to estimate the expansions of various bodies; in meteorology to determine the quantity of water held in folution by the air, a quantity which varies with its temperature and its denfity, according to a law not yet ascertained. Lastly, in the construction of tables of astronomical refraction, and in the application of the barometer to the measurement of heights. it is indifpenfible that we should accurately know the temperature of the air, and the law of its dilatations.

The author's inducements to this work.

Though these considerations were of themselves sufficient to render the enquiry into an object of such general utility desirable, yet the difficulty of the researches it demands would have prevented my attempting it, if I had not been strongly engaged by Cit. Berthollet, whose pupil I have the honor to be. I am indebted to him for the necessary means of executing this work, in which I have often been affished by his

The first serves to introduce water, the second admits a thermometer to indicate the temperature of the vapor, and to the third is adapted a recurved barometrical tube, to measure the elasticity of the same vapor. A vacuum is made in the boiler by means of an air pump communicating through a tube provided with a stop cock.

The apparatus of Ziegler differs little from that of Bettancourt; but as Ziegler did not make a vacuum in his boiler, there is a great difference in their experiments and refults. See the Hydraulic Architecture of Prony, Vol. II.---G. L.

advice,

advice, and that of Cit. Laplace. These great authorities will add to the confidence to which my work may be entitled.

The experiments which I have undertaken on the law of The experithe dilatation of gafes and vapors, and the march of the ther-ments related momentum not being not convoleted the march of the ther-only to the commometer not being yet completed, the only object of this me-parative expanmoir will be to examine the dilatation of gales and vapors at a fion of different determined elevation of temperature, and to shew that it is at a given tem the same in all these fluids. But before I relate my experi-perature. ments, I think it proper to give a flort account of what has . been done on this fubject; and as I shall make observations on each of the methods that have been employed, I shall prefix this amount to the examination of one of the principal causes of uncertainty to which this species of experiments is liable. Although it is very important, and feems to have remained unknown to the generality of philosophers who have examined the dilatations of gafes, its influence will be fully admitted when fimply pointed out. What I shall fay of atmospheric air will apply to the other gases.

This cause of uncertainty is owing to the presence of water Water in the in the apparatus. If a few drops of this fluid be left in a re-apparatus is the ceiver full of air, and its temperature then raised to that of uncertainty. boiling water; this water when converted into vapor, will

occupy a volume about 1800 times greater than its primitive bulk, and will drive off by this means a very confiderable portion of the air contained in the receiver. It necessarily hap-Explanation of pens then, that when these vapors are condensed, and come this effect.

to occupy a volume 1800 times lefs, a dilatation much too great will be attributed to the air remaining in the receiver; because it will be supposed that it was this air which at the temperature of boiling water occupied all the capacity of the receiver. If the temperature be not raifed to this degree, the fame cause of uncertainty will not the less exist, and its intenfity will be relative to the height of temperature: for in this case the water will not entirely evaporate, but the air will dissolve more of it in proportion as its temperature is raised, and it will confequently receive a greater or less augmentation of volume, independent of that occasioned by heat; fo

that as it passes to a lower temperature, the volume of air

Vol. III .- NOVEMBER, 1802.

which occupied the whole receiver will diminish for two rea-P

fons, 1st. by the loss of its caloric; 2d. by that of the water i held in solution. Too much dilatation will be here also attributed to the air.

Other liquids, or even folids may affect the sefult.

In general, whenever liquids, or even folids, as for example, muriate of ammonia are inclosed in gases capable of disfolving them, or of evaporating at the temperature to which they may be exposed, errors in the determination of the gases ous dilatations must necessarily result.

ART. II. Short Account of what has been afcertained concerning the Dilatation of Gases.

History of the science of the expansions of gases, &c.
Amontons.

The dilatation of atmospheric air by heat was known before the time of Amontons; but it appears that this philosopher was the first who wished to know the extent for a given elevation of temperature. For this purpose he enclosed air in a ball soldered to the extremity of one of the branches of a reversed syphon, and he plunged this apparatus into a bath of hot water \*. The air dilated by the heat compressed the mercury, and elevated it into the other branch of the syphon; so that he judged of the spring of the air by the height of the mercury above the level of the ball.

From the various experiments made on unequal volumes of air he concludes: (Mem. de L'Acad. 1699, 1702.)

Amontons's refults on common air

- 1st. "That the heat of boiling water has limits, beyond which it does not pass.
- 2d. "That unequal volumes of air equally augment the force of their fpring by equal degrees of heat, and on the contrary.
- 3d. "That the heat of boiling water does not augment the force of the spring of air beyond the power of sustaining about ten inches in height of mercury."

mechanically compressed. He then proves, that however compressed a volume of air may be, the heat of boiling water always augments the force of its spring about one third; that is to say, that a volume of air compressed, for example, by a column of mercury of 60 inches, will sustain (including the weight of the atmosphere) at the temperature of boiling water, a column of mercury of

\* The air enclosed in the ball, not being able to escape when mercury is poured in, is found a little less in volume than it naturally is; but if no other pressure is desired than that of the atmosphere, it will be very easy to avoid this slight inconvenience.

about

about 80 inches. He concludes from this, "That the same Much force from degree of heat, however small it may be, can always more fmall increase of temperature in and more augment the fpring of air, if this air bears a weight condensed air. more and more great."

If Amontons had begun to reckon from a degree of heat Amontons had better determined than that which he called temperate, which hot a comparawas then hardly possible, there might have been deduced from terhis experiments a sufficiently exact dilatation of atmospheric air; nevertheless, as he made them comparatively on volumes of air of very unequal denfity, it may be concluded that however dense a volume of air may be, the augmentation of spring it receives from the same degree of heat, is always relative to that it possessed before the experiment,

Nuguet, in endeavouring to verify the refults of Amontons, Nuguet's expefound others very different. In one of his experiments, the riments differed volume of air dilated by the heat of boiling water, and the Amontons, primitive volume were as 2 to 1, and in two others as 16 to 1. His apparatus confifted of a bottle, which he plunged reverfed into a water bath, and elevated its temperature to that of boiling water. It is easily feen that this apparatus is ex-because his air tremely defective, as the air was always in contact with the was in contact water; and besides, Nuguet had suffered some water to re- with water. main in the bottle. It is no wonder then that he obtained fuch disproportioned and extravagant results. (Mem. de L'Acad. 1708. Lahire.)

This great difference between the refults of Amontons and Lahire's repeti-Nuguet on the dilatation of atmospheric air, and the confide-tion's experiration that they had submitted it to experiments under uncom-ments. mon circumstances, engaged Lahire to pursue the same object. His apparatus was the fame as that of Amontons, excepting that the ball contained a tube, which he closed after having introduced mercury. By this means, the mercury being on a level with the ball and fyphon, the air on which the experiment was made, was not more compressed than the surrounding atmosphere. With this apparatus Lahire found, that air dilated from the degree of temperate to that of boiling water, could not fuftain a column of mercury of one third the weight of the atmosphere; he then found in another, the thermometer being lower, and the barometer higher than in the first experiment, that the air dilated by the heat of boiling water, could not fustain a column of mercury as great as in

from those of

the first. These results are evidently contradictory; but Lahire not suspecting any error, concluded that it must be neceffarily allowed that the nature of air was yet undiscovered.

Lahire objects to the experiments of Nuguet on account of water.

Lahire, in order to give a reason for the difference between his refults and those of Nuguet, a difference much too great not to be owing to some foreign cause, remarked, that Nuguet had left fome water in his apparatus; and from hence he judged that it might be this water, as it changed to vapor, and expelled a portion of the air enclosed in the bottle, had presented so great a dilatation. He was entirely confirmed in his opinion by the refult of an experiment made in Nuguet's method, a little water being left in the bottle; for he found that the volume of air dilated from the degree of temperate to that of boiling water, and the primitive volume was as 35 and a half to 1. (Mem. de L'Acad, 1708.)

Stancari also obferved the in-

At the same time M. Stancari of Bologne proved, that water surved the inthese two philosophers then is owing the knowledge of the influence of water on the dilatation of atmospheric air; but although they placed it by their experiments in the clearest light, it has been fince generally misconceived. The great variations in the refults of philosophers on the dilatation of gases, may likewise be attributed to the little attention that has been paid to this influence.

The dilatations fequence in barometrical admeasurements.

It is known that elevations in the atmosphere are given by of air are of con- the logarithms of the correspondent heights of the barometric column. If the denfity of the air were always the fame, it would be eafy to determine the height of one place above any other known place by barometrical observation. Whence it becomes of importance to diffinguish the causes that may infldence the density of the air, in order to make the necessary corrections in the heights given by the barometer.

Delucon the barometric meafurement.

Deluc, who has thrown such great light on this branch of philosophy, discovered that heat was one of the causes. That he might well distinguish its effects, he first fought the temperature at which the logarithms give the heights without correction, and he found by comparing feveral observations made in fituations whose heights he had exactly determined; that this takes place at the temperature of 16° \$ of the thermometer divided into 80 parts, which he calls the fixed tempegive the height. rature. To correct the effects of heat above and below this

Temperature of which the logs.

fixed

fixed point, he compared the heights found by the logarithms with those which he had measured, by attributing to heat the excesses and deficiences of the first for the second; and he concludes "that near the fixed temperature, the correction Correction for for a degree of the thermometer is to the height of the place temperature. :: 1: 215" (Recher. fur les modif. de l'at. IV. part. ch. III.)

Colonel Roy discovered a much greater dilatation in air. Colonel Roy's According to him near the 15° of the thermometer divided allowance into 80 parts, the air dilates 172 of its volume for each degree. He likewise found that humid air dilates much more than dry air; but Saussure observes, that Colonel Roy having introduced into his manometer either liquid water, or the vapor of water, confounded two things which ought to be but he produced feparated, namely, the conversion of water into an elastic more vapor the fluid, and the dilatation of air united with such vapor. (Phil. greater his heat. Transact. 1777, p. 704.)

Saussure fixes the dilatation of air near the 6th degree, at De Saussure's Tre of its volume for each degree. His experiments were refult. made in a great receiver, in which were enclosed a thermometer and barometer, to indicate the variations of the temperature of the air, and the correspondent elasticity it would acquire. To discover the influence of water on the dilatation of air he enclosed in his receiver air more or less dry, and avoided the production of new vapors. He was so far from He avoided the finding this air more dilatable than very dry air, that he on the production of new vapor; and contrary thought very dry air was a little more dilatable than in this case very moist; but always holding its water in perfect folution. found dry air ra-(Essai sur l'Hygrométrie, p. 108.)

Till this period philosophers had confined themselves to the The gases were dilatation of atmospheric air, and the first who examined that not yet exaof the other gases was the celebrated Priestley. The following Priestley's expeis his process:

After having filled a phial of gas over mercury, he adapts The gas was reto it a bended tube, one of its branches being very inclined, ceived over merand he leaves a little of the mercury in the neck of the phial, expansion nothat the expansion of the gas may make it pass into the tube, ticed in a re-This done, he places his apparatus in a box of wood, joins a curved tube. thermometer with it, and takes it into apartments of different temperatures. The dilated air causes the mercury to pass through a greater or less space in the tube, and by this space meafured

measured in inches, Priestley discovered the dilatations of the different gases. As all these experiments were made with the same phial and tube, which he probably always inclined in the same angle, they gave a comparative result between the dilatations of the different gases, but not the absolute dilatations; because for this purpose he ought to have known the capacity of the part of the tube passed through by the mercury relative to that of the phial, and to have known exactly the inclination of the tube. Priestley says nothing of this. I will no longer then discuss his experiments; particularly as Priestley himself does not place much considence in them, and wishes them to be repeated in a better manner.

but not with great attention:

His results as to the increments only.

Supposing equal volumes of different gases, the expansions measured in inches on the tube would be for 4°, 44 of the thermometer divided into 80 parts.

								Inches.
Common air		7		•				1,32
Hidrogen gas	-		-		-		-	2,05
Nitrous gas -		-		-		-		2,02
Carbonic acid gas	-		70,	*	- "		-	2,20
Muriatic acid gas		-		-		-	-	1,33
Oxigen gas -	-			-		-		2,21
Azote gas -			7				-	1,65
Sulphureous acid gas	s			-		- 1		2,37
Fluoric acid gas	-				-		-	2,83
Ammoniacal gas				-		-		4,75

(Exper. and Obser. &c. Book VII. Sect. VI.)

Vandermonde, Monge, and Berthollet on common air and hidrogen.

Guyton and Duvernois. In a memoir printed among those of the Academy for the year 1786, Citizens Monge, Berthollet, and Vandermonde, concluded from experiment, that for one degree, atmospheric air dilates 184.83 of its volume, and hidrogen gas 181.02.

At length Cit. Guyton confidering the variations in the refults on the dilatation of atmospheric air, and that there were no direct experiments to determine the dilatation of gases at degrees of heat a little elevated or otherwise, undertook with Cit. Duvernois to examine this object. As their work is the most recent, I shall endeavour for a moment to discover what may have affected their results.

Their

Their apparatus was composed of a body with a bended Their experitube, by which means, the air expelled by heat from the body, The gas was exwas received in a jar on the pneumato mercurial apparatus. truded from a body by the heat of a bath, and be made, was plunged into a bath at the temperature of melt-this excess reing ice, and was kept down by an iron cover. The bath was ceived our mere fuccessively heated to the degrees 20, 40, 60, and 80, and they collected in separate jars the products of the dilations for each of these degrees. They then determined the volumes of air expelled from the body by measuring them in their respective jars after having reduced them to the temperature of melting ice, and they determined the volume of that remaining in the body.\* But fetting afide that their apparatus obliged to af-Objection. certain many data which must have been an objection to the exactness of their results, I remark that after the immersion The mercury in of the bended tube in the mercury, as they did not introduce the tube diffurb-new air into the receiver to displace the mercury which had entered the tube by reason of the pressure of the mercury in the trough feveral degrees of heat must be required before any bubble of air can come out; fo that had they reckoned from divisions more nearly together, as from 5° to 5° they would have concluded that from zero; the first degrees of heat do not cause the different gases to undergo any dilatation. Hence they found for the first twenty degrees a dilatation much too weak for the gases in general.

- This cause of error, tho' great, would not have inade the Their retort or results of Citizens Guyton and Duvernois so far from the truth body does not appear to have had there not been a still greater. I suspect then that their re-been dry enough, ceiver was not fufficiently dry, and that a little water might have been introduced with the gas. In fact, a decigramme of water would have been sufficient to have influenced their refults very confiderably, particularly in the higher degrees; for, as it changed into elastic sluid, it would have expelled a great quantity of the air from the receivers.

The increasing progression which they obtained in all the And from this gases is this way explained, whilst they ought to have obtained cause the expansions appeared greatest at the reduced to the temperature of melting ice. I shall observe that higher temper-Cit. Guyton expresses himself thus relative to the dilatation of atures.

\* Annales de Chimie, Vol. I.

hidrogen

hidrogen gas, (Annales de Chimie, tom, I. page 284.) "The " four products of the dilatation were in this case received in " a vessel which was surrounded with another vessel filled with "ice. Notwithstanding this, the mercury of the trough in-"dicated by the thermometer 2, 3, 4, 6, degrees above zero, " whilst the water of the bath was at the same time at 20, 40, " 60 and 80 degrees, which may have occasioned some inac-" curacy in the valuation of each of these products; but this " cannot be of great consequence, the dilatation being very "little in these first degrees."

Hence it may be inferred that these philosophers did not pay more attention to reduce the volumes of the other gases to zero; and if that was the case, this would be another cause of uncertainty in their experiments.

By comparing the volumes of gases remaining in the retort with those expelled by heat, Cit. Guyton and Duvernois found that oxigen, hidrogenous and carbonic acid gafes and atmofpheric air underwent a diminution, and they attributed the cause of it to the combinations which had taken place during the time of performing the experiments. Yet when I used very pure mercury, deprived of oxide, I could not observe any fensible action between this metal and the gases from the temperature of melting ice to that of boiling water.

(To be concluded in our next.)

### XIV.

Description of a new Secret Lock. By J. B. BERARD.

# Preliminary Reflections.

Great utility of locks.

SOME machines are so eminently useful as to require no difcussion in that respect. They are highly interesting to the whole of fociety. Locks are of this number. They are a defence supplementary to the laws, and a remedy for the consequences of immorality. From the lowest to the highest ranks in the state, there is no one who does not derive daily advan-'tage from this invention. And accordingly different artifts

Secret locks;

<sup>\*</sup> From his Melanges Physico Mathematiques. Paris, an 9.

have fuccessfully employed themselves upon them, to the production of fecret locks, exhibiting different degrees of inge-

nuity.

But it is not sufficient that this instrument should contain a ought not to adfecret difficult to be developed by the skilful method of con- mit of partakers in their secret. cealment. The proprietor of this lock must always have, as partakers with him, not only the maker, but also his friends, his domestics, and generally every person who possesses a lock constructed on fimilar principles. The question to be solved remained as follows:

To invent a lock which cannot be opened but by the pro-Enunciation of prietor, of which the construction shall be simple and easy, the problem of a defended against all internal derangements, and, lastly, ca-

pable of being readily opened by the proprietor.

By reflecting on the conditions which a perfect lock ought Condition that to include, I was naturally led fome years ago to this refult; the fecret should be variable at namely, That the fecret of the lock ought to be variable at pleafure. pleasure\*, or at least that the secrets should be so multiplied, that it should be morally impossible for the workman who conftructed the lock, or any other person, to conjecture which among them all the proprietor might have chosen at the time. I foon afterwards contrived feveral kinds of locks founded on this general principle.

By continuing to attend to this object, I foon found that Others as well others before me had arrived at the fame refult, and had pub- as the author had purfued this object.

lished locks formed on the theory of combinations.

As simplicity is one of the most essential conditions of this Simplicity an

pursuit, and there are thousands of ways of applying the prin-effential requiciples of combinations to the construction of locks, I thought it would not be without utility to explain the different refults I have obtained. For it will not be till after having composed the most happy combinations, that we can hope to arrive at that fimplicity, without which a lock of this kind would be more curious than ufeful.

I shall first describe the lock which appears to me to be the The author's best among those I have thought of. I shall afterwards com- lock to be compare it with that invented in 1777, by Cit. Reguier, and de of Reguier.

\* This principle, which is appplied in the well-known lock of Braviali, was explained to me in conversation, much at large, by him, in 1797.

fcribed

feribed as the most persect, in the art of lock-making (Serrurie) in the Encyclopédie Méthodique. I shall shew that the lock of Reguier, as well as many others, contains an impersection, or rather a radical defect, which escaped the inventor, and renders its other properties of little value \*. It will be seen how I have avoided this defect in mine. And, lastly, I shall concisely describe some other systems of locks which seem to offer particular advantages, but which, nevertheless, do not possess the same degree of persection as that to which I have given the presence.

## Description of the new Lock.

Particular defcription of the new lock: The front of this lock confifts of a plate 230 millimetres † in length, 67 in width, and 4 or 5 thick. Along the middle line are difposed five nuts or screw heads, of 18 in diameter. In the centre of these nuts is a square stem of 5 in the side, which ends in a screw.

This square stem carries immediately on the other side of the plate, a round plate of 38 in diameter and 3 thick, having a square hole in the center, and having 24 teeth in its circum-ference, with spars between them equal to the teeth.

Upon this round plate is placed a cylindrical cup or ferril, having its bottom in contact with that wheel, and connected with it by the ftem, which passes through a round hole in its center, and binds it down by the screw nut. This ferril, of which the thickness is 2 and the height about 29, has a prominent stud at its stat face, which entering between any two teeth of the wheel, serves to vary at pleasure the relative positions of the wheel and ferril at the time of screwing them together. Lastly, The serril has 24 teeth cut in its edge, having a space of about 4 between each. These teeth are about 3 in depth, excepting one of the intervals which is cut to the depth of 6. We shall hereaster explain the use of this depth.

\* Cit. Reguier has, nevertheless, the honour of being the first who nearly resolved this problem.

+ To avoid fractions, I have not reduced these measures. As the metre is 39.371 inches (Philos. Journ. II. 250), it will be sufficiently near if the reader multiplies the numbers in the text by 4, and then cuts off two figures: Thus 230×4=9.20 inches, or 9½ inches.—N.

The

The face of the lock is fitted up, as we have observed, with Construction of five screw heads. The middle stem, instead of being provided a scret lock accessible only to with a wheel and ferril like the others, has a wheel of 45 in dithe proprietor. ameter, 4 in thickness, and has 24 teeth of the depth of 3.

Fig. 1. Pl. II. reprefents the face of the lock. A B is 230, A A 67, and the five ferew heads, R, are spaced at equal distances.

Fig. 2 represents the inside of the lock without its bolt. The plan of the four ferrils with their screws, together with the middle toothed wheel and the heads of the nuts, are visible, but the four flat wheels are concealed beneath the ferrils.

The ferrils are inclosed in a rectangular box, 2 in thickness, and about 28 in height at the longer sides. The shorter are 34 in height, and have an excavation or notch in them to the depth of 8. The sides of this box are at the distance of 6 from the ferrils endwise, but only 2 as to the breadth. The distance between the middle wheel and the nearest ferrils is also 2, but that between the contiguous ferrils is 5.

In the four inner angles of the box are fixed four pillars, **PPP**, in thickness 6, in width 13, and about 35 above the plate. These pillars have at their extremities a rectangular groove for the bolt to slide in, and they are fixed by screws to the long faces of the box, and by rivets to the end plates. The square frame of the box is fastened to the face by two cocks **Z**, which receive screws.

The bolt is constructed as follows: A thin bar of iron is taken, of which the width C C, Fig. 3, is 67, and the length C D, which is variable towards D, is 310 from F to C. It is fitted with two edge bars C F, C F, of which the width is 5, and another C C, only 4. At the end which is to form the part of the bolt to shoot out, a plate F F D D is fixed, of the same breadth as C C, but in length arbitrary. The depth of the three edge bars is  $6\frac{\pi}{2}$ , and cannot be seen in the figure. That of the end plate depends on the strength intended to be given.

The two lateral edge bars C F have near their extremities a projection x x', x'' x''', which exceed the edge bar by 3, and do not come down to the face of the bolt by an interval of 3. In these four spaces the extremities of the pillar slide, and they themselves receive the projections x x', x'' x''' in their grooves.

Construction of The distances of all these points from the extremity C C of the a secret lock ac-bolt are, namely, little c x 20, c x' 60, c x" 226, c x'' 274, the proprietor. C F 310.

In order to complete the bolt, it only remains to fix its rack and its four pins corresponding with the four ferrils.

The rack is a rectangle, of which the length is M N, Fig. 5, equal to 67, the breadth or depth M M' 33, and the thickness, which could not be shewn in this figure, is 4, and may be seen in Fig. 3. This parallelopepidon is fixed perpendicularly upon the bolt at M' N', Fig. 3, by two projecting rivets K K, Fig. 5.

The rack is notched so as to form in its length M' N', 13 teeth of 5 in depth; so that it resembles a comb with very short teeth. The spaces are wider than the teeth, and they are rounded on the side which locks into the wheel,

The fingle tooth or pin which is intended to correspond to each of the ferrils, is formed of four parts, stand up at right angles, as is seen in Fig. 4. O H is the pin properly so called, rivetted perpendicularly in the bolt by the small square extremity. H. I I are two small wedges which precede the tooth. O H is 11, and I I 16.

The distance of the tooth or stud O from the line C C, Fig. 3, is 39; that of O' is 118; that of O" is 169; that of O" 248.

The little wedge which precedes the tooth, is 6 in length and 3 in height. We shall hereafter see the use of this.

From the position of these sour study it follows, that when they are lodged precisely in the thickness of the serrils, the two extreme ferrils must have their long notch or aperture directly towards the extremities of the lock, and those of the intermediate ferrils will, on the contrary, have their apertures both turned towards the middle wheel. We shall see also that it follows, that when the lock is open, the first pin O is out of the lock in the groove cut in the door; the tooth O' is within the second ferrol; the tooth O'' in the space above the wheel; and the tooth O''' within the sourch ferril. And the contrary is the case when the door is shut.

In order that the bolt may ftop at the same point when the door is open, there is a small protuberance X or X". The teeth O, O', O", O", are then sufficiently distinct from the ferrils to allow of their being moved or set.

And so likewise in order to stop the bolt always at the same Construction of point when shut, there is fixed at Q, near the short edge bar, a a secret lock accessible only to strew, the head of which stands higher than the sace of the bolt, the proprietor, and stops it by taking against the box. When the bolt is required to be taken out, this screw is drawn back a little, and it passes clear of the edge bar. The same effect might be obtained by putting an interior stem to the handle, which stall stop the bolt when the door is locked, which takes place when the points X and X" coincide with the saces of the pillars towards the knob Q. This knob is also of use to open or bolt the door within. The position of the knob Q is such, that the circumsference of its screw stem is a tangent to the line X X.

The key remains to be described: This is a circular plate or disc, Fig. 6, of 50 in diameter and 3 in thickness. Its circumstenence is divided into 24 parts, carrying circularly the 24 letters of the alphabet, leaving out K. It has a handle in the form of a semiellipsis, to turn it with the singer and thumb, and this handle carries on the other side of the plate two small steel points TT, of about 3 in length, which enter into the holes of the screw heads. Lastly, This plate has an opening in its center about 20 in diameter, to allow the points to be visible when applied to the screw heads.

There are also marked on the face of the lock five small stars or lozenges, to serve as station-points for the key; and also five small holes near one of the two holes of the screw heads. These small holes serve as an indication to apply the key always in the same manner, by putting the same point in the same hole.

The position of the four stars corresponding with the ferrils, must be determined with great care and precision. For this purpose, the ferrils must be placed so that their openings shall be in a right line, and the holes of the screw heads in a vertical line. The bolt is then to be introduced by engaging it in the grooves of the pillars, so that the sour teeth or pins may be lodged in the thickness of the ferrils. The turn-screw is then to be inserted successively in the holes of each screw-head; observing that the letter A, having its mark distinguished by a diamond, shall be turned towards the same side as the small indicative point placed on the screw-head, and then a point must be made on the sace of the lock to correspond with the letter A of the key. At this point the star is to be made, which will

ceffible only to the proprietor.

Construction of be very nearly in a vertical line: but it may easily be imagined a fecret lock ac- that it will not be exactly fo, unless the study of the bolt should exactly fill the spaces, so as to leave no shake. The fifth star corresponding to the center wheel, is merely ornamental.

> In this disposition of the lock we may affirm, that it is adjusted to the word A A A A.

> In order to change the secret, and adjust the lock to the word long, for example, the first screw-head must be turned round to the left, until the letter L of the key coincides with the star. In the fame manner the letter O is made to coincide with the flar of the fecond fcrew-head; the letter N with that of the third; and that of the letter G with the fourth. The flat wheels, remaining in that position, the ferrils are then to be disposed in a right line, and the fecret will confift of the word long. bolt is then to be put in its place, and the handle turned to advance its stop.

> When the lock is to be flut, the screw-heads must be difposed so as to form the word long; and in this position the bolt may be moved backward and forward by turning the middle If one or more of the screw-heads be deranged, the bolt is locked fast. But in order to open it, the same disposition must be made, and the bolt may again be moved.

> We may observe that, when the screw-heads are deranged, the bolt nevertheless is capable of moving through the space of 13, arifing from the fize of the ferrils; which affords an advantage for uting it as a simple bolt, to be put backwards and forwards within the chamber.

> > (To be concluded.)

### SCIENTIFIC NEWS,

Cheap Method of producing Light.

NR. EZEKIEL WALKER informs me that he has difcovered a cheap method of producing light, which feems to poffefs advantages much superior to the common modes of illumi-This light generates no fmoke, and requires not the aid of fnuffers.

# On the Reflection of obscure Heat.

C. Pictet's experiment related in his essay on fire, to prove Pictet's experithe reflexibility of obscure heat is well known. It consists in ments of reflect. placing opposite to each other two concave metallic mirrors. mirrors. In the focus of one is placed a hot but not luminous cannonball, in the focus of the other a very fensible air thermometer, and the latter is foon feen to rife rapidly.

The same philosopher has since published some other experi-

ments on the same subject.

Having employed a lighted candle instead of the cannon-Glass intercepted ball, he placed between the two foci a plate of very thin clear the heat but not the light. transparent glass, and which intercepted the light very little: the thermometer indicating the transmission of heat, stopped that instant.

The two mirrors were placed at the distance of 25 metres Distance of mir-

(yards) one from the other, in order to determine whether the rors 25 metres. time of the propagation of the radiant heat from one focus to the other could be appreciated. A heated but not luminous ball was suspended at one of the foci, before which a screen was placed. At the infant that this obstacle was removed, the fluid of the thermometer, which was before perfectly at rest, began to move, and no fenfible interval could possibly be perceived between the suppression and the effects of the transmitted heat.

C. Pictet relates this experiment in the Bibliothèque Britan- The heat was nique, in support of the opinion he had advanced in his essay instantly transon fire, that light and heat are not the fame. This opinion has fince been renewed by M. Herschell.—Bulletin des Sciences. No. 62.

On certain Facts commonly urged against the Doctrine of two Electric Fluids. By Cit. TREMERY, Engineer of Mines. (Bull. des Sciences, No. 63.)

Among the facts which have been offered in support of The experiment Franklin's hypothesis of a single electric sluid, the most re- of passing a markable is the following, (commonly called Lullin's card, Lullin's card, from the inventor of the experiment).

Having placed between two metallic conductors a card, which touches each of them, by its opposite faces, in two different ductor.

different points, a ftrong electric charge is passed through the apparatus. At the inftant of the operation, a luminous flash passes from the positive conductor, slides along the surface of penetrated oppo- the card, and perforates it opposite the negative conductor. fite the neg. con-This happens even when the card is perforated beforehand.

opposite the positive conductor.

From this fact it has been concluded, that in order to support the theory of two fluids, we must suppose that one of them escapes from bodies and produces light, while the other remains inherent \*. Cit. Tremery destroys this reasoning by the following experiment:

This does not happen in rarefied air.

He places the card and the two conductors under the receiver of an air pump, and in proportion as the air is exhausted, the place of fracture happens nearest the positive conductor; and when the air is half exhausted, the point is precifely at the middle between the two conductors. At every discharge a luminous trace issues from each conductor along the furface of the card to the place of interfection.

Inferences.

From this experiment, Cit. Tremery concludes that the atmospheric air, in its ordinary state, resists the passage of the negative more than the positive sluid t. These resistances are diffinguished in different ratios, with regard to the two fluids as the air becomes less dense, and more rapidly with regard to the negative than the positive.

Hence Cit. Tremery deduces the general refult, that the infulating property of non-conductors cannot be the same for both electricities.

From this explanation, he thinks it eafy to reconcile the theory of two fluids with the few facts opposed to it by his adverfaries.

\* Or if one fingle pass from one conductor to the other, the attraction or power of the receiving conductor will be very oblique to the card at first, and most direct at last; whence the place of fracture is inferred to be at the receiving or minus extremity.

† Yet a negative point discharges its electricity more easily than 2 positive. See Fig. 5, Pl. VIII.

W. N.

# Â

# JOURNAL

o F

# NATURAL PHILOSOPHY, CHEMISTRY,

AND

# THE ARTS.

DECEMBER, 1802.

### ARTICLE I.

Description of an Improvement of Woulfe's Apparatus, by which the great Inconveniencies of that Method of connecting the Vegsels are obviated. In a Letter from Mr. John Murray.

Edinburgh, Sept. 18, 1802.

### To Mr. NICHOLSON.

SIR,

I SEND you a drawing of an improvement in Woulse's apparatus, which I believe I may recommend with considence to chemists.

In the apparatus on the common construction it is extremely It is scarcely difficult, I might perhaps say impossible, to have a series of possible to connect the state of the possible to connect the state of the possible to connect the state of the possible to connect the necessity of employing luting to close the joinings, without luting an operation extremely troublesome when done with accuracy. This difficulty of sitting the bent tubes to the bottles by grinding, arises from the construction of the apparatus itself; for if The cause why connected in the state of the tube be ground into the bottle from which it issue, it will be found scarcely possible to grind the other leg with accuracy into the other bottle, so as to sit it at the precise to the point. III.—December, 1802.

point at which the two bottles shall be on a level; and this difficulty is increased still more when a series of bottles are to be joined in this manner. The difficulty is indeed fo great; that an apparatus of this kind perfectly air tight, though it would be of the greatest advantage in many chemical operations, has perhaps fearcely ever been procured, and the attempt to have it made was relinquished even by Lavoisier.

Luting is most

Dr. Hamilton's excellent apparatus.

Various attempts have been made to obviate this inconvecommonly used nience, but with so little success, that the original apparatus in which the joinings are closed by a lute, is still the one employed by chemists. Of the improvements that have been proposed, I may remark that that by Dr. Hamilton, of which a description is given in his translation of Berthollet on Dyeing, and which does not feem to be fufficiently known to chemists, is by far the most convenient. It may be simplified by having the bent tubes to iffue from the receivers instead of being fitted to them by grinding, and for many purpofes it answers exceedingly well. Its principal deficiency is, that no great degree of pressure can be obtained in it, proportioned to the quantity of water put into the receivers.

Cit. Girard's improvement . very expensive, and why:

Cit. Girard has lately proposed another improvement, of which an account has been given in the 4th Vol. of your Journal (p. 41.) On ordering his apparatus at the glass manufactory, I found that it could not be made but at a very great expence. This was stated to be owing partly to the trouble which would attend the bending the long tube attached to the bottle, but still more to the difficulty of giving this tube the fame curvature, as the larger tube fixed in the bottle, into which the former is to be introduced. The execution of this by the mode Cit. Girard described, would not be so easy as he appears to have supposed it; and a number of bottles would probably have required to have been made, to obtain a feries of three or four adapted to each other.

This difficulty obviated by the author.

It fometime afterwards occurred to me, that every difficulty might be obviated by a more fimple method, which I have fince found is extremely eafy in the execution. The apparatus constructed on this method is represented in the annexed drawing. Plate XIII.

A is an adopter ground into the bottle B, and connecting a Description. The contrivance retort with it. C is a straight tube ground into an opening of Girard apin the fide of B, and into a fimilar opening in the fide of the plied perpendibottle bottle D. E is a bent tube passing from the neck of D, and cularly at the fitted to it by grinding; and it is in the mode of connecting woulfe's tubess this with the succeeding bottle G, that the present improvement confifts. F is merely a common Woulfe's bottle with two necks. But in the first neck a tube G is soldered, when the bottle is made, which descends to within 11 inch of the bottom, the height of the bottle being fix inches. The long leg of the tube E descends into this tube, reaches beyond the extremity of it, and has at its termination a flight curve. It is obvious that when water is put into the bottle F, so that the extremity of G is immerfed in it, if any vapour or gas pals from D, to F by E, it will be discharged from the extremity of the latter, and from the degree of bending will be thrown beyond the extremity of G, and as this is immerfed in the fluid, the gas cannot escape, but whatever part of it is not absorbed, must be collected in the upper part of F. It passes in a similar manner to H, and from H to I, being in each bottle transmitted through a portion of water. Any permanent gas, not absorbed, passes through the bent tube sitted to I, and may be collected in a jar placed in a small pneumatic trough.

The advantages of this apparatus are obvious. All the Advantages of joinings are air tight without lute, and the tubes are at the the apparatus. It is air tight fame time fo far free, that they are in no danger of being and fafe. broken by any flight agitation or derangement. In making it, it is most convenient to have the tubes first ground in, and when perfectly dry to have them bent by the blow pipe.

The first bottle in the apparatus being defigned to receive Observations any fluid condensed in it during distillation, has no tube of fafety, because at the commencement at least of the distillation, there would be no fluid in which its extremity would be immerfed. On this account it is requifite that this first bottle should be connected with the second by a straight tube, as represented in the drawing, and not by a bent tube as in the other bottle, else upon any condensation in the retort or first bottle, the liquor would rush backward from the second. this fecond bottle a tube of fafety is fitted by grinding, which is necessary to guard against condensation. In the other bottle this is not necessary, as the open tubes are sufficient to admit air.

The only inconvenience in this apparatus is, that the fluid This apparatus in the bottles is liable to be forced up in the open tubes, in liable to have Q 2 the the fluid pressed upwards between the tubes.

the tube G for example in the bottle F, by the pressure of the gas accumulated in it from the refistance of the column of water in the other bottles; and this even proceeds fo far that part of the liquor is forced out entirely. This can in some measure be obviated by diminishing the quantity of fluid in the last bottles, or by introducing into them little more than isfufficient to immerse the extremities of the tubes. In this way however the pressure obtained, by which the absorption of a gas is promoted, is very inconfiderable. The fame inconvenience is present in the apparatus of Cit. Girard, and can be obviated only in the fame mode. In the present apparatus, another improvement may be introduced, by which it is more effectually removed, that of having the straight tubes which are foldered into the bottle of a greater length. This is represented in the drawing at K L, the open tube A rifing five or fix inches above the neck of the bottle. It is rather more difficult to folder in the tube in this mode, but the advantage from the increased height is so great, that the apparatus should perhaps always be constructed in this manner.

Remedy, by lengthening the external tube.

How to convert a common Woulfe's apparatus into the prefent.

Instead of fixing the straight tubes into the bottles when they are made, they might afterwards be fitted in by grinding, and then a common Woulse's apparatus could be easily converted into this improved one. The other mode however is preferable, as we are more certain of having the apparatus air tight, and it is so easily done that it adds little to the expence.

I am, with respect, your's, &c.

J. MURRAY.

Observations and Experiments upon Oxigenized and Hyperoxigenized Muriatic Acid; and upon some Combinations of the Muriatic Acid in its three States. By RICHARD CHENEVIX, Efg. F. R. S. M. R. I. A. From the Philosophical Transactions for 1802.

(Concluded from Page 185.)

Third Species. Hyperoxigenized Muriate of Barites.

THE earthy bases seem to follow, in the order of affinities Hyperox. mur. for this acid, at a great distance from the alkalis. They are of barites. all superfeded by the two just mentioned; and it is much more difficult to accomplish their union with the acid, than is the case with potash or soda. The most advantageous method is, to pour warm water upon a large quantity of this earth, procured by Mr. Vauquelin's method; and to cause a current of oxigenized muriatic acid to pass through the liquor, kept warm; fo that the barites already dissolved being saturated, a fresh portion of it may be taken up by the water, and presented in a state of great division to the acid. This salt is soluble; cryin about four parts of cold, and less of warm water. It cry- stallizable. stallizes like the muriate of this earth; and resembles it so much in folubility, that I could not separate them effectually by crystallization repeated several times. At first, indeed, I Purification of despaired of ever obtaining any of the earthy hyperoxigenized earthy hyperoximuriates; (not muriates in a state sufficiently pure for analysis. If we con-by diff. of solufider them as a genus distinct from the alkaline hyperoxige-bility.) nized muriates, a leading character may be, their great refemblance to their respective species of earthy muriates. thought, however, that I might, if not by direct, at least by double affinity, decompose the one without the other; and phosphate of filver occurred to me as the most likely agent. If by phosphate of filver, which by phosphate of silver be boiled with muriate of lime, of harites, double affinity &c. a double decomposition ensues; and muriate of silver, decomposes the together with phosphate of the earth, both infoluble, are pre-nuriates, and leaves the hycipitated. To increase the action, the phosphate of filver may perox. muriates be distolved in a weak acid, such as the acetous; and, though in solution the earthy phosphate be at first retained in solution, it will be

**feparated** 

be free from copper.

The filver must separated by expelling the acid. The only condition absorb lutely necessary is, that the filver employed be free from cop-For, in preparing phosphate of filver by phosphate of foda, and by nitrate of filver thus impure, copper would be thrown down by the phosphoric acid; and the phosphate of copper would be afterwards decomposed by muriate of lime. Muriate of copper would therefore remain with the earthy hyperoxigenized muriates; or, what is still worse, a part of the muriatic acid being eafily expelled from oxide of copper, the hyperoxigenized muriatic acid would be driven off from Properties of the its basis, by the more powerful agency of the former.

falt has all the properties enumerated as belonging to the ge-

earthy hyp. ox. mur.

frequently than

those with alk.

bafis.

nus of hyperoxigenized muriates; and, with heat, the acid is expelled by all acids above the benzoic. I had hoped that, without diffillation, I could procure the acid from the falt by means of fulphuric acid, which would have left an infoluble falt with barites; but hyperoxigenized muriatic acid is so easily decomposed by light, that I have not yet obtained it, to my emit light more satisfaction, disengaged and pure. A fact well worthy of attention is that the stronger acids difengage this acid with a flash of light, more frequently from the earthy than from the alkaline hyperoxigenized muriates; a phenomenon which, I suppose, depends upon the relative proportionate affinities, and confequently the greater rapidity of the difengagement But, where all is hypothesis, it is useless to draw any inference from a finole fact

Component parts, hyp. ox. mur. barites.

Hypero	xigen	izec	l mu	riati	c ac	id			4		47
Barites			-	11	1		4	3		Ų.	42,2
Water		4		)	-		(II.				10,8

Hyp. ox. mur. frontia.

Fourth Species. Hyperoxigenized Muriate of Strontia.

Deliquescent, and more foluble in alcohol than the muriate.

The foregoing observations apply to the formation of this falt, to the mode of obtaining it pure by phosphate of filver, to its conduct with the acids, to the rank of its acid in the order of affinities, and to its other properties. It is deliquefcent; and more foluble in alcohol than muriate of strontia. melts in the mouth immediately, and produces cold. Rals assume the shape of needles.

It is compo	fed of,												Comp. parts
Hyperoxi	igenize	d n	nuria	tic	acid		•		-		,  -	46	hyp. ox. mur. of ffrontia.
Strontia		,	-	-		-		•		-		26	Or anjoinings
Water				Ŧ	-		•		•		-	28	
		. 1										-	
												100.	

Fifth Species. Hyperoxigenized Muriate of Lime.

Hyp. ox. mur. lime very deli-

This falt is obtained pure, in the same manner as the other quescent, and earthy falts. It is extremely deliquescent; liquifies at a low soluble in alcoheat. by means of its water of crystallization; and is very soluble in alcohol. It produces much cold, and a sharp bitter tafte in the mouth.

It is compol	ed of,				9		. 110 18
Hyperoxi	genized	muri	atic ac	id			55,2
Lime		-				**	28,3
Water	-	-	-	•	•		16,5
							1.0

Comp. parts hyp. ox. mur. of strontia.

100,0.

Sixth Species. Hyperoxigenized Muriate of Ammonia.

Hyp. ox. mur. ammonia.

From the property which oxigenized muriatic acid possesses The ox. mur. of decomposing ammonia, this combination may be thought acid decomposes paradoxical. For, how can an acid much more active than union of the oxigenized muriatic acid exist with ammonia, which is de-hidrogen and ftroyed by the latter? But this argument may be opposed by oxigen; because this last attracthe sum of affinities that act in either case. If the affinity of tion, &c. excomposition of oxigenized muriatic acid and of ammonia, to ceeds thosewhich might form ox. gether with the affinity of oxigenized muriatic acid for ammo-mur, of ammonia, to form oxigenized muriate of ammonia, be not more nia. powerful than the affinity of oxigen for hidrogen, of azote for ox. mur. acid caloric, and of muriatic acid for ammonia, the divellent affi-does not; benities will prevail; and this is what actually happens. But, tion of compos. although oxigen may be held with less force of attraction in to form hyp. ox. oxigenized than in hyperoxigenized muriatic acid, yet the affi-is ftrongeft. nity of the latter acid for ammonia may increase in a much This salt formed greater ratio, and favour the quiescent affinities. If carbonate by double affiof ammonia be poured into any earthy falt of this genus, a nity: e. g. car-bonate of amm. double decomposition takes place; and hyperoxigenized mu- + hyp. ox. riate of ammonia is formed. This falt is very foluble in water, mur. of earth and in alcohol. It is decomposed at a very low temperature, earth and hyp.

afford carb. of and ox. mur. of am. It is very foluble in water and alcohol, and decomposed at lowing to the great quantity of oxigen contained in the acid; it temperature. being more than is necessary to combine with the quantity of hidrogen contained in the alkali, and therefore some of the

Component parts acid is difengaged, without decomposition. All the attempts I have made to ascertain the proportions of its principles, have been fruitless. The formation and existence of this salt, as I before said, are very strong proofs of what I have advanced respecting the state in which hyperoxigenized muriates at first exist; and very fully prove the different degree of assimity exercised by each acid toward the basis.

Seventh Species. Hyperoxigenized Muriate of Magnesia.

Hyp. ex. mur. of magnetia.

Its chemical and physical properties are nearly the same with those of the 5th species, only that, in addition to the other bases, lime and ammonia cause a precipitate in this salt.

Component parts. Its proportions are,

em	Zec	1 11	ıuı	iai.	ic a	CIQ				-		-	60
	-		-		_		-		-		-		25,7
													14.3
-				-		-		-		-		-	14,0
						t.		-		,			
												-	_
	eni	genize	genized n	genized mur	genized muriat	enized muriatic a	enized muriatic acid	genized muriatic acid	renized muriatic acid	enized muriatic acid	renized muriatic acid	enized muriatic acid	enized muriatic acid

Eighth Species. Hyperoxigenized Muriate of Alumina.

Hyp. ox. mur. of alumina.

It is decomposed by phosphate of filver, in the usual way, I sound nothing in by phosphate of solution but hyperoxigenised muriate of silver \*; and all the filver, hyperoxigenized muriate of alumina had been decomposed.

is deliquescent. This falt, however, appears to be very deliquescent, and is and soluble in alcohol.

This falt shall be particularly mentioned and described in another part of this Paper. For the present, it is sufficient to say, that it is very soluble in water; and, in that property, as in many others, is totally different from muriate of silver.

foluble

100,0

foluble in alcohol; but I could not ascertain the proportion of its principles, because I did not obtain it sufficiently free from the fimple muriate.

Ninth Species. Hyperoxigenized Muriate of Silica.

Hyp. ox. muze filica does not

I am inclined to think this falt does not really e xist A cur-exist. rent of oxigenized muriatic acid, fent through fome filica which had been precipitated from an acid by ammonia, and collected moist from the filter, did not feem to dissolve any portion of it. In all barites and ftrontia, prepared according to Mr. Vauquelin's method, a portion of filica from the crucibles is attacked, and taken up, by whatever acid those earths may afterwards be dissolved in: and, in all potash of commerce, there is some filica; but I have never perceived that any portion of this earth had been dissolved by this acid.

The very small portion of earth which, in attempts to form General observathe different species of this genus of salts, is taken up by acids, hyp. ox. mur. and the still smaller portion of the salt so formed, which is really in the state of hyperoxigenized muriate, render the operation fo tedious, that I have confined myfelf to form what was necessary to determine their analysis, in such a manner as I believe to be nearly accurate. It cannot, therefore, be expected that I make myself responsible, without a right of appeal to further experiments, for the accuracy with which the crystalline forms, and other physical properties, may have been stated. It is impossible to obtain fatisfactory crystals from a very small portion of falt; and I have attached myself more particularly to chemical than to physical characters, as being a much more important and certain mode of determination. For the fame reason, I have not examined the combination of the new and rarer earths with this acid. not doubt, that whatever chemist undertakes a further investigation of these extraordinary bodies, will be amply repaid for his labour.

I have mentioned, in a former part of this Paper, that all All muriates muriates lost a portion of their acid at a red heat. I exposed by ignition. one hundred parts of muriate of potash, in a crucible, to a red heat, for some minutes, and found that they lost five. I diffolved them in water, and they manifested alkaline properties. Treated by nitrate of filver, they gave a precipitate, which shewed one per cent. less of muriatic acid, than 100

ter of crystallization is inaccurately determined. If the acid of the falt be determined by precip. with filver before and may be known how much of the lofs by the latter process is acid, and how much water.

parts of the same salt that had not been exposed to fire. A violent heat may be necessary to expel the last portion of water of crystallization from certain salts, as we know particularly Whence the was is the case with sulphate of lime. But, if any of the acid can be expelled at the same temperature, there is no longer any certainty. The quantity of water, as stated by different chemists, varies much; and, from some experiments I have made, I do not believe it to have been accurately determined. The method I used to ascertain this, was as follows: I exposed a given quantity of the salt to a violent heat, and noted after ignition, it its loss of weight. I then precipitated, by nitrate of filver; and thus knew, how much the quantity of muriatic acid which this falt contained, was less than that in a like portion which had not been exposed to heat. I subtracted the difference in this quantity, from the total loss of weight in the falt exposed to heat; and the remainder I confidered as water. It was upon refults obtained in this manner, that I founded many of the proportions I have given in this Paper,

Observations on elective actractions.

It is stated in the tables of Bergman, corrected by Dr. Pearson, that lime and strontia prefer acetous to arsenic acid. But arfenic acid can expel hyperoxigenized muriatic acid from its basis, although the acetous cannot act in the same manner; therefore, this order of affinities is erroneous, It was not till lately, that we had potash and soda so pure as to be relied upon in delicate experiments; and it is not furprifing that we find mistakes with regard to their taking the acid from barites, strontia, and lime. But real potash and soda both precipitate even barites from hyperoxigenized muriatic acid. If ever it becomes eafy to obtain hyperoxigenized muriate of barites, we may prepare that earth from it in the humid way, and more near to purity, than in the method proposed by Vauquelin.

Metallic muriatic falts.

# METALLIC COMBINATIONS OF MURIATIC ACID, IN ITS DIFFERENT STATES.

The action of hyperoxigenized muriatic acid upon metals, is, as may well be expected, rapid, and without difengagement of gas. It appears to diffolve every metal, not except-Hyp.ox. mur. acid diffolves all ing gold and platina. If the metal be prefented to the acid metals: inflames at the moment when it is difengaged from the falt, inflammathem and protion enfues; and the phenomena of light and heat vary acduces muriates. cording

cording to the metal; but the falts thus produced are merely muriates. In order to form real hyperoxigenized muriates, It must combine it is necessary to take the metal in its fullest state of oxidize-mum oxides, to ment, and combine it with the acid, either by double decom-produce h. ox. position, or by passing a current of oxigenized muriatic acid muriates. gas through the oxide suspended in water. The acid is thus separated into muriatic and hyperoxigenixed muriatic acid; and, in these states, combines with the metallic oxide. The metallic hyperoxigenized muriates are different, in every re- These greatly spect, from the metallic muriates. Red oxide of iron is dif-riates. folved with difficulty. Oxide of copper more easily. Red Habitudes of oxide of lead exhibits the same appearances, during its combi
Iron. Coppers nation with this acid, as with nitric acid. When nitric acid is Lead. poured, even in excefs, upon red oxide of lead, only a part of the oxide is dissolved, unless heat be applied; and what remains becomes a blackish brown powder. But, if metallic Red oxide of lead be added, in a just proportion, all the red oxide difap-lead is too oxigepears, and none of the brown powder is formed; neither is wholly diffolved there any difengagement of nitrous gas, when the metallic in nitric acid. lead is diffolved. The precipitates caused in either case, by more oxigenated pouring an alkali into the nitric folution, are yellow. Hence and the other it appears, that red oxide of lead contains too much oxigen to part diffolves. be diffolved by nitric acid. One part of the oxide takes up the excels of oxigen, and becomes brown; while the portion which loses oxigen, becomes yellow, and is foluble in nitric acid. The presence of metallic lead promotes the total solution of the red oxide, by taking up the superabundant oxigen. I found that a current of oxigenized muriatic acid gas, like the Ox. mur. acid nitric acid, diffolved a part of the red oxide, and caused the acts similarly. brown powder to be formed, upon which it could not act. Hyperoxigenized muriate of lead is much more foluble than Hyp. ox. mur. muriate of lead; and the acid is very flightly attracted by the of lead: foliable. basis.

But, of all the metallic falts formed by the combination of Mercurial muthe muriatic acid, in any of its different states, none so much riates. deferve attention as those which have for their bases, the oxides of mercury. The nature of the falts which refult from the combination of common muriatic acid with the different oxides of this metal, has been stated in the most contradictory manner, by different chemists. But, as the knowledge of hyperoxigenized muriatic acid has thrown fome light upon the

Calomel and corres. sublimate. true state of calomel and corrosive sublimate\*, I must beg leave to dwell at some length upon this important part of my subject.

It would be useless to repeat the opinions of the old authors, who have treated of corrosive sublimate and of calomel. They are to be found in the works of those respective chemists, and I must refer to them for particulars.

Opinions refpecting corrofive fublimate,

In the Memoirs of the Academy of Sciences of Paris, for 1780, we find a Paper of Mr. Berthollet, upon the causticity of metallic salts; in which he appears to think, that the acid in corrosive sublimate is in the state of what was then called dephlogisticated marine acid. In 1785, when he had examined the oxigenized muriatic acid with more care, he renounced his former opinion; and gave the reasons why he no longer adhered to it. Some late experiments of Mr. Proust shew, that this chemist thinks as Mr. Berthollet now does. And these may be ranked among the first of modern authorities.

Foureroy confiders it as hyperox. muriate of mercury;

Notwithstanding those opinions, Mr. Fourcroy, in his Systeme des Connoissances chimiques, still considers corrosive sublimate as a hyperoxigenized muriate of mercury; and designs it throughout by that name †. This chemist, one of the founders of the methodical nomenclature, is too well acquainted with its principles, to apply the term hyperoxigenized muriate to any thing but a combination of hyperoxigenized muriatic acid. It is evident, therefore, that he considers the portion of oxigen, which, in equal quantities of corrosive sublimate and calomel, is greater in the former, to be combined with the acid, and not

<sup>\*</sup> I regret very much, that I am under the necessity of using these unmeaning terms. But the French nomenclature has made no distinction between salts formed by metallic oxides in different states of oxidizement, except by the colour, which is an extremely defective and unmeaning method. At all events, this metal is so uncomplaisant as to retain the white colour, in its different oxides combined with muriatic acid. I prefer, however, using the old name, to proposing any provisional substitute that might be found defective. This will be farther explained in Remarks upon Chemical Nomenclature.

<sup>+</sup> I have faid before, that this acid was talked of by many chemilts, as if the existence of it had really been proved.

with the oxide of mercury. As foon as I have flated fome but erroneously. experiments that prove Mr. Fourcroy's opinion to be errone-For, ous, and endeavoured to establish the analysis of corrosive fublimate and of calomel, I shall take notice of a falt hitherto unknown, which really is hyperoxigenized muriate of mer-Cury.

I took a portion of corrofive fublimate, and precipitated by Corrof. fub: prepotash. The liquor was filtered; and, upon being tried, cipitated by potash. nothing but muriate of potash was found. No re-agent of potash and no could discover the smallest trace of hyperoxigenized muri-hyp. ox. m. acid.

atic acid.

Sulphuric, nitric, phosphoric, and many other acids, poured The acids difenupon corrosive sublimate, did not disengage either muriatic, or gaged nothing hyperoxigenized muriatic acid. Nitrate of filver, poured into a folution of corrofive fublimate, gave an abundant white precipitate.

From these experiments it is evident, that muriatic acid, not Confeq. muriahyperoxigenized muriatic acid, is combined with the oxide of m. acid exists in mercury in corrofive fublimate.

To determine the proportions of this falt, I took one hundred Component parts, and precipitated by nitrate of filver. I then took another tation with pothundred, and precipitated by potash. The result of these two ash and with niexperiments was fuch as to establish the proportions of corro-trate of silver. five fublimate as follows:

Oxide of mercury -Muriatic acid -18 100.

But, the acid of this falt not being charged with a fuper- To measure the abundance of oxigen, we must look for the excess in the me-oxigen, 100 grs. tallic oxide. I took 100 grains of mercury, and dissolved them dissolved in niin nitric acid; then poured in muriatic acid; and, at a very tric acid; mur. gentle heat, evaporated to dryness. I afterwards sublimed, poured in, and in a Florence flask, the falt that remained, and obtained the falt (cor. 143,5 of corrofive fublimate. But, 143,5 of corrofive fubli-fublimed = mate contain 26 of acid; which will leave 117,5 for the 143.5; of which mercurial oxide; and, if 117,5 contain 100 of mercury, 100 acid, and confeq. of the oxide will contain 85. Therefore, the oxide of mer- 15 per cent. of cury in corrofive sublimate, is oxidized at the rate of 15 per the oxide was cent.

Component parts of calomel.

To determine the proportions in calomel, I dissolved 100 grains of it in nitric acid. The phenomena of the folution have been fo accurately described by Mr. Berthollet, that I shall not repeat them. I precipitated by nitrate of filver; and obtained a quantity of muriate of filver corresponding with 11,5 of muriatic acid. The oxide of mercury I obtained apart. Therefore, calomel is composed of,

Oxide of mercury	14	/ 70+		88,5
Muriatic acid			-	11,5
	7		-	-
				100.0.

Calomel was converted into cor. fubl. by mur. acid, evapor. and fubl. It had gained weight, of which, from the former expts. the proportions of acid and of oxigen were known;

To afcertain the flate of oxidizement of the oxide in calomel, I took 100 grains, and boiled them with nitro-muboiling in nitro-riatic acid; then evaporated very flowly, and sublimed as above. The calomel was totally converted into corrofive fublimate, and weighed 113. But 113 of corrofive fublimate contain 20,3 of muriatic acid, of which, 11,5 were originally in the calomel. The total addition of weight was 13. But the quantity of acid in these 13, amounts to 20.3 - 11.5 =8,8. Therefore, 13 - 8.8 = 4.2, remain for that part of the additional weight which is oxigen. On the other hand, 100 of calomel contain the the same quantity of mercury as

portion of oxigen in the oxide 10.7 per cent.

whence the pre-113 of corrofive sublimate, = 79. These 79, with 11,5 of acid, are equal to 90,5, and leave 9,5 for the quantity of oxiis deduced, viz. gen contained in calomel. It would appear, from these experiments, that corrofive fublimate contains 6,5 per cent. more acid, and but 2,8 per cent. more oxigen, than calomel. But this quantity of oxigen is combined with a much greater proportion of mercury; and forms an oxide of a very different degree of oxidizement. For, 88,5:9,5::100:10,7. Therefore, we may establish the following comparative table:--

CALOMEL.	CORROSIVE SUBLIMATE. Comparative
The oxide of mercury in calo- mel is composed of,	The oxide of mercury in corro-flatement. five fublimate is composed of,
Mercury 89,3	Mercury 85
Oxigen 10,7	Oxigen 15
100,0.	100.
	And corrofive fublimate is com- posed of,
Mercury79 (oxide of)	Mercury 69,7 (oxide of)
Oxigen 9.5) mercury (88,5)	Oxigen 12,3 mercury \ 82
Mercury 79 Soxide of Oxigen 9,5 mercury 88,5 Muriatic acid - 11,5	Muriatic acid - 18
	Apparit Company Company
100,0.	100.

These proportions are different from those given by Lemery, Geoffroy, Bergman, &c. But, without calling in question the accuracy and skill of these chemists, it is fair to affert, that the pure materials used by modern chemists, are more likely to lead to fure refults, than the impure re-agents of the ancients.

In these falts we find another instance, that, in proportion as The most oxig. metallic oxides contain a greater quantity of oxigen, they re-oxides faturate the largest quanquire a greater quantity of acid to enter into combination with tity of acid. them.

The method I have followed, to ascertain the proportions Observations to just stated, may appear, at first view, not to be the shortest that thew that the I might have adopted. But I have tried others, and I have ceffes were found found none so accurate. It is impossible, synthetically, to the most accuconvert a given quantity of mercury into calomel, in such a rate and concise. manner as to be certain that none of it is in a different state from that required. And, if we would attack calomel analytically, the action of the alkalies, without which we cannot proceed, is fuch as to alter the nature of the oxides. I have also made many comparative experiments, by dissolving calomel in nitro-muriatic acid, (which converted it into corrofive fublimate,) and then precipitating by ammonia; but I have not found these trials so successful as those I have described. The nature of the precipitate from corrofive sublimate by ammonia, certainly differs, according to the excess of acid that may be prefent; and mercury feems to have the power of exifting in many degrees of combination with oxigen. The only precaution

precaution absolutely necessary, in this mode of operating, is; that while the mercurial falt is in an open vessel, it should not be exposed to a degree of heat capable of volatilizing any part

London Pharmacopæia prescribes too much mercury in the formula for cor. fub. But this is prudent.

The quantity of mercury ordered in the London Pharmacopæia, to convert corrofive sublimate into calomel, is 9 pounds of mercury for every 12 pounds of corrofive sublimate. But, from the above experiments, it would appear, that a smaller quantity of mercury might strictly answer. However, from the refults of minute investigation, we should not conclude too hastily upon preparations on the great scale; and, I rather think, that the excess of mercury ordered by the Pharmacopæia is a ufeful precaution.

In attempts to reduce the mercurial falts by other metals, threw down mercury, and copper precipitated calomel from cor. fub.

In my experiments, I attempted to reduce, by means of copper, iron, or zinc, the mercury contained in the mercurial falts. Iron did not answer the purpose: zinc precipitated the iron failed, zinc mercury a little better; and copper produced a change which I did not expect. If a bit of copper be put into a folution of corrofive fublimate, a white powder shortly falls to the bottom; and that powder is calomel. When washed, it does not contain an atom of copper, or of corrofive fublimate.

Calomel is the fame whether produced in the humid or dry way: and alfo cor. fub. Observations on Scheele's humid mel. It is not pure, but contains a sub-ni-

Before I conclude these considerations, I must say, that whether calomel be prepared in the dry or in the humid way \*, iŧ

\* By the humid way, I do not mean precifely the method of Scheele. That chemist desires us to boil the acid with the mercury, process for calo- after they have ceased to act upon each other at a low temperature. By this method, the nitric acid takes up an excess of mercurial oxide; and the nitrate of mercury thus formed, precipitates by water. trate of mercury. Therefore, when this nitrate of mercury is poured into the dilute folution of muriate of foda, according to the formula of Scheele, the action, on the part of the folution, is twofold.

> 1/t. The water acts upon one part, and precipitates an oxide, or rather an infoluble fub-nitrate of mercury. And,

> 2dly. A double decomposition takes place between the nitrate of mercury and the muriate of foda. It is with reason, that the medical world have supposed the calomel of Scheele to be different from that prepared in the humid way; for it is, in fact, calomel, plus an infoluble fub-nitrate of mercury. In the first part of Scheele's procefs, there is disengagement of nitrous gas, together with oxidizement and folution of some of the mercury. When he boils the acid upon the remaining mercury, there is no further disengagement of

it does not feem to differ chemically; nor does it contain any fensible portion of water of crystallization. The same may be faid of corrofive fublimate.

It now remains to speak of the real hyperoxigenized muriate Hyper-ox. mur. of mercury. I passed a current of oxigenized muriatic acid of mercury: obtained by passes gas through some water, in which there was red oxide of mer- fing ox. m. gas cury \*. After a short time, the oxide became of a very dark through water containing red brown colour; and a folution appeared to have taken place. oxide of mer-The current was continued for fome time; and, when I thought cury. that a fufficient quantity of the oxide had been dissolved, I stopped the operation. The liquor was evaporated to dryness; and the falt was thus obtained. There evidently was in the mass a great proportion of corrosive sublimate, as might be expected, from what I had observed to take place in the forma- It is more soluble tion of the other falts of this acid; but, by carefully separating than cor. sub. and is separable the last formed crystals, I could pick out some hyperoxigenized by crystallizamuriate of mercury. I then crystallized it over again; and, tion. in this manner, I obtained it nearly pure. This falt is more foluble than corrofive fublimate: about four parts of water retain it in folution. The shape of its crystals, I cannot well

gas; yet more mercury is dissolved. The nitrate of mercury, therefore, rather contains an oxide less oxidized after ebullition than before it. The true difference is in the sub-nitrate of mercury, precipitated, as I before said, by the water in which the muriate of soda was diffolved. And the orange-coloured powder, which remains after an attempt to fublime Scheele's calomel, is to be attributed to the fame cause. To prepare calomel in the humid way, uniform as Preparation of to itself, and in all respects similar-to that prepared in the dry way, calomel in the it is necessary, either to use the nitric solution before it has boiled, humid way. or to pour some muriatic acid into the solution of muriate of soda, previously to mixing it with the boiled folution of nitrate of mercury. In the first case, no precaution is necessary; and, in the latter, the oxide of mercury, which the nitrate of mercury has, by boiling, taken up in excess, finds an acid which is ready to faturate it. All the mercurial oxide being thus converted into calomel, none of that sub-nitrate of mercury can be present.

The objections made by a medical gentleman against Scheele's calomel, when this paper was read before the Royal Society, led me to reconsider the subject, and to undertake the investigation detailed in this note.

\* I used either of the red oxides of mercury indiscriminately.

determine. When fulphuric, or even weaker acids, are poured upon it, it gives out the usual smell of hyperoxigenized muriatic acid; and the liquor becomes of an orange colour. This is a sufficient proof, that corrosive sublimate is not a hyperoxigenized muriate of mercury.

Dark brown oxide of mercury not taken up, supposed to be different from the red oxide.

I have just mentioned that, in the formation of this falt, the oxide of mercury, which was not diffolved by the acid, became of a very dark brown colour. I procured a portion of this oxide, which feemed different from the red oxide. It however retained the form, and the crystalline appearance, of the latter. It was foluble in nitric acid, without difengagement of gas; and was precipitated from it, in a yellow oxide, by all the alkalis, except ammonia. It formed corrofive fublimate with muriatic acid; and the precipitate by the alkalis, was the same as that from corrofive fublimate, made with the red oxide. Yet I am inclined to think, that the dark brown oxide differs in some effential point from the red; but I have not yet made fufficient experiments to prove this opinion. At all events, the present object being to examine the mercurial oxides only as combined with muriatic acid, it would be foreign to the purpose, to enter into too minute an investigation of the other thates of the metal. This, and some other objects hinted at in this Paper, must be referved for future inquiry.

of filver

Hyper-ox. mur. In treating the earthy hyperoxigenized muriates with phofphate of filver, as I mentioned before, I observed that the liquor fometimes contained in folution oxide of filver; which, upon examination, I found to be combined with hyperoxigenized muriatic acid. As the falt which is thus formed is different, in every respect, from simple muriate of silver, it may be of some importance to consider it with attention. In the first place, it will afford the most convincing proof of the difference between muriatic and hyperoxigenized muriatic acid; and, in the next place, it particularly deferves to be remarked, for possessing, in the most eminent degree, one of the great chafoluble in 2 parts racteristic features of the genus to which it belongs. Hyperoxigenized muriate of filver is foluble in about two parts of

warm water, and crystall. by cold:

flightly fol. in alceho!.

warm water; but, by cooling, it crystallizes in the shape of fmall rhomboids, opaque and dull, like nitrate of lead or of It is fomewhat foluble in alcohol. Muriatic acid decomposes it; as does nitric, and even acetous acid: but the refult of this decomposition is not, as might be expected, nitrate or acetite of filver. At the moment that the acid is expelled Acids decompose from hyperoxigenized muriate of filver, a re-action takes place it, but leave muriate inftead of among its elements: oxigen is disengaged; and the muriatic seizing the filacid remains in combination with the oxide of filver. If this veracid the compared with the manner in which nitric and acetous acids act upon hyperoxigenized muriate of potash, it will give a strong proof of the proportionate affinities of all these acids for oxide of filver, in comparison with that which they exercise towards the alkali.

Hyperoxigenized muriate of filver, when exposed to a very Hyp. ox. mi of moderate heat, begins by melting, and then gives out a configen by heat, and derable quantity of oxigen gas, with effervelcence; and mu-becomes muririate of filver remains behind. These phenomena however ate. differ much, according to the degree of heat applied. When With sulphur it hyperoxigenized muriate of filver is mixed with about half its explodes; weight of fulphur, it detonates in the most violent manner; -and does not, like hyperoxigenized muriate of potath, require the addition of charcoal, to possess a very great force of explofion. The flightest pressure is sufficient to cause this mixture by the slightest to detonate; and I think I shall be within bounds, when I pressure, -flate, that half a grain of hyperoxigenized muriate of filver, with a quarter of a grain of fulphur, explodes with a violence at least ten times at least equal to five grains of hyperoxigenized muriate of pot- as frongly as ash, with the due quantities of sulphur and charcoal. The of potash. -flash is white and vivid, and is accompanied by a sharp and quick noise, like the fulminating filver so ably described by Mr. Howard; and the filver is reduced to the metallic state. and vaporized.

I think it right to add a few remarks upon what I have Order of attractermed the proportionate affinities of acids and of bases, one tion of acids for oxides is frequently other. It is a law, not indeed universally, but frequently very differently observed, and very well worthy of consideration, that ferent from that the acids are attracted by metallic oxides, in a very different order from that in which they are disposed to unite to alkaline and earthy bases.

Nitric acid, which holds so high a place in the order of Acids commonly affinities for alkalis, is expelled from metallic oxides by most attract oxides more the less acids. Phosphoric, fluoric, all the vegetable acids, except two they act on the or three, and the animal acids, attract the latter bases more metals: affrongly. Nay, we shall find, upon an attentive examination,

Hyp. ox. m. acid follows this rule.

that acids commonly attract metallic oxides, in the inverse ratio of their action upon metals, or, in other words, in proportion to their own affinity of composition. Thus, the phofphoric and fluoric acids fometimes rank before the fulphuric; and the nitric, as I before faid, is generally very low. Hyperoxigenized muriatic acid feems to follow the fame rule; and takes its place, in the order of affinities for metallic oxides, after many of those acids which it can expel from earths and alkalis.

The metallic butters have their metal and not their acid extraordinarily exigenized.

The other hyperoxigenized muriates, I have not yet fufficiently examined. I shall, however, mention at prefent, that I have afcertained the muriatic falts, formerly known by the strange name of butters of the metals, to be muriates, and not hyperoxigenized muriates; and the extraordinary proportion of oxigen, to be combined, not in the acid, but in the metallic oxide.

In the course of different experiments, I have known hyperoxigenized muriatic acid to be formed in two cases, where I could not have expected it.

Precipitation of from mur. acid by potash, gave oxigen to the latter, and form-

In the analysis of some menachanite from Botany Bay, given oxide of titanium to me last year by the President of the Royal Society, I obferved, that while the oxide of titanium was precipitated from the muriatic acid in which it was dissolved, the excess of oxiedex. m. potash, gen in the oxide passed over to the muriatic acid and the potash, already in the liquor, and that hyperoxigenized muriate of potath was formed. I have attempted the fame experiment with black oxide of manganefe, but could not fucceed.

Whennitro-mutilled from platima, the metal is oxided, and ox. and byper-ox. m. acid are formed.

There is, however, a still more extraordinary formation of riatic acid is dif- this acid, in the distillation of nitro-muriatic acid upon platina. Oxigen is absorbed by the metal; yet, not only oxigenized, but also hyperoxigenized muriatic acid is formed. I have repeated the experiment feveral times; and am well convinced of the fact, however contrary to theory it may appear. I have tried the action of oxigenized muriatic acid, upon nitric acid, in the hopes of forming hyperoxigenized muriatic acid; but there was no action to this effect among their elements.

This fact imperfeetly feen beforc.

The fact of the production of a peculiar gas, by the diftillation of nitro-muriatic acid upon platina, has been observed by Mr. Mr. Davy, in his Researches \*. But, as hyperoxigenized muriatic acid was not known at that time, he could not fay the real nature of that gas. Had Mr. Davy carried his ingenious experiments a little farther, we should have been much earlier acquainted with the last degree of oxigenizement of muriatic acid.

Mr. Berthollet terminates his Paper upon hyperoxigenized Berthollet confimuriate of potash, by saying, that he will consider muriatic acid as a radical, acid as the radical; oxigenized muriatic acid, as correspond- &c. ing with fulphureous and nitrous acid; and the acid which he conjectured to exist in this falt, as corresponding with sulphuric and nitric acid. I fliall now conclude, by flating the arguments in favour of each denomination, and the analogies upon which they are founded.

Muriatic acid is for us a fimple body; but it has acid pro- The arguments perties of the strongest kind; therefore, from analogy, we stated on this suppose it to contain oxigen. But may not this be too hasty a conclusion? Are we not very doubtful concerning the existence of oxigen in prussic acid? And are we not, on the contrary, certain that fulphurated hidrogen, which possesses many of the characteristics of acids, does not contain any? Of the oxigenizement of fluoric and boracic acids, we have no proof: but then we cannot affirm that any one of these acids exists in three states of combination with oxigen; and the muriatic is the only radical of which we admit this fact. We must not, however, pretend to limit the number or degrees of combinations between combustible bodies and oxigen; but we can fpeak with certainty only of those things which are proved. Besides its acid properties, this substance has others, common Mur. acid, like to oxigenizable bodies. With 16 of oxigen, it forms an acid, the properties of which, in many of its properties, is to its radical what the oxigenizable bofulphureous is to fulphur. Like the fulphureous, it is volatile; dies, and may not contain oxhas little attraction for falifiable bases; destroys vegetable igen. blues; and is capable of further oxigenizement. With 65 of oxigen, it becomes more fixed, like fulphuric acid; has a stronger affinity for salisiable bases; and acquires more truly acid properties. Upon these considerations, I submit to

Dr. Priestley, also, mentions a peculiar gas, produced by distilling a folution of gold in aqua regia.

the chemical world, whether, in the prefent state of our knowledge, it be not more philosophical to say,

Nomenclature of Muriatic radical, or this hypothesis. fome fingle word of the same import,

of the same import, Muriatous acid, Muriatic acid, Muriatic acid

Oxigenized muriatic acid; Hyperoxigenized muriatic acid,

It is not proved that every acid contains oxigen.

I am fully aware that, at first fight, this may appear extraordinary; and the more so, as we have no positive facts that prove muriatic acid to be a simple body. All we can, therefore, consider fairly, is, in favour of which appellation does the sum of analogies seem to preponderate. And, to give the canse a candid investigation, we should begin by considering, whether the presence of oxigen in all bodies that have acid properties, has been rigidly demonstrated; and not determine by this law of the French chemistry; till we are well convinced it has not been too generally assumed.

Nomenclature should be governed by the faces; and not the contrary. If a nomenclature be not subservient to the uses of science, and does not keep pace with its progress, the relation between substances and their names will become so relaxed, that consussion will be brought about, by the very means we take to avoid it; and if, while we continue to extend our acquaintance with chemical bodies, nomenclature remains confined within its former limits, the bonds that unite these two parts of the science must inevitably be broken.

#### HI.

On the present State of the British Museum. By a Correspondent\*.

The British Mufeum stated to be of no public utility.

cial manuer;

IT will probably be confidered a prefumption, even bordering on impertinence, to remark, that no establishment is of less utility to the public, and none a greater reproach to the nation, than the British Museum in its present state; but as the remark is made with the purest motive, it is hoped its apparent petulance will be excused.

That it is not That this magnificent and valuable collection, bought by the open to the public for their own use, and entertained at their own explicinany benefi-

\* Whose name accompanied the paper.-W. N.

pence, should be rigorously shut up from the public themselves, is so extraordinary and disgraceful a fact, that I am wholly at a loss to account for it, particularly when I resect that the sact occurs in a nation who prides itself, above all things, on the free enjoyment of its liberty and its property.

It may be objected to me, that the Museum is not literally because the times that up from the public, and that, on the contrary, it is in the of inspection are power of any individual to procure a sight of it. True; it is in nient to trudents; the power of any one to procure a cursory view of it, after the formality of inscribing his name on the register of admission, and waiting until the keepers think proper to appoint him a precise day and hour, when it is very likely his other engagements may deprive him of the possibility of attending. But, and the actual suppose the appointment procured, and the time convenient, process is that of a multitude hurwhat gratification or instruction can be derived from prowling ried through a through a suite of apartments, at the heels of a keeper, who suite of rooms. Is anxious to dispatch you as soon as possible, and in company of a dozen people, each of whom is eager to catch a transient glance of every object within his reach, and each exclaiming at the "wondrous surprising" things which surround him.

The library, the most valuable and most extensively useful The library faid part of the British Museum, is precisely that part which is most only at the plean completely secluded from the public. To visit this it is indisfure of the lipensably necessary to be introduced to the librarian, and to obstrain. tain his permission to consult the collection. How different is Not so on the it on the Continent! there, scarce a town of any note but has Continent. its public library, to which free access may be had at all times, by every individual, whether stranger or native. How humilinesselection ating the comparison! and how singular that London, where the sciences and the arts are cultivated with greater success probably than in any other spot of the globe, should be destitute of a single public library.

If I might presume to suggest a plan for rendering the British Suggestion; that Museum a real benefit to the public, I would propose that the the library and library, and the collection of objects relative to natural history, be separated. Exc. should be formed into two separate establishments, to each of which the public might have the freest access during as

great a portion of the day as possible.

The library might be open all the year round, from eight in Regulations for the morning until four in the afternoon, and again from fix the library.

until ten at night. It should be furnished with tables, pens,

and

and ink: the floor should be matted, and the room properly heated during the winter season. The books should be delivered by the affistant librarians to those who might require them; but on no account should any book be permitted to be taken out of the library. A regular register might be kept of the books delivered out each day, with the names of those who required them; and the delivery might be cancelled as they were returned to the librarian. It might perhaps be proper, on account of the value of the manuscripts, to require the express permission of the principal librarian for their perusal.

Concerning the expences and other objects.

A proper fund should be provided for the necessary expence of the library, which might be under the direction of a committee: fuch fums as were over and above the expence of the maintenance of the library, might be applied to the purchase of fuch books as the committee should see expedient. But to contract, as far as possible, the expence of supplying the library with new publications, the donations of public spirited individuals should be solicited, and the names of the donors registered in a conspicuous manner on each book they presented. The committee might be empowered to fell, by public auction, from time to time, fuch duplicates as might be received, &c. and be thought unnecessary to retain in the library. The accounts of the library might be adjusted every quarter; but they should conftantly remain open for the inspection of the public. A regular catalogue of the library should be printed in octavo; and supplements might be added from time to time, as the new acquifitions should render it necessary: one or more of these catalogues should be constantly kept in the library, for the purpose of general reference. A binder and proper affistants should be attached to the library, through whose hands each volume should pass at least once every year; and it might not be beneath the attention of the committee, to hold out premiums for any improvements which should be made in the binding of books, fo as more effectually to preferve them from the attack of the worm, and from gradual decay.

The collection of natural history.
Direction:

The collection of objects relative to natural history, &c. should be under the superintendance of a separate committee; and this establishment, if properly conducted, could hardly sail to become one of the finest and most instructive repositories in Europe. No nation is so well situated for establishing a national Museum as our own: our extensive commerce, and

the public spirit of individuals, would shortly supply us with the choicest specimens from every quarter of the globe; and at this very moment thousands of valuable specimens are rotting in obscurity, for want of a national Museum to receive them.

The arrangement of this establishment requires considerable Arrangement of knowledge and ability; it would therefore be the highest prefumption in me to attempt to dictate on the subject. I should Five classes. Conceive, however, that the whole collection might be advanables; minerals; tageously separated into sive classes. The first might comprecuriosities; obhend the animal, the second the vegetable, and the third the jects of arts mineral kingdom; the fourth might comprehend objects of pure curiosity, and such as were illustrative of the state of society, of arts, and of manusactures of various nations; the sight comprise such objects as tended to the improvement of, or were simply illustrative of the arts, sciences, and manusactures of our own country.

In the arrangement of the objects of natural history, I would Natural history. fuggest that the most extensive scale should be assumed, and that Subjects. spaces should be left for the insertion of such objects as were wanting, and were likely to be procured: the places of these objects should be supplied by accurate drawings or models, if possible. To each object might be attached its proper name, both in Latin and in English; and a descriptive catalogue of Catalogue.

the Museum should be printed, which might at the same time

ferve as an introduction to the study of each class.

The donations of public spirited individuals should be soli-Donations. cited for this establishment, as for the library; and the names of the donors should be regularly inserted in the public prints. The committee should also be empowered to dispose of such duplicate specimens as might be sent, by public auction.

A laboratory for the preparation of objects of natural history Laboratory. should be annexed to the establishment; and the committee might be directed to offer premiums for improvements in the art of preparing and preferving these objects. The committee might also be enjoined to publish concise directions for travellers, for collecting, preserving, and transporting the different objects they might meet with; copies of this work should be given to all who might request them.

I am inclined to think that if this establishment was conducted Consequences of on an extensive and a liberal scale, it would shortly comprise such an extensive and liberal most

establishment.

conduct of this most of the private collections in the kingdom; for it is prefumed that there are few lovers of natural history who would not prefer the honourable applause of completing the national Museum, to the selfish pleasure of possessing an impersect and comparatively useless collection of their own, especially as they would enjoy the credit of the collection equally as if it remained in their own private possession.

Extension of the Muleum.

By degrees as the collection enlarged, and as its utility was more fenfibly felt, it is to be prefumed that the nation might be induced to purchase ground, and to erect appropriate and extensive buildings to contain it: in short it would seem, were this plan to be adopted, that we might indulge ourselves with the prospect of speedily possessing one of the finest, the most interesting, and most instructive establishments in the universe.

J. G.

## ANNOTATION. W. N.

Though the importance and value of public establishments

have been strongly felt and admitted in all ages, yet there has

not, as far as I know, been any direct investigation of the means

Public eftablishments are of great importance.

They have vigour at firft;

but foon decay.

Is this degene-

Endowments, Sec.

of giving them permanence and effect. A public library, college, inftitution, or fociety, may be radically defective in its conflitution, and, nevertheless, flourish for a time by virtue of the ability and active exertions of its founders; and it is certain, that every constitution whatever will afford only a perishable establishment, unless it contain some inducement by which fuch exertions shall be invited and continued in its favour. appears to be extremely difficult, and perhaps in strictness impossible, to produce motives of this kind by mere regulation; and truftees, managers, inspectors, or governors, being bound by the letter of the flatutes, can feldom be expected to make advances in support of the spirit of the foundation. racy inevitable? unavoidable confequence, that schools, universities, and sqcieties, shall degenerate into mere instruments for conferring degrees or titles? Are all establishments perishable unless endowed? Is it from the operation of inevitable causes that various endowments are productive of little or no effect, and that most unendowed establishments disappear in a few years?

Instances. Gre-Among the former we may mention the Gresham foundations sham College and others.

and various public libraries in this town, with feveral colleges, schools, and societies, which it would answer no good purpose to name without more ample comment; and among the latter the colleges of Warrington and Hackney, begun with great spirit, but in a few years abandoned. Or may we not rather The history of hope, that an investigation of constitutional and historical facts, may probably respecting all the public means of instruction in this and the flew how to neighbouring countries, would affift us in the means of giving preferve their full energy to these important establishments? How are the Correspondents collections and libraries on the Continent regulated as to supply invited to state of materials, preservation from injury or embezzlement, and continental liready access by the student? What may be the state of the braries, the Brimuseums and libraries in London, and our universities, in these &c. in order to and other respects? Can the collective power of any body of make progress. men be eminently applied to forward literature and the sciences; and if fo, why has it generally failed? &c. Thefe and many other interesting subjects of inquiry, might be cleared up by Correspondents of Journals, who may not have leifure for the direct and ample discussion of a subject of such magnitude and extent.

### IV.

Of the Properties of the Earth Yttria, compared with those of Glucine; of Fossils, in which the first of these Earths is contained; and of the Discovery of a new Substance of a metallic Nature (Tantalium). By A. G. EKEBERG\*.

THE first part of this Memoir contains the particulars of the First part of the experiments made by M. Ekeberg, to confirm the difference Memoir. Existing between glucine and the earth last discovered in gadolinite, and named yttria or gadoline; but as the particular nature of each of these earths is well known to chemists, it would be superstuous to say any thing in this place on the subject.

<sup>\*</sup> Kongl. fwænsk. Vetenskups, academiens nya Handlingar; fser eer 1802, Qwartal I. Pag. 68.

This abridged Memoir is from the Journal de Van Mons, but I take it from the Journal de Physique, LV. 281.

As there is novelty to recommend the fecond part of this Memoir, we shall communicate it to our readers in a faithful translation.

Mineral substance newly investigated, containing a metal hitherto unknown.

The author remarks, that although the mineral substance which he has discovered contains yttria, it cannot be classed in a fystem of mineralogy as a species of that earth, on account of the more abundant presence of another equally remarkable fubstance, which must be carried to the numerous class of me-He found this substance in two fossils obtained from different fituations: in one place it was united with iron and manganese, and in the other with iron and gadoline.

Infoluble in all the acids; foluble by fusion in alkali;

precipitable by acid, but not foluble ;

the precipitate ignition;

foluble by heat with alkaline phosphate or borate; but gives no colour. It is reducible to a metallic button by charcoal;

and is then oxivable, but not foluble by acids.

It appears to be a new metal;

infoluble, and not, like it, reducible; nor tungsten, because it is nei-

This new metallic substance is characterized by its infolubility in all the acids. The only re-agent that has any action on it, is fixed caustic alkali. When exposed to the fire with that alkali, the mass being afterwards lixiviated, it partly diffolves in water, and may be precipitated from this folution by means of an acid, but the precipitate is not in the least attacked, however great the quantity of acid employed. When feparated by filtration and dried, it has the form of an ex-

tremely white powder, which does not change colour even at white, even after a red heat. The remainder of the mass being treated with acids, affords the fame powder. Its specific gravity, after ignition, is 6,500. It is fufible with the blow-pipe, by the addition of alkaline phosphate and borate of foda, but it does

not communicate any colour to the flux.

When exposed to a strong heat in a crucible, without any mixture except pounded charcoal, it is reduced to a moderately hard button, having a metallic brilliancy at its furface, but its fracture is dull and blackish. The acids have no otheraction on this kind of regulus, than that of causing it to return to the state of white oxide. The circumstances of the reduction, as well as the specific gravity of this singular substance. feem to place it among the metals, and I have sufficient reason to be perfuaded that it is not the same as any of those hitherto known. The fubstances with which it is likely to be confounded are the oxides of tin, of tungsten, and of titanium. which are foluble in the caustic alkalis, and in all circumnot tin, because stances resist the action of acids. But the oxide of tin is easy to be diffolved and reduced; tungften is eafily detected by its folubility in ammonia, and by the blue colour it communicates 7'5 7 .415.11.2VI .58.5070

to phosphate of soda; the oxide of titanium gives to borax the ther soluble in colour of hyacinth, and becomes foluble in acids when fused ammonia, or gives a blue cowith carbonate of potash \*. lour to phof.

Before the author describes the chemical analysis he has un-foda; dertaken of these two substances, which he regards as ores of because it does the new metal, he has chosen to describe their external cha-not colour bo-rax, nor become racters. In order to avoid circumlocution, he has ventured to foluble in acids give them a generic name. By availing himself of the usage when sufed with carb. potath. which admits of mythological denominations, and in order to External characexpress the want of power of this new metal to become fatu-ters. rated with the acids into which it is plunged, he has given it The new metal the name of tantalium; for the ore composed of tantalium, of denominated tantalium. iron, and of manganese, he proposes the name of tantalite; and its ores tantalite for that containing yttria, the appellation of yttrotantalite, and yttrotantawhich cannot be thought more heavy than that of fiderotitanium, already adopted.

He received the specimen of tantalite from M. Geyer, Di-Locality and rector of Mines, who has affured him that this substance has gangue, &c. of been known fince the year 1746, and has been regarded as a problematical variety of tin garnet (zinngraupen). It is found near the farm Brokaern, in the parish of Rimito, in the government of Abo, in Finland, in a large mountain, on the banks of the Baltic. The gangue is composed of white quartz and mica, with veins of red feld spar in large plates; substances which compose the sides of the gangue. The titanite is diffeminated in the form of garnets.

The specimens he has seen were in detached crystals, of Size and figure the fize of a nut, the most regular among which seemed to of crystals. approach the octahedral form: they contained particles of feld fpar and mica.

Their furface is even, polifhed, and blackish.

Fracture compact, and of a metallic brilliancy; colour of Fracture. the fracture not alike all through; it varies between a greyish blue and the black of iron.

When pulverized, it is of a blackish grey approaching to Powder; grey brownish. brown.

\* The author nevertheless entertained some suspicions of the identity of this new substance with the latter metal, and nothing but a comparison with the titaneous iron of Norway, which he decomposed with this view, convinced him that this identity does not exist.

Gives sparks.

Sp. gravity. Locality and gangue, &c. of yttrotantalite.

It is fufficiently hard to give sparks with the steel. He did not remark whether it is attracted by the magnet.

Its specific gravity is 7,953.

The yttrotantalite is found in the fame place and in the fame gangue as gadolinite. Klaproth fays, that the latter is found incrusted in a mass of granite; but though the constituent parts of granite are occasionally found with it, it is not the less infifted that feldspar is its true gangue, as is clearly seen in the instances of the great quarry at Ytterby. The mica and quartz found in it form diffinct parts, and do not enter into combination with the feldspar; (he speaks only of the masses of the principal rock, without disputing the possibility of particles of the three substances being found mixed); but in general it is a rock of feldspar intersected by large veins of mica in a direction almost perpendicular; and the gadolinite and yttrotantalite are to be fought near these veins. The first is commonly found attached to one fide of a vein of filvery mica, and the rest of its mass enveloped with feldspar. The fecond is never found immediately adherent to mica. The maffes or kidnies which it forms, are enveloped with a thin crust of feldspar. feparated from the general mass by slight layers of a greyish black mica. The gangues charged with these kidnies are seldom insulated: They are found in clusters, separated from the principal rock and from one another by fimilar coatings of mica. These substances are generally thus disposed by nature, and it is very feldom that they are found diffeminated in grains in the rock of feldspar.

Size and fracture of the maffes.

The largest masses he has found of the ore of yttrotantalium were not of the fize of a nut. Its fracture is granulated, of the black colour of iron, with a metallic brilliancy.

Hardness.

Its hardness is not confiderable; it may be feraped with a knife, though difficultly.

Powder.

When in powder it is of a greyish colour.

Not magnetical.

The magnet does not attract it.

Sp. gr.

Its specific gravity was found to be 5,130; but as no specimen was found totally free from feldfpar, it was fufpected that its real weight is more confiderable.

On gadolinite.

The abridger concludes his extract by giving an outline of 'that part of M. Ekeberg's Memoir, which treats of the analysis of gadolinite and the comparison of yttria with glucine, which feveral writers have affirmed to be of the same nature.

Gadolinite

Gadolinite when pure is hard enough to give fire with fteel. It is hard; It is found crystallized in an imperfect manner like fome crystallized; species of garnets.

It contains glucine.

contains glueine,

Besides the distinctions between gadoline and glucine esta-Gadoline and blished by Klaproth and Vauquelin, these earths differ in the glucine differ following peculiar properties.

The specific gravity of gadoline differs considerably from in spec. gravity. that of glucine, which is only 2,967, whilst that of gadoline is 4,842. This last earth is the heaviest of all the known earthy substances, since it is 0,842 heavier than barites, which only weighs 4,000.

It is foluble in the alkaline carbonates.

It is absolutely insoluble in the caustic alkalis.

It is not precipitated by the fuccinates as glucine is.

Gadoline is fofuble in alkaline carbonates. not in pure alk. nor (like glucine) precip. by fuccinates.

#### V.

On the Probability that the Eruptions of Lunar Volcanos may reach the Earth. By J. B.

## To Mr. NICHOLSON.

SIR,

Cambridge, Nov. 6, 1802.

HE fact of stony and metalline substances having at differ- Stones fallen on ent times fallen on the earth, feems to be confirmed by fuch the earth. strong evidence, that a doubt can scarcely remain as to its reality; permit me through the medium of your Journal, to offer to your readers an attempt to account for this phenomenon. The existence of lunar volcanos, at least equal in mag-Lunar volcanos. nitude to those of the earth, has been proved by ocular demonstration. The velocity with which a body must be pro-Velocity of a jected from the moon's furface fo as to be carried within the projectile to be thrown from the sphere of the earth's attraction, is about 12,000 feet per se-moon. cond; now when we confider the height to which volcanic fubstances are projected during an eruption, and how very confiderable a part of swift motion must necessarily have been destroyed by the resistance of the atmosphere, I think it by no means improbable, that the force exerted may be confiderably greater than would be necessary to carry a body 12,000 feet in

The lunar atmosphere relists little.

may be thrown to the earth.

vaeuo. The atmosphere of the moon cannot offer fo great tefistance to the motion of bodies as that of the earth, as it is certainly much rarer than ours, and probably its height is not Whence bodies much greater than that of the lunar mountains. From all these circumstances taken together, I think it by no means improbable that the volcanoes of the moon may occasionally project bodies with fusicient force to carry them to the earth.

I remain, SIR.

your's, was a second of the

I. B.

## OBSERVATIONS.-W. N.

Remarks on the above letter;

Against the conjecture of J. B. it seems to be a principal difficulty that the stones fallen on the earth are thought to have arrived in a flate of ignition, which they must have preferved through their immense passage from the moon. If the meteors or ignited globes be allowed to be stones of the kind in queftion, their extreme velocity in fome inflances may be thought to favor his hypothesis, But this subject, like every other in which the data are few, affords wide range for speculation. When we reason from our want of knowledge, there is scarcely any branch of science more favorable to the process than chemistry. We are ignorant of the component parts of the earths. the metals, oxigen, azote, hidrogen, &c. and of the very existence of light, caloric, electricity, and magnetism, as independent bodies. Some, if not all of the former may be, and most probably are, compounded. We see very sudden or inftantaneous separations of water, light and electricity from the atmosphere. May not iron and nickel be among its ingredients, and may they not by causes equally unknown, be no less rapidly separated, with the emission of light and heat? Even if these metals exist in the atmosphere disseminated, sufpended, or diffolved like aqueous vapors, (and we know they emit effluvia perceptible to the fmell) their quantity may elude all chemical tests from its minuteness, and still be sufficient to answer the phenomena. A cubic foot of air does not weigh 100 grains, and if one thousandth part of this were metal, the quantity contained in the air to the height of 500 feet over a fingle field of ten acres would be about a ton and a half, of which a small part being precipitated, would afford a prodigious shower of stones.

with other very loofe conjectuies.

VI. Enquiries

#### VI.

of spides yes and Employed the

Enquiries concerning the Dilatation of the Gases and Vapors.

Read to the National Institute of France. By CIT. GAY

LUSSAC.

# (Concluded from Page 216.)

THE following is the table of refults of Citizen Guyton and Duvernois, in which they have placed between brackets those in which they place little confidence.

	From o° to 20°	From 20° to 40°	From 40° to 60°	From 60° to 80°	From o° to 80°
Common air	1	1	1	(1)	1
dilates to	12,67	5,61	2,49	$\left(\frac{1}{3,57}\right)$	1,067
Vital air	1	1	11	$\left(3 + \frac{1}{1,73}\right)$	4+ 1
	22,12	4,92	1,53	(37 1,73)	2,09
Azote gas	1	1	1	$5+\frac{1}{57,2}$	5+
Tizoto gus	29,41	5,41	1,82	57,2	1,062
Hidrogen	1	1	$\left(\begin{array}{c} 1 \end{array}\right)$	$\left(\frac{1}{58,82}\right)$	1
gas	11,91	6,92	6,85	58,82	2,55
Nitrous gas	1	1	1	$\left(\frac{1}{-1}\right)$	1
	15,33	9,00	3,739	6,88	1,65
Carbonic	1	1	4-1 ×	$\left(\frac{1}{3,69}\right)$	1+
acid gas	9,049	5,099	2,31	3,69	$1+\frac{1}{106,3}$
Ammonia-	1	1	11	$\left(3 + \frac{1}{4,69}\right)$	5+-1
cal gas	3,58	1,75	1,35	4,69	1,248

Before I proceed further, I ought to premife, that though I had frequently found that oxigen, azote, hidrogen, carbonic acid gas, and atmospheric air, dilate equally from 0° to 80°, Cit. Charles had remarked the same properties in these gases fifteen years ago; but having never published his results, it was by mere chance that I became acquainted with them. He likewise endeavoured to determine the dilatation of gases Vol. III.—December, 1802.

which are foluble in water, and he found in each a particula dilatation differing from those of the other gases. In this respect my experiments differ much from his.

Apparatus of Cit. Charles. A barometer with a large vacuum, and a refervoir below for the gas to prefs on the mercury.

This was inaccurate, because the rise of the mercury gave room for expansion.

The apparatus of Cit. Charles confisted of a barometer whole chamber or upper space was of great extent, The gas on which the experiment was to be performed was enclosed in the refervoir, or lower space of the barometer, at the temperature 0°, and under the pressure of 28 inches of mercury. When the barometer was plunged in boiling water, the mercury rose in the tube, and the excess of the whole column above 28 inches indicates the elasticity acquired by the gas; but Cit. Charles having had the politeness to shew me his apparatus, I saw that the tube of the barometer was too large in proportion to the capacity of the refervoir; fo that the elevation of the mercury above 28 inches did not indicate all the fpring the gas had acquired, because in that case its volume in the refervoir ought to have remained conftant. It appears then to me, that the real dilatation of the gases cannot be decided from his experiments.

## ART. III. Description of Apparatus.

Apparatus for the gafes. A globe with an iron cock; filled as ufual, and immerfed in water to be heated. The gas is fuffered to expand and pafs out, and the expansion meafured.

A receiver B (Fig. 1. Pl. IX.) has a cock of iron R, to which may be adjusted a bended tube I, D, (Fig. 2). The key of the cock carries a lever L L bored at its two extremities in order to receive two cords, by means of which the cock may be opened and flut under the water. In order to introduce the gas into the receiver I use a bell glass M, (Fig. 1.) to which are adapted a cock and bended tube T, plunged in a vessel Q, S. When water is poured into the vessel, and the cock opened, the gas compressed in the bell escapes through the tube and fills the receiver B, reverfed on a bath P, O, of mercury. The receiver B being full I shut the cock: I adjust the tube I, D, (Fig. 2.) to it, and I fix it in a cylindrical cage of iron E, F, G, H, which I then carry to a copper veffel A, D, filled with water. In order that there might be no communication between the exterior air and the gas inclosed in the receiver, when the cock is opened I cause the extremities of the tube I, D, of one or two millimetres (thirtieths of an inch) to be plunged in a fmall bath K, X, of mercury. This done I heat the bath, and from 10° to 10° for example, I open the cock and shut it immediately. The gas dilated by

the heat coming brifkly out of the receiver, quickly drove off the atmospheric air that filled the tube, and after the 40° the cock may be fafely left open during the remainder of the operation: I prefer however alternately opening and shutting it, because I find that the gas of the receiver acquires the temperature of the bath better. After 15 or 20 minutes of ebullition, a time sufficient for the whole to acquire the same temperature, I disengage the extremity of the tube I, D, from the mercury in order to re-establish the equilibrium of pressure between the exterior air and the gas of the receiver, and I then close the cock. After having cooled the bath with ice or water I take away the apparatus, I disengage the receiver from which I take the tube I, D, and the lever L, L, and I plunge it altogether in a bath of a known temperature, where I leave it a sufficient time to acquire its temperature.

When the cock is opened a volume of water enters the re-The relative ceiver, which when reduced to a level is precifely equal to bulks of the gas at different tem-that of the gas expelled by the heat. When the cock is that peratures deter-I withdraw the receiver; I dry its exterior furface with care, mined by and I weigh it in this state. I then weigh it first full of water, weighing. and then empty, and note the refults of each weight. By this knowledge I obtain the capacity of the receiver by fubtracting its weight when empty, from its weight when filled with water; and also the volume of the water which represents the volume of air expelled by the heat from the receiver, by fubtracting again the weight of the receiver when empty, from its weight when containing this water. Hence it is very eafy to determine the proportion of the first volume of the air to that of the dilated.

This method has the advantage of being very exact; for as This method is the volumes are determined from the weights, any error that very exact. is committed in this determination must be very trifling, even if the scales employed should be but moderately sensible.

The apparatus I have just described is simple enough in it- Another appafelf; but as it contains cement, and a cock which ought to be ratus nearly as made of iron on account of the mercury, it is rather difficult easily conto execute. It will not therefore be amiss to describe another structed. apparatus which I employ, and which on account of its great fimplicity and easy execution, very nearly combines all the advantages of the former.

the dilated as

A body with a life It is a simple globe or body, and the neck of which must be long neck; gra- at least one decimetre (about four inches) long. After having -filled it with the gas by the method already described, I

cury;

plunged in mer- plunge its neck about two centimetres (three quarters of an inch) in mercury contained in a common glass vessel, and I fecure it by an iron frame like the preceding apparatus. If I

communicate with the outer

were to plunge it in this state into a bath of heated water, the dilated gas in order to escape would have to overcome not only the pressure of the mercury in the glass, but likewise that A small tube to of the water of the bath. To remedy this inconvenience, I introduce into the neck of the receiver the extremity of a very air when requi- fine bended tube, taking the precaution to keep the upper aperture closed till the tube is plunged beneath the mercury.

.To support the tube I tie a string about its middle, to the end of which I fuspend a weight, and I pass it over a support in fuch a manner, that the cord by its action shall have a tendency to draw the tube upwards. The apparatus being thus disposed I plunge it in a glass vessel, the water in which must be on a level with that in the bath: I open the extremity of the tube for an inftant, that the equilibrium of pressure with the external air may be restored, and I then shut it. As there

Observation of the mercury in the neck.

is a scale marked on the neck of the receiver whose divisions are extremely minute, I find exactly the level or station of the mercury in the neck, and I note it, because it is at this flation that the capacity of the receiver terminates. The lower extremity of the tube must rise clearly above the surface of the mercury, which otherwise would flow into the tube and resist

the passage of the dilated gas. After all these operations, which are longer in the description than the execution, I carry When the apparatus to a bath of hot water, and I open the upper

ratus is plunged in boiling water, extremity of the tube, after having plunged it into a small bath the expanded gas of mercury, as in the preceding arrangement. When the reescapes by the ceiver is at the temperature of boiling water I withdraw the tube which is then withdrawn. tube, the extremity of which must be previously disengaged Water is fuffered from the mercury, and I cool the bath. The mercury then to enter as it rifes into the receiver; but it is eafy to substitute water as the

cools, temperature diminishes. The capacity of the receiver, and the volume of water which replaces the gas expelled by heat, is determined in the manner already described; excepting and the expan-

Con determined that in this determination to the weight of the empty receiver. by weight.

must

must be added that of a cylinder of water terminated at one and by the line of the station of the mercury, and at the other by the neck of the receiver.

I might add a few more details, but I shall suppress them for the sake of brevity, and because persons very little accus-

tomed to manipulation will eafily supply the defect.

However, as it is important, from what I have already faid to exclude water entirely from the apparatus, I shall mention how I succeeded in completely effecting this pur-

pole.

If the receiver was vifibly moift, I first dried it with blotting Methods of expaper, and then heated it so as to evaporate part of the water cluding moifture from the it might still contain, and by means of bellows, to which I gas. adjusted a glass tube, I expelled the vapour by a current of air. These last operations being repeated several times on the receiver and tube, both were found perfectly dry. The mercury which I used in my experiments was always employed very dry and pure.

In all the experiments, the refults of which I am about to The terms of give, I always reduced the gases (the dilatation of which the following could be determined by the apparatus here described) to the experiments temperature of melting ice. For this purpose I had a bath were freezing and boiling war, with ice in it, into which the receiver was wholly immersed, ter. after having withdrawn it from the bath where it was placed for the experiment, and was left there for about half an hour, during which I often stirred the ice. The other fixed temperature, at which I rested for the same gases, was that of boiling water.

I made some experiments at other temperatures; but they Experiments at require to be again repeated, and will besides form a part of other temperatures require to a work I have begun on the law of the dilatation of gases and be repeated. vapors. I shall therefore confine myself to the dilatation of gases at a fixed elevation of temperature, which shall be that comprised between the degree of melting ice and the boiling point. With regard to the vapors, I shall compare their dilatation with that of the gases.

# ART. IV. Experiments and Refults.

When I used the two apparatuses here described, but of-Expansion from tener the second than the first, and avoiding all the causes of freezing to boil-uncertainty

Of atmospheric uncertainty that I could suspect, I obtained the following fix refults from fix experiments on atmospheric air \*.

					137,40
From the temperatu	re of	melti	ng ic	e	137,61
to that of boiling v	vater	, equ	al vo	7-1	137,44
lumes of atmospheric	c air	repre	fente	1 3	137,55
by 100, became	-	10		yatti.	157,48
					137,57
The mean refult is at	out	-	-		137.50

for each degree,

If the total augmentation of volume be divided by the number of degrees which produced it, or by 80, it will be found by making the volume at the temperature O equal to unity, that the augmentation of volume for each degree is 473.43, or 166.66 for each degree of the centigrade thermometer.

not agree with Deluc.

This refult does As Deluc obtained 1 for his coefficient, it feems at first fight that our refults are the same; but if it be observed that he began at the temperature 1603, and that I began at 001 it will be feen that our refults are confiderably different. I shall explain this difference elfewhere, and fnew that the coefficients of dilatation vary according to the temperature of commencement.

drogen ;

Expansion of hi- Hidrogen gas obtained from iron by weak sulphuric acid was submitted to two experiments: in one, by an elevation of temperature from the degree of melting ice to that of boiling water, 100 parts became 137,49; and in another by the fame elevation of temperature 100 parts became 137,66. The mean of these two results is 137,52, which differs very little from the mean result of the dilatation of atmospheric air.

> \* My receiver contained about 350 grammes of water (a little more than 12 ounces) water a first in belief the en value in

> + Though the difference between these results are not very confiderable, I think I might have rendered them less so, if I could have kept an account of the state of the barometer at the moment of the ebullition of the water. I have, however, always had the precaution to be affured of its thermometrical state at the moment of ebullition, in which I can affert that I never found any very fenfible variations. A variation of an inch of the barometer would in fact be necessary to occasion one, of a degree, in the place of the boiling point; which can felcom happen. However this may be, this mean refult 137,50 must be very near the truth, inship (da)

Oxigen gas obtained from oxigenated muriate of potath was dilated three times, and gave the following refults.

oxigen,

Azote gas obtained from the decomposition of ammonia by oxigenated muriatic acid gave the following mean refults.

-azote.

By connecting the preceding refults, and by comparing the Tabulated redilatation of oxigen, hidrogen, and azote gafes with that of fults of four gases shew the atmospheric air, we obtain the following table: same expansion

From the temperature of melting ice to that of boil- ing water, 100 parts of	Increase their volume to	Differences	
140, y = 4	Parts	. Verbi	
Atmospheric air -	37,50	- 0 - 1 24	
Hidrogen gas	37,52	+0.02	
Oxigen gas	37,48	-0.02	
Azote gas	37,49	-0.01	

The trifling differences observed in the preceding results may proceed from the impossibility of rendering the circumstances rigorously the same in each experiment; and as they extend to only two ten thousandth parts of the original volume, it may be concluded with fafety, that the dilatation of atmofpheric air, oxigen, hidrogen, and azote gafes are the fame from the temperature of melting ice to that of boiling water.

In order to determine the dilatation of gases soluble in wa- Apparatus for ter I changed the apparatus, I used two tubes graduated at rates soluble in the same time on the same bath of mercury with a very small were compared measure. Every time I used this apparatus, I took the pre- by exposure to caution that the quantity of mercury should be the same as cury in a tube, when the tubes were graduated. I must even observe, that if while atmospheany accident happens to the vessel containing the mercury, the larly treated. tubes must be again graduated on another bath. It would

even be proper to cut them out of the same tube of glass and to make them of equal heights, so as to render all the circumstances as similar as possible.

The tubes with the two elastic fluids were heated in a stove. Into one of these tubes I put atmospheric air to the 100th division, for example, and in the other the gas on which the experiment was to be made likewise up to the 100th division. 100 equal measures of these gases were thus submitted to experiment. I then carried the apparatus into a stove, the temperature of which I regulated at pleasure, and I observed the course of the dilatation of the gases. Whatever attention I paid to the observation I never could perceive any difference, and I always remarked that the same divisions were passed through in the same time.

The gases were purified and dried before they were tried.

76-11-00

The gases thus examined were never directly received in the tubes; but they previously remained some time in an intermediate vessel into which I introduced some desiccative body, muriate of lime for example, and I then passed them into the tubes, compressing them with mercury, which I poured by means of a tube of safety adapted to the intermediate vessel. If these general precautions be neglected the dilatation will almost always be too great; because the influence of undissolved water will not be guarded against, or of some other body capable of easily taking the gaseous form.

Carbonic acid gas; One hundred measures of carbonic acid gas obtained from marble by means of sulphuric acid, were compared with the like portion of atmospheric air. From the 5th to the 90th degree (Reaumur) the expansions were the same,

and muriatic

One hundred measures of muriatic acid gas obtained by means of concentrated sulphuric acid from muriate of soda (very dry) being compared with 100 measures of atmospheric air from the 3d to the 86th degree, the expansions of the gases were found to be absolutely the same. This experiment, as well as the preceding, were repeated several times, and the same results were always obtained.

and fulphureous and nitrous gafes expand precifely the fame as atmospheric air. Priestley, Guyton, and Duvernois found ammoniacal gas expand more

and fulphureous Sulphureous acid gas and nitrous gas undergo the fame diand nitrous gafes latation by the action of heat as atmospheric air.

Doctor Priestley, and Citizens Guyton and Duvernois, found that ammoniacal gas dilated very considerably. With a view to discover the cause that could have induced them to draw this conclusion from their experiments, I introduced (without preparation) into one of my tubes ammoniacal gas, obtained

from

from the decomposition of muriate of ammonia by common than common lime, and in the other tube an equal volume of atmosphericair. air. The ammoniacal gas dilated progressively more than the because water atmospheric air in proportion to the elevation of temperature; and a neutral ammoniacal salt so much so that its volume was soon double that of the atmosf were present. pheric air, but on observing the surface of the mercury, and the fides of the tube after the temperature was lowered, I remarked some traces of a liquid and some crystalline speeks, which could be nothing but muriate or carbonate of ammonia, and it all disappeared when the temperature was sufficiently elevated. I performed this experiment again by fuffering the ammoniacal gas to remain fome time in an intermediate vessel containing caustic potash, and then its dilatations from 0 to 95° exactly followed that of the atmospheric air. In this experiment I observed the surface of the mercury and the sides of the tubes when the temperature had returned to 0, but I could not perceive any indication of liquid, or any crystalline speck. This experiment was repeated feveral times, and always with the fame event.

Hence it may be seen that not only a liquid, but another Water is not body capable of taking the gaseous state, may render the extherefore the periment liable to error. These causes ought therefore to be error. avoided with the most scrupulous attention.

The experiments which I have just related, and were all Hence the diffemade with great care, incontestibly prove that atmospheric air, the gases do not oxigen, hidrogen, azote, nitrous, ammoniacal, muriatic acid, affect their exfulphureous acid, and carbonic acid gafes expand equally by panfion by heat. the same degrees of heat; and that, consequently, their greater or less density under the same pressure, and at the same temperature, their more or less considerable solubility in water.

I therefore conclude that gases in general, provided they be all placed in the fame circumftances, dilate equally with equal degrees of heat.

and their particular nature, in no respect influence their dila-

These experiments on the dilatation of gases naturally led Vapors. me to examine that of vapors; but suspecting from the preceding refults that they dilated like the gases, I determined to make experiments on one vapor only, and I chose in preference that of fulphuric ether, as being the most easy to manage. obtained'

I employed the two tubes I have already mentioned, atmof-

The vapor of ether was tried in the tube ; in atmosph. air.

They expanded alike between 60 and 1000 Reaumur,

When near its ethereal vapor was condensed rather more.

comparison with pheric air being always the standard of comparison. This apparatus having been kept fome time in a stove whose temperature was about 60°, I introduced ethereal vapor into one of the tubes, and into the other atmospheric air, so that they each corresponded to the same division. I then elevated the temperature of the stove from 60° to 100°, and I had the fatisfaction to find, that whether the temperature was raifed or lowered, the ether and the atmospheric air always corresponded at the same time to the same divisions. This experiment, at which Citizen Berthollet was prefent, was repeated feveral times, and I never could observe any difference in the dilatation compared with that of atmospheric air. I remarked, boiling point the however, that at some degrees above the point of the ebullition of ether, its condensations were a little more rapid than those of atmospheric air. This is owing to a phenomenon which a great number of bodies present in passing from the liquid to the folid state, but which does not present itself a few degrees above that of this period.

The conclusion may therefore be extended to vapors;

This experiment, by proving that the vapor of other and the gases dilate equally, shews us that this property in no refpect depends on the peculiar nature of gafes and vapors, but only on their elastic state, and we must consequently conclude that all gases and vapors dilate equally at the same degrees of

The vapors will be more compressible near their point of liquefaction.

Since all the gases are equally dilatable by heat and equally compressible, (gas condensable) and as these two properties depend on one another, as I shall prove elsewhere, vapors dense flu dity or being equally dilatable with the gases, ought also to be equally compressible: but I must observe that this latter conclusion cannot be true, unless while the compressed vapors perfectly retain the elastic state; which requires that their temperature be fufficiently elevated to enable them to refift the preffure which tends to make them assume the liquid state.

That dry and moist air dilate alike .- Qu. ?

I have afferted on the authority of Saussure, and my own experiments confirm it, that very dry air, and air holding more or less of water in solution are equally dilatable; confequently I am authorized from the whole of the facts to draw the following conclusions.

Ift. All

If, All the gases, whatever may be their density or the Conclusions. quantity of water they hold in folution, and all vapors, equally Expansion of dilate by the same degrees of heat.

2d, The augmentation of volume which each of the perma- Augmentation nent gases receives, from the temperature of melting ice to of volume. that of boiling water, is equal to \$\frac{80}{213,33}\$ of the primitive volume, if judged by the thermometer divided into 80 parts, and 100 of the same volume, if by the centigrade thermometer.

It remains, in order to complete this course of experiments, Future experito determine the law of the dilatation of gases and vapors, in ments on the laws of dilatation order to ascertain the coefficient of dilatation for any degree at at other tempea known heat, and to be affured of the real progress of the ratures. thermometer. I shall employ myself on these new researches; and when they are terminated, I shall have the honour of communicating them to the Institute.

#### VII.

New Theory of the Constitution of mixed Gases elucidated. Letter from Mr. J. DALTON.

## To Mr. NICHOLSON.

SIR,

Manchester, Nov. 18, 1802.

" would

IN a paper of mine which you had the goodness to publish in New theory of your Journal for 1801 (Quarto Series, V. 241), I announced the conflictution a new theory of the constitution of the atmosphere. This has sphere, fince been published on a more enlarged scale, and elucidated by a plate, &c. in the Memoirs of the Literary and Philosophical Society of Manchester, Vol. V. Part 2. Notwithstanding this, I am informed by fome of my chemical friends, that they do not clearly understand the hypothesis itself, and confe- not clearly unquently are not able to judge of its merits or defects: And a derstood. late writer (Dr. Thomson), in his System of Chemistry, Vol. III. Page 270, speaking of the uniform distusion of the different gales of the atmosphere, makes the following observation: Even Mr. Dalton's ingenious supposition, that they neither " attract nor repel each other, would not account for this equal distribution; for, undoubtedly, on that supposition, they

tion.

Intended order of difcuffion.

"would arrange themselves according to their specific gra-Farther explana-" vity." Now as I am perfuaded that no one acquainted with the principles of mechanical philosophy would have written the above if he had understood my hypothesis, it seems to call from me a further explanation. I propose, therefore, 1st, To state in as clear a point of view as the subject will admit, the principles which I assume: 2d, To shew that the consequences which I have deduced from them are legitimate, and particularly that mixed elastic sluids ought not to arrange themselves according to their specific gravity: and, 3d, To demonstrate that the supposition of the gases constituting the atmosphere being held in a flate of equal diffusion by chemical affinity, is not only inconfistent with the phenomena, but is completely abfurd.

#### 1. PRINCIPLES ASSUMED.

Principles affumed. 1. That particles of fimple elastic fluids repei inversely as their distance.

1. I take for granted that the particles of simple (unmixed) elastic fluids repel one another with forces inversely as the distance of their centres, the temperature being given. is a mathematical deduction from the allowed fact, that the space occupied by any gas is inversely as the compressing force: (See Newton's Principia, B. II. Prop. 23). The absolute distances of the centres of such particles must vary according to circumftances, and cannot eafily be determined; their relative distances in a liquid and aërial state sometimes may. Mr. Watt has shewn, that steam of 212° and pressure 28 inches, is 1800 times lighter than water; confequently the diftances of the particles of fteam are to the distances of the said particles in a liquid flate, as 12 to 1 nearly, in that particular case. Vapor in the vacuum of an air-pump, at common temperature, will have its particles about 4 times the distance, or 48 to 1.

2. I suppose that in mixed elastic fluids the heterogeneous 2. In mixed elastic fluids the particles do not repel one another at all at fuch distances as heterogeneous particles have no they repel those of their own kind; but that such particles, distant or gaze- when brought into actual contact (to use the common lanfiant repulsion; but only that of guage), refift each other in all respects like inelastic bodies, contact, formerly This is the peculiarity of the hypothesis, and what appears not called the refiftance of impene- to be generally understood. If I may explain by analogy, the most striking will be found in magnetism. Two like poles of Illustration from the vifible effects magnets repel one another with the same force whether any other of magnetifin.

other bodies intervene or not, and do not affect those other bodies; in the same way I conceive two particles of any one gas repel one another with the same force whether particles of other gases intervene or not, and do not affect those other particles. A magnet is amenable to the common laws of motion in its collision with other bodies, and when it is brought into feeming contact with them; fo is a particle of one gas when it is brought into seeming contact with a particle of another species: and in this case the bodies may be said to have a repulfive power; but this power is essentially different from the others, in that it extends to no definite distance. Further, conceive a very fine capillary tube placed perpendicular to the horizon, into which let a number of correspondent finall magnetic wires, or particles, be inferted with their poles of the same denomination together, or more strictly, as near as their repulfive power would admit, one particle above another, the air having intercourse amongst them. Then, as the magnetic particles would not actually touch one another, by reason of their repulsion, they would feem to be supported by the intervening air, whereas in reality they are supported and kept at certain distances intirely by the repulsion inherent in themfelves, and their own gravitation: and thus I conceive par-Particles of gas ticles of gas support those of their own kind above them, support each other by repultions, were they visible, they might seem to rest upon others sion, in the same immediately under them; and the ground, or lowest folid or manner; whether liquid surface, by supporting the lowest particle of each kind, another gas be has the weight of the whole to fustain. These observations, present or not. together with a view of the plate above alluded to, must, I think, be sufficient to satisfy any one what the hypothesis is. And it may be proper to add, there is fomething much refembling polarity observable in the ultimate particles of bodies at the instant of transition from the liquid to the folid state; wit-

# 2. CONSEQUENCES.

ness the congelation of water.

It is plain from the above account, that I conceive any one If a gas be one gas to be conflituted of perhaps one part folid matter, and one part matter and a thousand mere thousand or more parts vacuity or pore, if it may be so called; space, no differ and that into this vacuity we may throw as many other gases ence of effect will follow from as we please, without materially disturbing the first, provided other gases in we do not absolutely fill the vacuity with solid matter, meaning that vacuity, &c.

by '

And one gas will not raife any other, whatever dividual specific gravities;

by that expression common liquids or folids. Thus we might have had a dozen gases in our atmosphere instead of three or four, all in the same compass, and each retaining the same density it would have had alone. The heavier gas has no more tendency to raife the lighter, than a quantity of shot has may be their in- to expel the air from its interstices. If therefore Dr. Thomson, or any other, can shew how one fluid, which is not displaced nor any way acted on by another, should by its reaction cause that other to move into a higher or lower station; then the mathematical world will be obliged to reconfider their doctrine of statics. Till then I must take the negative of the proposition, and conclude, that elastic sluids of the greatest and least specific gravity imaginable, on the supposition I hold, will alike take the lowest and the highest stations, regardless of each other; or in other words, they will arrange themselves in the fame order as if thrown into a complete vacuum. The great difficulty respecting the uniform diffusion of the gases being removed, I think on my hypothesis the other phenomena can require no explanation to any person conversant in pneumatics,

and the chemical I will take one inftance: It may by asked, How does sulphuret absorption of any of potalli abstract oxigenous gas out of any mixture; lime fingle gas will be the same as if water, carbonic acid gas, &c. &c.? The answer is obvious: it were alone. Exactly in the same way as if the gas in question was the only one in the vessel, and the operation going on in a close vessel.

3. It is contrary to fact that gales are held together by chem. affinity: For they mix uniformly, and

## 3. GASES HELD TOGETHER BY CHEMICAL AFFINITY, ABSURD.

On this head it will be proper to premise certain facts:

1. When two gafes of different specific gravity, such as oxigenous and hidrogenous, are put into the same vessel and agitated; if they be suffered to stand some time after, they still continue uniformly mixed.

they occupy the same space after mixture;

remain fo;

2. They occupy the same space before and after mixture; that is, one measure of each put together occupy two measures, the temperature and pressure remaining the same. Mr. Davy feems to think this principle is not firictly true in regard to a mixture of azotic and oxigenous gas; but the deviation from it, if any, is extremely small.

3. The compound is subject to the same laws of rarefaction and they expand, &c. as before. and condensation as the simples.

There

There are but three suppositions we can make essentially Where gases different respecting the mutual action of heterogeneous par-mix, the particles must eiticles of gas: 1st, When two gases are mixed their particles ther, (1.) repel, may reciprocally repel one another, just as they act on their own or, (2.) have no kind in an unmixed state: 2d, They may be neutral, or have attract. neither attraction nor repulsion for each other: 3d, They may have a chemical affinity or attraction for each other. The ad- The gafes do not vocates for the chemical adhesion of gases will agree with me fin: if they in exploding the first, because where nothing but repulsion is have no action manifest, we can ascribe no effect to attraction: The second, ory is good; which is the one I adopt, is obviously inconsistent with their and, hypothesis; and as for the third, I can conceive no other explanation than the following: 1/t, Two or more heterogeneous if the particles particles may unite and become a new centre for the caloric attract and unite to adhere to; but in this case the gases are no longer two but gas, and not two, one; and oxigenous and hidrogenous would become aqueous as in the affumed case: vapor: This therefore is not a case of two gases being held together by chemical affinity. 2d, The two gafes may fepa- Or if they atrately retain their caloric and still be held by chemical affinity; tract and do not that is there may be an aguilibrium between the remarks that is, there may be an equilibrium between the powers of come nearer toattraction and repulsion; but this is evidently inconsistent with gether, contrary the third law of condensation and rarefaction observed in such or fact above compounds.

I am, &c.

J. DALTON.

I thank you for the notice you have taken of my Essays, and particularly for the note subjoined to that on the Expansion of Gases. I have lately read a Paper to the Literary and Philosophical Society of this place, in which I have shewn that the quantity of carbonic acid gas found in a given volume of atmospheric air, is not more than  $\frac{1}{1000}$  part of the whole; and that the said gas is held in water, not by chemical affinity, but merely by the pressure of the gas abstractedly considered on the surface, forcing it into the pores of the water.

A Method of increasing the Quantity of Light afforded by Candles. and to obviate the Necessity of shuffing them. In a Letter from Mr. EZEKIEL WALKER.

#### To Mr. NICHOLSON.

SIR,

Feeble light of

WHEN our apartments are illuminated with common frequently four-candles, which cannot be regularly attended with the fouffers, they produce much smoke, and their glimmering light scarcely ferves the most ordinary purposes. My first attempts to im. prove this fombre mode of illumination were made many years ago. Those however were not productive of any useful discovery, but some experiments which I made in the year 1797 afforded refults that rather exceeded my expectation.

Method of remedying this in candles for illuminating apartments.

Experience foon convinced me of the utility of the method which I had discovered, of using candles in such situations as above described, but in every attempt to produce the same effect upon my table, I was unfuccefsful, till the beginning of the present season. This was owing to an oversight, which might be deemed very unaccountable, did not the history of inventions contain many inflances of the fame kind.

Great effects are frequently produced by causes that are in general deemed trifles. And, if a trifling alteration be made in the method of using common tallow candles, they will become excellent substitutes for those of wax.

If a candle be a steady light, and requires no inuffing;

&c.

A common candle, weighing in of a pound, containing 14 inclined it gives fingle threads of fine cotton, placed fo as to form an angle of 30 degrees \* with the perpendicular, and lighted, requires no fnuffing; and what is much more valuable for some purposes, it gives a light that is nearly uniform in strength without the least smoke. These effects are thus produced:

When a candle burns in an inclined position, most part of because the wick protrudes into the flame rifes perpendicularly from the upper fide of the the air, and becomes burned,

> \* Candlesticks may be made to hold the candle at this angle, or they may be so contrived as to hold the candle at any angle at pleafure. I have candlesticks constructed on both principles.

> > wick.

wick, and when viewed in a certain direction, it appears in the form of an obtuse angled triangle. And as the end of the wick projects beyond the flame at the obtuse angle, it meets with the air, and is completely burnt to ashes; hence it is rendered incapable of acting as a conductor to carry off any part of the combustible matter in the form of smoke. By this spontaneous mode of snussing, that part of the wick which is acted upon by the slame, continues of the same length, and the slame itself very nearly of the same strength and magnitude \*.

The advantages which may be derived from candles that The light does require no fnuffing and afford no fmoke, may be readily understood; but these candles have another property which ought not to be passed over in silence. A candle snuffed by an instrument gives a very sluctuating light, which, in viewing near objects, is highly injurious to the eye; and this is an inconvenience which no shade can remove. But when a candle is snuffed spontaneously, it gives a light so perfectly steady and so uniformly bright, that the adjustments of the eye remain at rest, and distinct vision is performed without pain, and without uneasiness.

Candles on which I have made experiments, are described Candles which in the following were subjected to experiment.

TABLE.

No.	No. of candles to the pound avoirdupois weight.	Length in inches.	No. of fingle threads of fine cotton in the wick.
	14	0.7	10
2	12	8.5 9.	10 12
3	10	9.7 <i>5</i>	14
4	8	10.	20
5	6	10.25	24
Mould -	6	13.	

Numbers 1, 2, and 3. These candles, when lighted and placed to form an angle of 30° with the perpendicular, require

The wick's not being uniformly twifted throughout, may occafion a little variation in the dimensions of the flame.

no fnuffing: they give lights which are nearly equal, and combustion proceeds so regularly, that no part of the melted tallow escapes unconfumed, except from accidental causes.

Remarks on the phenomena.

No. 4, placed at the angle mentioned above, and lighted, requires no fnuffing: it gives a light very little stronger than No. 1, but its colour is not quite so white, nor its slame so steady.

No. 5. This candle, placed at an angle of 30°, and lighted, requires no fnuffing; its flame is rather fluctuating, and not fo white as No. 4, nor is its strength of light much greater than No. 1. The melted tallow fometimes overflows when the air in the room is put in motion; yet the light of this candle is much improved by the candle's being placed in an inclined position.

The mould candle, treated in the same manner as those above, affords a very pure steady slame, without smoke and without fnuffing, and its strength of light is about equal to that of No. 1.

#### ON ECONOMY.

The economy of candles as to light and confumption of the fuel.

My experiments have not been sufficiently numerous to determine with precision which of these candles affords the most light at a given expence, but the few experiments which I have made feem to indicate, that the quantity of light is directly as the quantity of combustible matter expended.

EZ. WALKER.

Lynn, Nov. 19, 1802.

## REMARKS.-W. N.

On candles in general.

On candles in general fee an effay by W. N. in the Philof. Journal, quarto, I. 67; also my First Principles of Chemistry, Art. Wax: Hermstadt on the different materials, wax, tallow, fpermaceti; Philof. Journ. quarto, V. 187.-I have always fuspected from the general facts, that the light of small candles is cheaper than that of larger, and have little doubt of the truth with regard to fuch as exceed the diameter of three quarters of Outline of very an inch in tallow. We are much in want of a feries of ex-

defirable experi-periments on the different descriptions of candles, under the general ing them.

general titles of Wax, Spermaceti, Tallow, and mixed; and tabulated, to shew, 1. Average weight of the wick; 2. Number of threads; 3. Weight of the whole candle; 4. Diameter; 5. Length; 6. Time of combustion per inch; 7. Ditto per ounce; 8. Quantity or intensity of light when newly snuffed; 9. Ditto after burning half a minute, or till the usual period of fnuffing; 10. Average intensity; 11. Expence of material per hour to produce the quantity 1.00 of light; 12. Ditto in the money price of the article; 13. Station of the barometer; 14. Ditto of the thermometer; 15. And ditto of the eudiometer. I am absolutely unacquainted with any good standard of com-Standard of the parison of light. The sun perpetually varies with the weather, unknown, its altitude, &c. It seems probable that an earthen lamp, having a wick of known texture and weight, cut to a determinate elevation by a gage, and charged with very pure olive oil, might, at like stations of the barometer and thermometer, give light fo nearly equal, for the first two or three hours, as to afford refults confiderably exact; of which the means of feveral repetitions would be of great practical utility. The compari-Method of coma fon of lights by the shadows has been often mentioned; but the paring lights by fimplicity and value of the method (not yet univerfally known) will justify a short repetition. When the shadows of the same object fall upon each other, not precifely, the bordering shades will be feverally illuminated by one light only: move the lights till these bodies appear equally dark at the point or place where they meet in an angle. Then measure the distances of the lights from that point or place, and the intensities are inversely as the squares of those distances. It is to be noticed that the obstacle which forms the shadow, a book for example, must be held so that the ray from each light may fall on the wall nearly in the fame angle; and this will always be the cafe if the farther light be nearly behind the other.

In a subsequent letter, Mr. Walker informs me, that he carefully measured his intensities by the method of shadows, as mentioned in Prieftley's Optics. I conclude this to have been the method here described; though, for want of a good index, I can only find the method of Bouguer in that work.

#### IX.

Description of a new Secret Lock. By J. B. BERARD.

(Concluded from Page 222.)

cessible only to the proprietor.

MAY also in this place state the reason why I have made a fecret lock ac- the key as well as the ferrils with 24 rather than 25 divisions. In the first place, it affords a small diminution in the dimensions of the lock. Another more effential advantage is, that in the division of 24, each notch in the ferril is opposed to another diametrically opposite, which facilitates the introduction and taking out of the bolt. We may observe, that for the same reason the teeth of the ferrils, being rather rounded at the tip, facilitate the entry of the little wedge, which is likewise rounded underneath.

The lock being thus conftructed, it only remains to fix it to the door. For this purpose, an opening is contrived of the fize of the box of the lock, in the horizontal cross piece, without cutting the vertical frame of the door; and round it a groove is made, of 2 in depth and 6 in breadth, on which the face of the lock refts: the lock is placed in this opening without the bolt; and if it is defired, four nails or fcrews may be driven into the wood through four holes made in the face of the plate; but this is not necessary. The bolt is then placed on the other fide of the door, and if the wood be too thick, it may be cut away till the bolt flides fair and eafily over the wood. This is very eafy to be done, and then the face of the bolt ought to lie flush with the edge of the door.

It is obvious that the height of the ferrils must be proportioned to the thickness of the door. If the door be 30 m. thick, for example, the ferrils (which ought to be 1 under the furface of the lock) would be about 29.

I must also observe, that in order to render the motion of the screw-heads more easy and uniform, a pasteboard of 1 in thicknefs, with five holes punched in it, must be laid upon the inner furface of the plate of the lock; and above the pasteboard, another flip of very thin brafs, likewife perforated with five holes: then the interior flat wheels have a very eafy motion over the brass which rests on an elastic plane, which yields to Construction of the unavoidable irregularities of execution in a work of a mocessible only to the proprietor.

It must likewise be observed, that the heads of the screws, as well as the sace itself of the lock, ought to be case hardened, in order to protect it from external injuries, and to prevent the enlargement of the holes which receive the screws. The surface ought to be highly polished, and the heads of the screws should be blue.

# Advantages of the new Lock.

1st, As in Reignier's lock, which likewife has fcrew-heads, ferrils, and teeth without wedges, these pieces cannot be put at distances extremely precise, it necessarily follows that one of the pins or teeth, the fourth for example, will always meet its ferril before the others meet theirs. Let us suppose that with a turnfcrew we endeavour to turn the fcrew-head which moves the bolt, whilst with another hand and turnscrew we fuccessively turn the other screw-heads, they will all turn freely, except a fourth, which will be immoveable: It will be concluded that the fourth fcrew-head is not in the position to open the door, that is to fay, that the deep cut of the ferril is not opposite the pin of the bolt: that figure or letter of the fcrew-head will be noticed among those which do not form the fecret. The position of the screw-head must therefore be changed, and the above process repeated till the pin is felt to fall into the notch. The same experiment must then be made with that screw-head among the remaining ferrils which alone is now pressed by its pin. And after eighty attempts at most, if there be eight forew-heads, each having eleven figures, we shall know the eight figures which compose the secret.

Here then is a regular and infallible process to discover in less than an hour the combinations with which the lock is closed. This desect, which is common to the generality of compounded locks, does not exist in mine.

When my bolt is moved, the wedges which precede the teeth enter at the same time into the cuts of the ferrils, which are of such a height that they graze the interior surface of the lock: all the four ferrils then become immoveable, and it is impossible to discover which is the ferril that is first met by the correspondent tooth, because their fixation is produced at the

ceffible only to the proprietor.

Construction of same time from the teeth and the wedges. No external exaa fecret lock ac-mination can therefore shew the difference in situation of the four ferrils. It is obvious that in order to infure this effect, the wedges ought to fit the cuts between the teeth in the ferrils exactly; which is very eafily obtained in the construction. Another advantage resulting from the wedge is, that if any of the deep notches or openings be not exactly in a line with the rest, the wedge easily adjusts it, and the bolt is not in the least impeded on that account; this advantage renders it unnecessary that the workmanship of the lock should be exquisite, a condition which, in a work of a moderate price like the present, would render it useless.

2d, My lock is eafily opened, it having only four fcrewheads: it has nevertheless \$31,776 \* combinations, a number fully fufficient to fet the most fuspicious miser at ease.

We may add, that it is much easier to recollect one word than

a number, which must be the case in Reignier's lock.

3d, When characters are engraven externally on the screwheads, they are either difficult to be perceived, or may be eafily damaged. A key with characters engraved on it, is I think a great improvement: a faving is likewise afforded, which is a confideration of fome confequence in a machine of common nie.

4th, The method by which I vary the position of each ferril with respect to its screw-head, is at once simple and solid.

5th, The mechanism by which the bolt is attached to the lock, is concealed, and cannot be damaged. As nothing is feen of the lock but a smooth plate on either fide, it would be very proper for a bureau or 'scrutoire.

Another fingular property refulting from this mechanism is, that it can only be taken to pieces by the proprietor, whether the door be open or shut: it could not even be done by a perfon that in the apartment.

6th, The lock may easily be set: as the interior screwheads do not turn on wood, there is no danger that their motion should become less regular or easy on account of wear. The same may be observed of the bolt, which slides in a groove of iron.

<sup>\*</sup> This is the fourth power of 24.

7th, The lock is very folid; for there is no opening into Conftruction of which any tool might be introduced. The face of the door a fecret lock acmay likewife be furrounded with a strong metallic covering let the proprietor, into the wood, and the back part of the door may be covered with iron plates, the only method of preventing any one from cutting the wood round the lock.

8th, If it should be thought inconvenient to place the lock in the thickness of the door it may be easily avoided. The stems of the screw-heads need only be made longer, and another plate similar to that of the face, added: the screws in that case, after having passed through the sace as well as the door, would pass into the lock. This addition requires an increase of work, but it affords no real advantage.

## Lock for a Bureau or Secretary.

Though my lock, as I have described it, can be easily fixed to a secretary, yet I think it useful to indicate the changes convenient to be made in it.

1st, The height of the ferrils must be diminished, that the bolt may be intirely sunk in the stap of the secretary.

2d, In order to diminish the length and breadth of the lock, ten divisions in the ferrils may be made instead of eighty: they need then be only 25 in diameter; but then the secret will consist in a number of sour figures instead of a word of sour letters, which is not so convenient, as I have already observed.

3d, The box of the lock may be suppressed, and the lock will still have solidity sufficient for a bureau; the price will likewise be diminished.

4th, Instead of engraving characters on the face of the lock, I preser, as has been seen, to mark them on the turnscrew itsels; but the external damage is not to be feared in a secretary as in a door. It is therefore more convenient that the sigures or letters should be on the sace itself of the screw, and a knob or small handle may be contrived on the sace of the screws, in order to turn them. In this case no key or turnscrew need be carried about; this is an advantage of some consequence.

# Concerning fome other Kinds of Compound Locks.

It is possible, as I have already observed, to apply the principles of these combinations to the construction of locks in a

number

Construction of number of methods. That which I have described is the last a secret lock ac-result of my endeavours on this object; it seems to me to be the proprietor. It think, however, that it will not be useless to make known some more, if it were only to prevent others from making unnecessary attempts on the same subject.

1st, The front part of the lock presents 25 small rectangles, 10 wide by 15, disposed in a right line, and forming in all 250. The 24 letters of the alphabet are marked on these rectangles: they are included between the two partitions of the horizontal channel or groove, and they all touch one another. They may be pulled outwards to the distance of 4 millemetres, like the stops of an organ. Each of these stops has a little spring at top, which gives it an easy and smooth motion, and prevents it from dragging with it any of its neighbours. Each stop, in the inside of the lock, has a projection in the form of a tooth.

The bolt has, on its fuperior and horizontal face, 24 dove-tailed grooves with dovetails, each of which receives a register or stop of the same form, having a tooth at top. These stops of the bolt may be put into their grooves with the extremity having the tooth, or the other end, just at pleasure. The bolt is enclosed in a box, the visible and interior face of which moves in a groove, in order to take out and replace the bolt at pleasure.

The lock is thus used. If it is to be opened, the exterior stops forming the secret are brought forward, then the 25th stop is drawn, and this renders the rest of the stops immoveable. After having returned it, it is turned by means of a knob or handle which it carries, and the bolt is thus made to move.

When the fecret is to be changed, the interior registers are to be placed in the grooves cut in the bolt, by the extremity which has no tooth. They are then inverted with respect to the others, and their teeth become in a line with those of the exterior stops.

The number of combinations in this lock is equal to the eightieth power of 2, nearly 17 million: It is convenient for a bureau, but the workmanship must be exquisite: Besides, it is not convenient for a door, because stops which project several millimetres may be easily damaged. I cannot here enlarge upon the other impersections which it presents. In this system

the

the 24 small screw-heads might be substituted for the stops, Construction of which would not then require ferrils. This arrangement would a fecret lock acceptible only to not be preferable to the other.

the proprietor.

2d, Let us imagine three rings lodged one within the other, and forming one and the same plane with a fourth fixed to the door, and an immoveable circle in the centre; fuch is the face of the lock. The three intermediate rings turn freely and feparately by means of a button, which must be made to correfound with one of the letters of the alphabet marked on the outer circle fixed to the door. Each carries interiorly a ferril capable of 24 positions. When the notches of the three ferrils are disposed in straight lines, the bolt is made to move by means of a knob or handle in the middle of the face.

This lock \* refembles in some respects the first I have defcribed. In this the ferrils are concentric, and in the other's they are separate. It has an agreeable form, and would be proper for a fecretary, and its execution does not require much precision. It is not very convenient for a door, because it is

likely to be damaged.

. 3d, Let us imagine five forew-heads on the face; let each of these screws carry a small pinion within, which moves a notched bar or rack. This rack must have ten teeth, one of which is withdrawn, in order to leave a clear way or space. The fifth fcrew must move a fifth rule carrying four teeth, and at right angles to the other four. It is clear that this last rule, which I suppose to be the bolt, can never move except when the other four rules are in a convenient position. In this method the lock will be enormously large; and it will also be very difficult to prevent this fecret from being discovered from without.

4th. If toothed wheels, with an unequal number of teeth all taking in one another, be substituted for the flat wheels which are under the ferrils in the fystem of locks with screwheads, then, if one screw-head be moved, it will derange all the ferrils at once. This faves time, without diminishing the number of combinations. This advantage led me to approve of the construction for some time; but it is productive of so many ferious inconveniences, that I was obliged to abandon it.

<sup>\*</sup> I have feen some at Paris, nearly similar to this.

Conftruction of ceffible only to the proprietor.

It would be superfluous to examine a greater number of a fecret lock ac- systems. I shall conclude by one general observation.

All the compound locks that can possibly be made, may be divided into three species. In the first, straight or curve lines are made to move; in the fecond, plain or curved furfaces; in the third, folids. It is obvious that one folid offers as many arrangements or combinations as three lines together, and one furface as many as two lines.

These lines, surfaces, and solids, form interior stops made to receive motions (externally produced) in a straight or in a curve line.

It is obvious from the application of these principles, an almost infinite variety of locks may be constructed. I shall not undertake to extend the variations, but shall only fay, that I think I am convinced that the best system is that of moving the circumferences of circles internally by circular motions produced from without,

#### CONCLUSION.

I am fensible that what I have said concerning the locks, which I confider as less perfect, is not sufficient to direct their construction; but I have said enough to make their system known to artists, and to deter them from fruitless experiments. It must not be forgotten, that in this kind of machine nothing good or useful shall be done, unless simplicity, convenience, folidity, economy, and facility of execution, be combined.

It is under these several points of view that the problem of compound locks, which may be worthy of the attention of artiffs, ought to be confidered. I do not pretend to have difcovered the best, but I dare affert, that the lock which I have described approaches so near to perfection, as to entitle it to the fuperiority over the common locks which are more or less ingenious and complicated, and even to place it among the locks of combination, as one of the most simple that has yet appeared, or that I know of.

Whether I shall be allowed the small honour of having invented this arrangement, or the merit only of having brought it to perfection, is not to me a matter of confequence. I shall be satisfied if I have assisted others in improving this species of machines, the utility of which need not be proved. In fact,

the common locks which are brought to the greatest perfec-

tion, were rendered as fure and inviolable, as they are at present the contrary, they would still have this inconvenience: they are liable to be forced by introducing some tool or instrument; and even if the violator should fail in his attempt, the proprietor will nevertheless be reduced to the necessity of breaking open his own door if the infide of the lock should be ever fo little deranged. The advantage then of a lock which affords no entrance to a key is obvious, a problem which I think cannot be well folved, except by a circular motion. That which I have described, not only answers all the conditions of the theory, but has flood the test of experience. greatest obstacle to the propagation of locks of combination, proceeds from the prejudices of habit, still more than from their want of simplicity: and it is well known that this is the fate of all truths \*.

#### X.

Description of Dr. Young's Apparatus for exhibiting the Colours of thin Plates, by means of the Solar Microscope. the Journals of the Royal Institution, I. 241.

I HE colours of thin plates were observed by Boyle and The colours of Hooke, and more accurately analysed by Newton: but little thin plates have been little exaor nothing was added to the account that Newton gave of mined, and nethem, until fome attempts were lately made to explain them, ver publicly exand to build at the fame time on the explanation, the principal arguments in favour of a new fystem of light and colours. The phenomena themselves were very little known, except from Newton's description; it had happened but to few to observe them: and they had never been made conspicuous to a public audience in a form equally beautiful and interesting.

It appeared, however, that there would be little difficulty New attempt to in applying the apparatus for reprefenting opaque objects in exhibit them by the folar microscope, to the exhibition of these colours on a cope. large scale: but several precautions were necessary, in order to obtain the most advantageous representation; and, these precautions having been completely fuccessful, it may be of fome utility to give a detached account of them.

\* In the former Paper for Requier read Reignier, and in the note on p. 217, for Braviali read Bramah.

The

Methods of producing the colours.

Mr. Jordan's convenient method of exhibiting the foapy film.

The colours of thin substances must often have been seen in bubbles of water or of other fluids, and in the film produced by a drop of oil spreading on water; they were more particularly observed in the plates of talc, or of selenite, into which those substances readily divide. Sir Isaac Newton made his experiments principally on the colours of foap bubbles, and on those which are produced by the contact of two lenses. inspecting the colours of soapy water, the most convenient method is that of Mr. Jordan. He dips a wine glass into a weak folution of foap, and then holds it in a horizontal position against an upright substance, for example, a window shutter; the film covering the glass being in a vertical position, the gravity of the fluid tends to make it thicker at the lower part, and it becomes every where gradually thinner and thinner, till at length it bursts at the uppermost point. colours assume, in this case, the form of horizontal stripes. fimilar to the rings which are to be more particularly defcribed.

The colours from a dense plate are more vivid than those from a rare one; because the latthe light reflected from the confining mediums.

Remedy.

It has been observed by Newton, that the colours thus reflected from a plate of a denser medium, are more vivid than when a plate of a rarer medium is interposed between two denfer mediums. But the cause of this apparent difference is ter are diluted by probably, the quantity of foreign light that is generally prefent in the experiment, reflected as well from the upper furface of the fuperior medium as from the lower furface of the inferior, both these surfaces being often nearly parallel to the surfaces in contact. It becomes therefore defirable to remove this foreign light: this may be done effectually, by employing one glass in the form of a prisin, and coating the lower surface of the other with black fealing wax: the light reflected by the oblique surface of the first, is thus thrown into another direction; and the reflection of the inferior furface of the fecond, is either destroyed or rendered imperceptible. And, with these precautions, the rings of colours produced in the reflected light, may be rendered a very beautiful object by means of the folar microscope.

The plane glass is made prismatic, and the lower fulface of the lens coated b.ack.

The most perfectly plane glasses are those which are used for Hadley's quadrants: one of these may be ground in the direction of the diagonal of its transverse section, so as to make a thin wedge or prism; and the surface of the lens employed must be a portion of a sphere of from sive to ten feet radius.

The two glasses must be retained in their position by means of three screws; for, as soon as the pressure is removed, they repel each other with confiderable force; and, for this reason, neither of them ought to be very thin, otherwise they will bend before they are fufficiently wear.

For adjusting the glasses to the microscope, it is convenient This apparatus to fix them in a cylinder of fufficient fize to project beyond the may be fixed in the folar miglaffes and their fcrews, in order that they may be readily crofcope. turned fo as to reflect the light coming from the speculum into the direction of the axis of the microscope: it is obvious that, in this case, they must be somewhat inclined to the light, so that the focus of the whole image will never be equally perfect; and, instead of being circular, like the rings themselves, their images on the screen will be oval. In this manner, eight The effect or ten alternations of colours may eafily be observed; but their order and fequence is too complicated to be eafily underflood: for they are really composed of an infinite number of series of rings of different magnitude, each feries being formed by each of the gradations of light in the prismatic spectrum, which, near the centre, are fufficiently separate to form distinct appearances, either alone or in combination; but, after eight or ten alternations, are lost in the common effect of white light. For, when the glaffes are illuminated by homogeneous light with homogeneous light ous light, only, separated from the rest by the refraction of a prism, or otherwise, the rings of each colour occupy, together with the dark fpaces, the whole visible furface, their number being only limited by the power of the eye in perceiving objects fo minute as the external ones become, in confequence of the rapid increase of the thickness of the plate of air near the edges of the curved furface. This circumstance being once under- and still more flood, it is also capable of being illustrated in a manner still elegantly by a more elegant, by placing a prism a few feet from the microfcope, leaving only a narrow line of its furface exposed to the incident rays, and then throwing the rings of colours on it, in fuch a direction that this line shall pass through their centre. Care being taken to exclude from the prifinatic spectrum thus formed all extraneous light, it exhibits a most interesting analysis of these colours; for the line consists of portions of the Interesting efrings of all possible gradations of colour, each forming a broken feet. line, but not of the fame dimensions; and, by the prilmatic refraction, all these broken lines are separated and placed pa-

rallel

rallel to each other, on account of the different refrangibility of the light of which they confift. Thus the broken line of the extreme red, which confifts of the longest portions, is least refracted: the other reds follow, and are placed in contact with the first, and with each other, but on account of the different magnitude of the portions, somewhat obliquely. The dark spaces also are in contact, and form a separation between each portion of light. In the same manner, the green follows the red, with little or no visible yellow. The blue and violet are somewhat mixed: for these two colours are much less widely separated by thin plates than by the prism: for this reason, each portion of light formed by the contiguous lines of the different colours is bounded not by straight but by curved lines.

It is evident that, by drawing a line across this compound spectrum at any part, we may learn the component parts of the light constituting the rings at that part; for the prism only spreads the colours in a direction transverse to this spectrum; and it may be observed, that after the eighth or tenth alternation, the light transmitted at each point is so mixed, that we may easily understand how it appears white.

Colours by transmission are not so advantageously shewn.

The colours of thin plates, as feen by transmission, are also easily exhibited in the solar microscope; but, since it is utterly impossible to exclude the very great proportion of the light which is not concerned in their formation, they never appear so brilliant as the colours seen by restection.

Y.

### XI.

Extract of a Letter from R. Chenevix, Esq. on the Magnetic Property of Nickel.

Nickel is mag

I AM afraid that I was too hasty in denying the magnetic property of nickel. The fact is, I had obtained both it and cobalt in that state. But in profecuting a feries of experiments upon these metals, I perceived the real cause why several fragments which I had obtained did not exert any influence upon the magnet. I took a piece of nickel which was not attractable, and which exhaled arsenical vapours when

heated by the blow pipe, and dissolved it in nitric acid. I Examination of boiled the folution until the whole of the metal was converted a piece not maginto arfeniate of nickel; I then dropped in a fufficient quantion in nitric tity of nitrate of lead, and evaporated the liquor at a very acid; ebullition; gentle heat, but not quite to drynefs. Alcohol poured into added; evapor. this folution precipitated every falt but the nitrate of nickel, nearly to drywhich had been formed by the double decomposition of the added which arfeniate of nickel and the nitrate of lead. This is the me-diffolved nitrate thod which I used in the analysis of the arseniated copper ores of nickel only; of Cornwall; and the proportion of arfenic acid, contained in arfeniate of lead, was the standard to which I referred those ores to ascertain their relative proportions. These experiments were published in the Philosophical Transactions for 1801.

netical.-Solu-

After the preceding operations, I evaporated the alcohol of evap. of the af-After the preceding operations, I evaporated the account the folition of nitrate of nickel, rediffolved the metallic falt in cohol; folition in water; prewater, and decomposed it by potash. The oxide, well washed cip. by potash; and dried, was reduced in a Hessian crucible, lined with lamp reduction with charcoal? It was black; and in that state it manifested strong magnetic proper-firongly magnetic. ties. None of the processes of the reagents I had used could Arienic being have communicated iron to the nickel. I then allayed this the magnetic fame nickel with a small proportion of arsenic in a similar cru- property. cible, and found that it had loft its magnetic property. A fmall A little iron did portion of iron added to the mass did not render it attractable. not revive it. From these experiments, often repeated, I believe that arsenic has the power of concealing the magnetic property of nickel. Cobalt appears to be in the same case. I was therefore mistaken when I announced, that those metals were not in themfelves magnetic:

On the unexpected Production of Sulphate of Magnefia. J. Bostock, M. D.

### To Mr. NICHOLSON.

SIR,

Liverpool, Nov. 18, 1802.

I FEEL fomewhat doubtful whether the subject of the following communication may appear to you sufficiently interesting, or the experiments detailed with a sufficient degree of accuracy, to entitle it to a place in your Journal. I have however fent it for your inspection; if you think it of any value, it is completely at your fervice.

I remain.

your's, &c.

JOHN BOSTOCK.

Copious faline efflorescence on the inner furface of a brick wall.

ABOUT two years ago, a house was erected in the neighbourhood of this town by Tho. Earle, Efq. upon his estate of Brook Farm. The walls, which were chiefly constructed of stone, were lined with bricks, formed of clay procured from the foundation. Some months after the erection of the building, a large part of the bricks were covered with an extremely copious, faline efflorescence. Mr. Earle felt anxious lest this peculiar fubftance should prove injurious to the plaister, with which it was proposed to cover the walls, and was defirous to be informed respecting its nature. With this intention the following experiments were performed.

Analytic experi-

The falt was diffolved in water, filtered, and gently evapoments to deter-rated; it crystallized with difficulty, and the crystals were too mine its nature. indeterminate to afcertain precifely their figure. The falt thus purified had a bitter, acrid tafte; it neither effloresced nor deliquesced after being exposed for several months to the atmosphere. It was very soluble in water; 100 grains of water at the temperature of 55°, dissolved nearly 50 grains of the falt; and the same weight of water at the boiling heat, disfolved nearly three times that quantity.

1. To

1. To a dram of the aqueous folution, 20 quarts of folution of the muriate of barites were added, and a copious, denfe, white precipitate was infantly produced.

2. Equal parts of the aqueous folution and lime water, when added together, formed a cloud which gradually fub-

fided.

3. Twenty drops of pure potash were added to a dram of the solution, and a copious precipitate was instantly produced.

- 4. To a dram of the folution were added two drams of a folution of the carbonate of potash, a slocculent precipitate was immediately produced.
- 5. A very copious precipitate was produced, when twenty drops of pure ammonia were added to a dram of the folution.
- 6. The oxalic acid produced no action when added to the folution.
- 7. Ten drops of the nitrate of filver being added to the aqueous folution, produced fcarcely any effect; a flight milkines only was perceptible in the fluid.

From these experiments the nature of the salt is sufficiently It was sulphate evident; the action of the muriate of barites proves it to be a of magnesia. Sulphate, and the experiments with lime water and the alcalis, shew that the sulphuric acid is united to magnesia. From the 6th experiment we find that it contains no lime, and from the 7th, that it contains only a very minute portion of muriatic acid. It appears therefore to be the sulphate of magnesia, or Epsom salt, in almost a perfectly pure state. The sulphate of magnesia, which is usually met with in commerce, contains a considerable proportion of muriatic acid; this is proved by the very copious precipitate which it will be found to yield upon the addition of the nitrate of silver. To the muriatic acid which it contains, Bergmann ascribes the tendency which the common sulphate of magnesia possesses, of attracting moisture from the atmosphere.

The refult of these experiments were not what I expected; I imagined that the efflorescences which we occasionally observed on walls, were composed for the most part of the nitrate of potash. The purity of the salt was also a circumstance which appeared curious, as being contrary to the state in which we usually find it. The most singular circumstance, however, was its appearing at all in that situation.

Vol. III .- DECEMBER, 1802.

·II

Upon

The falt was not found in the clay of the bricks.

Upon an examination both of the water and the clay, with which the bricks were made, there are but faint indications of the existence of this salt, at the same time that the presence of the muriatic acid is very perceptible. Is it possible that the muriatic acid may have been diffipated during the burning of the bricks, and the fulphuric generated by the union of oxigen with some sulphureous matter existing in the coals? I fear this conjecture will be thought extravagant, but at prefent I fee none which is more probable.

Conjecture.

This incident did not injure the plaister.

It is necessary to add, that no injury to the plaister appears to have arisen from the presence of this sulphate of magnesia.

#### XIII.

Abstract of a Memoir on the Dew. By CIT. BENEDICT PREVOST \*.

Well known fact that dew falls more plentifully, or in preference, on glass and not on metals.

IT is known that a glass vessel is sometimes covered with dew in the midst of a vessel of silver, which at the same time remains dry; and that mercury in a china vessel is not lowered by the dew, though the edges of the vessel are very wet; and it is very generally concluded from various experiments of this kind, that glass is a substance upon which the dew is most plentifully deposited, while it is not at all precipitated on metals.

Repetition of the experiments.

Cit. Prevoft was curious to repeat these experiments, which did not feem to him to have been made with fufficient care, in order to determine to what degree of precision it might be allowable to support the usual conclusion. In this repetition he observes some very singular facts, of which we shall proceed to give the general refults.

Difcs of metal upon glass were often changed with dew; tho' the glass wet.

1. Difcs of tinfoil, of gold, of filver, of copper, &c. being applied or fluck to plates of glass, and exposed to the dew, were often found to be no less charged with dew than the more frequently glass itself on which they were placed; though it more commonly happens that they remained dry, while the glass was very wet.

> Read to the Society of Sciences and Arts of the Department of Translated from the Annales de Lot, fitting at Montanban. Chimie, XLIV. 75.

> > 2. In

2. In this case the preserving property of the metal fre-The metal is not quently extends to a confiderable distance round, so that in causes a dry the midst of the humidity, there is constituted a zone perfectly place round it. dry and almost always well terminated. But what is very remarkable is, that when the glass is moistened underneath, the place answering to the metal remains dry; so that the and even on the influence of the metal acts through the glass of several milli-opposite surface. metres (or twentieths of an inch) in thickness. And we shall foon fee that it extends much farther.

3. The metal does not in this case act as a shelter or de- An additional fence, for if the surface be covered with glass nothing particu- glass plate does not produce the

lar happens.

4. Having pasted (collé) against the upper part of a pane of metal. glass in a window exposed to the north, and on the inner fur-being pasted on face a metallic rectangle; and on the lower part of the fame the pane of a window within; pane externally a fimilar rectangle, the author observed, that and another on a when the moisture was condensed on the inner surface, as different part often happens in winter, the inner metal fo far from remaining ture condensed dry, was more moistened than the glass, while the surface within fell most corresponding with the external metal was perfectly dry, as metal, and none well as a square of several lines round about. It was however on the surface remarked, that the humidity in the square approached much the outer; nearer the corners of the metal than in the middle of the fides of the rectangle.

3. In the same arrangement if the humidity was deposited on and moisture the outfide (by cooling) the external metal remained dry, and without fell only on the furface the place corresponding with the internal metal was on the opposite the in-

contrary rather wetter than the rest of the glass.

4. When the humidity was formed both within and without, outer metal; no moissure was deposited either upon the external metal, or moissure on both sides did not

its correspondent place within.

7. When a smaller rectangle was pasted on the outer sur-metal; or its face in the middle of the place corresponding with the inte- A small external rior rectangle, and humidity was deposited internally, the in-metal on the terior rectangle was not entirely wetted as it would else have furface opposite been, but the place corresponding with the small external the internal merectangle remained dry.

8. If upon the small external rectangle a disc of glass be opposed metal. applied, a round wet spot is formed within in the middle of A small external the dry place of which we have just spoken, corresponding to middle of the

the place occupied externally by the glass.

effect of the

without; moif-

ternal metal. and none on the wet the external tals, produced

dryness on the glass plate in the external metal, 9. If produced wetness If the whole external metal be covered with glafs, its effect ceases.

9. If instead of this last mentioned small disc, a glass of the fame fize as the external metallic dife be applied fo as exactly to cover this last, the humidity is discovered over the whole of the interior rectangle in the same manner as if the glass were bare on the outfide.

Metal upon this last glass plate revives the ef-

a. If upon the rectangle of glass there be applied another metallic rectangle, no humidity is again produced on the corresponding interior surface, though it continues to be formed on the same side on the rest of the metal and glass.

Another glass destroys it.

b. But the humidity is again condensed, if on the last external metallic rectangle there be applied a rectangle of glass.

Another metal revives it, &c.

c. And this again disappears, or ceases to be produced, by the application of a third metallic leaf.

d. A third rectangle of glass causes it again to appear.

e, f, g, &c. &c. &c.

This alternation is limited.

Nevertheless there is a term at which the phenomenon becomes irregular; and this is when the whole thickness of the metallic bases and plates of glass amounts to 16 or 20 millimetres (three quarters of an inch).

10. Though the above phenomena clearly prove that the metal does not act as a defence or cover in the production of these effects, the author proves it still more strongly in the following manner.

Gilt paper opetal if the gold be outfide; otherwise not.

He applies to a pane of glass gilt papers of the same fize and rates like a me-figure, fome by their metallic faces, and others by the furfaces which are uncovered, all the other circumstances being the fame. Those papers only of which the metallic furfaces were exposed to the air produced the effects of metals, and the others produced no effect. This experiment repeated with pieces of glass coated with tin foil instead of gilt paper, produced the fame refults.

Infide metal at a plafs caufes conglass, and not the metal.

11. If in the infide of a chamber, and parallel to one of the distance from the panes of glass in a window, there be placed a disc of metal of denfation on the fix or feven centimetres in diameter (two inches and a quarter) fo that it shall be supported only by the middle, and the face shall be only a few millimetres (or twenty-fifths of an inch) distant from the glass, when the humidity comes on within there is none deposited on the disc, unless it be extremely near, but on the opposite glass it is deposited in much greater quantity than any where elfe. 12, If

12. If instead of this disc of metal a disc of glass be substi- Glass instead of tuted, the humidity is not accumulated opposite this disc any metal has no effect. more than on the rest of the glass.

13. The inverse phenomenon takes place on the outer sur- Outside, the re-

face under like circumstances.

14. If the metallic disc (of section II.) be varnished on the Varnish on the fide which faces the interior of the chamber, the phenomenon face farthest from the glass takes place in the fame manner. has no effect

15. If the fide next the window be varnished, or if both on the fide next the glass, defides be varnished, the phenomenon does not take place. ftroys the effect.

16. These properties of the metals belong also to other Other plates as conductors of electricity, but with fome modifications accord- well as metals. ing to their nature.

17. If a plate of metal be substituted instead of a square of Metal plate inglass, humidity is sometimes produced, but always much less stead of the pane

than upon glass.

18. If this plate of metal be thin and flat, and a disc of glass Pane of glass on be fastened to it, the humidity is produced in preserence, or a plate of metal.

more abundantly on the corresponding interior surface.

19. All these phenomena take place in the same manner, or The same phenearly so, when panes of glass sitted up with the metallic discs, nomena in the open air, where or apparatus before described, are supported upon short pegs the ground rein the middle of a field where the grafs has been mowed and prefents the clipped; in these circumstances the upper surface of the glass the upper space represents the outside, and the lower the infide of the chamber. the outer air.

20. The interior humidity may be excited provided the Artificial proweather be rather cold, by means of a flove, or by fprinkling duction of dew in a chamber; water on a red hot body; and in the country during dry wea- and u ther, the space on which the glasses are to be exposed may be doors.

lightly watered after fun-fet.

21. If different liquids fuch as water, mercury, alcohol, Other bodies oils, acids, &c or even fmall leaden shot be put into glass different from metals, vessels equal and alike in all respects to the half, or two thirds of their height, and they be exposed to the dew out of a window, the humidity will not be deposited on the lower part of the veffel, but only above the level of the fubflances contained. and at a distance which varies according to the nature of these fubflances, being greater for mercury than for water, greater for water than for oil, &c.

22. In all these phenomena when the humidity is too abun- The humidity must be modedant, the refults become confused and uncertain. rate.

Difficult to fhew whether the metals differ.

23. The author made feveral experiments to afcertain whether fome of the metals might not be better adapted to the production of these phenomena than others. Having, for example, exposed to the dew plates of copper gilt and filvered, and plates of filver gilt only on the half of one of their faces, he observed nothing decifive. It only appeared to him, that in general the drops were formed larger on the white metals. But these comparative experiments are difficult to make on account of the readiness with which most of the metals become oxided in moift air; which then gives more or less of the property of glass.

Differences of temperature on the two fides.

24. In making these observations Cit. Prevost almost always kept account of the changes produced in the temperature, or the difference between the internal and external temperature. He endeavoured to procure thermometers with flat bulbs, fufficiently fenfible to point out the difference of temperature of the two fides of the glass at the same instant, which he thinks would have afforded instructive results; but he has not yet succeeded. He only remarks that it is not necesfary that the external temperature should, as is commonly thought, be more elevated than the internal, in order that humidity should be formed on the outside of the glass, but that the contrary often happens.

Farther refearches.

25. These researches may be carried farther, for example, by making fimilar experiments in vacuo, or in other gafes as well as air, and by employing other liquids instead of water.

26. In order to reduce the principal facts contained in his memoir to a small number of propositions, Cit. Prevost gives the name of the armour of contact to a metallic leaf applied or glued against the glass, and armour of distance to that which is fome millimetres off from it. This being premifed, the following are the general facts:

General statement of the facts. Armour of me-

tal on the warm face condenses moisture whether without or within;

I. When a partition or frame of glass, which separates two great masses of air at unequal temperatures, is armed in contact on its warm face.

A. If humidity be deposited on this face, it accumulates on the armour, infomuch that there is more there than elfewhere.

B. And if humidity be deposited on the opposite face, or the cold face, it accumulates on this face in the place correfponding with the armour, fo that there is more there than elfewhere.

C. The

C. The same happens when the armour is at a distance; but on the glass if then if the humidity be deposited on the warm side, it is not distant. on the armour, but it accumulates opposite to it upon the glass.

II. When the partition is armed in contact with the cold Armour on the

cold face repels moisture on

A. If humidity be deposited on this face, it is not on the both sides, armour.

B. And if humidity be deposited on the opposite face, that is to say, the warm face, there is none on the correspondent place of the armour.

C. The same happens when the armour is at a distance on if at a distance, the cold sace. But then if the humidity be deposited on this sace, there is not only a want of it on the armour, but there is none on that surface of glass opposite the armour.

III. A. In order that the phenomena should cease to take Glass or varnish place, it is sufficient that the armours of contact, or the face of prevents the effects of the armours of distance opposite the glass, should be covered mour. with glass or varnish. Nothing remarkable then happens.

B. If this glass or varnish be again covered with metal, the Alterations of

phenomena again occur.

face.

C. They cease if this metal be again covered with glass.

D. They begin, &c.

IV. One fingle proposition will include almost the whole of these facts.

When glass separates two masses of air of unequal temperatures, General proposition armour gives it (or seems to give it) the property of accumution or statement.

Leting or repelling humidity, according as it is applied either on the warm or cold sace, and this influence extends through the glass and through other substances, to the distances of several centimetres.

V. Or still more simply.

Glass which separates two masses of air of unequal temperatures, Simple general accumulates (seems to accumulate) or repells humidity, according law of those as it is armed on the warm or cold face.

#### EXPLANATION.

I imagined for a long time, fays the author, that these phenomena depended on electricity, but they may be explained very naturally by the elective attractions which are exercised at a distance, and by the well known property of the metals of being good conductors of caloric.

BASES

#### BASES OF THE EXPLANATION.

General bases for explaining them,

- 1. The less the temperature of glass is elevated, the more humidity it attracts from the air.
  - 2. Metals attract it very little.
- 3. Glass sensibly exercises its action on the humidity of air, at a distance, and notwithstanding the interposition of different bodies, such as plates of metal, &c.

4. Metals give to class, near which they are placed, the property of more speedily attracting caloric from hot air, and on the contrary, that of yielding it more speedily to cold air.

N. B. When I say that metals give glass this property, I mean that they as as if they gave it them; which is evident by an examination of two thermometers, one of mercury the other of alcohol, which are plunged at the same time in air either colder or warmer than that whose temperature they indicate. The metallic thermometer arrives much sooner than the other to that of the new medium. Its glass then, if colder, must take up more speedily from the medium the caloric which it transmits to the metal, or, if hotter, it must more speedily give out that of the metal.

Remarks.

The first basis has been long established.

The second and third are proved by the 17th and 18th sections of the extract.

The last is a necessary consequence of the conducting property of metals.

This being premifed, it is eafy to comprehend that,

The laws applied to explain all the facts.

- A. When the glass is armed on its warm face (§ 27, 1st, A. B. and C.) it yields its caloric to the cold air more speedily than that which is not armed (basis 4), and consequently it attracts humidity more powerfully (basis 1), whether directly on the glass, or through the metal, or on the metal itself (basis 3), if this be in contact; but if it be at the distance of some millimetres, the humidity not meeting the metal on its passage, accumulates on the opposite glass in a greater quantity than elsewhere.
- B. If the metal be applied on the cold fide (§ 27, II. A. and B.) the glass most heated does not attract so much humidity (bases 4 and 1), and it accumulates on the unarmed part of the pane.

C. If, in this case, the armour be covered with a plate of glass, the plate cools more speedily than if the metal were not pre-

fent;

fent; but as the partition is more heated than if it were not present, there is no effect, and the totality of the double glass armed within, is in the same case with that unarmed; it therefore accumulates neither more nor less humidity.

A fecond armour on the plate of glass will cause the phenomena to re-appear; a second plate of glass on this new armour will again make it disappear, &c. For as long as the symmetrical glass shall have the interior armours, the causes of heat and cold will be found in equilibrio; but an additional armour will necessarily destroy the balance, and the heated glass will not attract humidity; which explains the sacts of § 27. III. A. B. C. D. &c.

D. If the glass be a med on both sides (§ 7), as it would not then be exposed to the air, either on the cold or warm side, it seems that it ought to attract as much humidity on the armours as on the rest of the partition. But though the glass excreises its action through the metal, this is nevertheless an obstacle which diminishes its force; humidity in this case will not then be so strongly attracted by the doubly armed glass as by the part persectly unarmed, &c.

#### CONCLUSION.

These observations are not only interesting, but they appear Conclusions to establish an important point in philosophy; namely, that glass exercises its uttraction for the humidity (which has a tendency to be deposited from the air) through metals. See likewise § § 19 and 26 for the generalization of this proposition.

# EXPLANATION OF THE FIGURES.

Fig. 1, Plate XV. relates to Art. 4 of the abstract. A is Explanation of the interior rectangle more deeply shaded than the rest of the square, which signifies that it is more moistened. B is the place corresponding to the exterior rectangle, with its surrounding part represented white, because dry.

Fig. 2 relates to § § 7 and 8.

Fig. 3, 4, 5, 6, 7, relate to Art. 7 and 9. I c fignifies interior fide hot; E f, exterior fide cold. The lines more deeply shaded represent the section of the metallic leaves; the white lines represent the section of the glass, and the dotted shades represent drops of dew or moisture.

Fig. 8 relates to Art. 11,

# SCIENTIFIC NEWS, ACCOUNT OF BOOKS, &c.

Abstract of the late Experiments of Professor Aldini on Galvanifm.

ments of Aldini.

Galvanic experi- M. ALDINI, Professor at the Institute of Bologna, and nephew of the celebrated Galvani, after having made his experiments at the National Institute of France, has visited London, and given an accurate account of his experiments and difcoveries to the Royal Society, before whom the fame was read on the 25th laft. I have the pleasure to communicate some of the principal facts which he has had the goodness to communicate to me, which appear calculated to throw much light on fome of the most difficult phenomena of nature,

The theory that nerated in animals,

perified by Al-

by fubflituting part of an animal instead of the metallic pile. The head of an ox convulfes

part of a frog.

dini;

Various philosophers have considered the metals as not abelectricity is ge-folutely necessary for the production of galvanism, and Mr. Davy has proved it in the pile: It has also been indistinctly apprehended or conjectured in the way of theory, that the galvanic or electric matter was excited, collected, or generated in the bodies of animals, where it was confidered as the great cause or instrument of muscular motion, sensation, and other effects highly interesting, but very little understood. Profesior Aldini has the diffinguished merit of having placed this propofition in the rank of established truths. He has succeeded in exciting muscular contractions by the simple application of the nerves to the muscles of a prepared frog, without the least sufpicion of any stimulus arising from contact. He has also given motion to the limbs of a fmall cold blooded animal by the galvanic energy of an animal with warm blood; an experiment never before imagined. He takes the head of an ox recently cut off, and applying the finger of one hand wetted with falt water to the spinal marrow, he holds in the other hand the muscle of a frog prepared (that is by dissection) in such a direction that its crural nerves shall touch the cervical muscles on the tongue of the ox. Every time of this contact strong contractions are produced in the frog. If a chain of feveral persons be formed holding hands, the same effect takes place; but the contacts do not produce any effect if the chain of connection be broken or interrupted. Here then we have the most decided substitution of the organized animal system in the place of the metallic pile: it is an animal pile; and the production of the galvanic fluid, or electricity, by the direct or independent energy of life in animals, can no longer be doubted.

The Profesfor has lately repeated the series of these experi-These experiments at Oxford, and shewed in the presence of Doctors Pegg ments repeated and Bancrost, that the nerves of a prepared frog, disposed in the manner here stated, approach very sensibly to the muscles of the warm-blooded animal, and exhibit a real attraction hi- Animal attractherto unknown in natural philosophy and physiology. invites philosophers to vary and repeat this phenomenon, which has been confirmed by different philosophers, particularly by the celebrated Felice Fontana of Florence. Galvanism verified. by these facts, is shewn to be animal electricity, not merely paffive, but most probably performing the most important functions in the animal economy. This power appears not to be confined in its operation to the motion of the muscles, but also appears to be of importance in the fecretions. Aldini has given Galvanism opestrength to this conjecture, by subjecting the urine to the power rates in the seof the artificial galvanic stream, and he found it productive of a separation of the principal combined parts of that fluid, which were confidered as of much importance by the celebrated Profesfors Senebier and Jurine, of Geneva.

A very ample feries of experiments were made by Professor A. which shew the eminent and superior power of galvanism beyond any other stimulant in nature. In the months of Janu-Assonishing efary and February last, he had the courage to apply it at Bo-steets on human subgrate to the bodies of various criminals who had suffered death at that place, and by means of the pile he excited the remaining vital forces in a most assonishing manner. This stimulus Contortions of produced the most horrible contortions and grimaces by the the muscles, &c. motions of the muscles of the head and face; and an hour and a quarter after death, the arm of one of the decollated bodies the arm listed was elevated eight inches from the table on which it was sup-up-ported, and this even when a considerable weight was placed in the hand. These experiments have since been consistend in various parts of Italy, particularly at Turin, by Professors Giulio, Vassali, and Rossi.

These are not experiments of pure curiosity, but offer very These experiencouraging prospects for the benefit of mankind, in disorders ments promise of the head, and in apoplexies. Prosessor Aldini means to apply the cure of dispart orders.

part of his time in London, in communicating these important fubjects of information to physicians, as he has already done in Paris, in which place he made fome applications of his discoveries, chiefly at the Hospital de Salpetriere, in company with Doctor Pinel\*. The application of galvanifm in melancholic infanity is absolutely new, and of great interest. He perfectly, cured two patients at Bologna of this diforder; and on that account he is more defirous of recommending the trial in an affliction fo diffreshing, against which the present system of physic has fo little to offer.

Infanity cured.

Apoplexy.

drowned.

Respiration ftrongly excited by galvanism.

Precautions.

intends to publift a large work.

It appears to be equally promifing in cases of apoplexy. Recovery of the Aldini thinks it may be highly useful in recovering drowned persons, and on that account he is desirous of communicating with the Society established at London for the recovery of these unfortunate individuals. An experiment lately made at Paris adds much to his expectations. At the Hospital de la Charité he shewed the pupils, that galvanism applied to the trunk of a dog, to the spinal marrow, and the intestines, caused the lungs to act in such a manner that the air that issued from the afpera arteria twice in fuccession, extinguished a large candle placed opposite. Now, as in most cases little more is required in drowned subjects, than to set the organs of respiration into action, it is to be prefumed that the action of galvanism may be of the highest utility in these cases. Many precautions are necessary to be used in the administration of this powerful remedy in lunacy or apoplexy, which will be de-Profestor Aldini tailed in a large work which Profestor Aldini intends to publish in this country before his return to Italy. In the mean time the reader will doubtless receive fatisfaction from this short notice he has enabled me to give of his labours on a subject which promifes greatly to extend the limits of natural science, and may be reasonably expected to add to the powers which man is enabled to exert for his own benefit over the numerous beings around him.

\* Whose Treatise fur la Manie is well known in this country.

Fixamen, &c." Chemical Examination of a New Gas, composed of Hidrogen, Carbon, and Phosphorus. By J. B. TROMMS-DORFF.

Mr. Trommsdorff obtained this gas during the decomposi- A new gas by tion of phosphoric acid by ignited charcoal. In its common Trommsdoeff. state it is mixed with carbonic acid, which may be separated from it by agitation in lime water.

The new gas is nearly of the fame specific gravity as common air; it is insoluble in water, and undergoes no change when mixed with oxigen, at common temperatures; but it detonates with that aërisorm sluid by the action of heat. It is possessed no agency upon the solutions of metallic oxides which are not reducible by heat, but it decomposes the sluid saline compounds containing gold, silver, or mercury. During its combustion with oxigen, water, phosphoric acid, and carbonic acid are formed, and hence Mr. Trommsdorff is inclined to conclude that it is a triple compound of phosphorus, hidrogen, and carbon; and he proposes to call it by a name, which may be translated by the term of phosphorated carbonated hidrogen gas.

A part only of the memoir from which this account is taken is as yet published. Concerning the action of the new gas upon metallic folutions, and other phenomena presented by it, the learned author promises to enter upon some additional details. Without wishing to anticipate any of his reasoning upon the subject; with the simple hope of throwing out a hint for suture discussions, we shall venture a general observation or two in relation to it.

If hidrogen exifts in the gas, it must apparently arise from the decomposition of water contained by the concrete acid or the charcoal; for, as would appear probable from the experiments of Desormes and Clement, well burnt charcoal contains no ascertainable quantity of combined hidrogen. By separately igniting the phosphoric acid and the charcoal, before they were made to act on each other, the water contained by them would be driven off; and, under such circumstances, it would be curious to ascertain if the gas of Mr. Trommsdorff would be produced.

As phosphorus had a very strong affinity for oxigen, we should be disposed, a priori, to conclude, in reasoning upon the facts lately discovered concerning the gaseous oxide of carbon, that this substance would be formed and evolved with the carbonic acid in the process of the decomposition of phosphoric acid by charcoal. Is it not possible that the new gas may be a mixture of a triple compound of carbon, phosphorus, and oxigen, with carbonated hidrogen produced from the decomposition of water united to the primary ingredients? There is nothing in the experiments detailed in the memoir which militates against this supposition, and it might be submitted to the proof of a new experiment at the same time with the theory of the author.

D.

Notice concerning certain peculiar Habits of the Shark and Pilot Fish, by Geoffeon, Professor at the Museum of Natural History.

Interesting narrative of shark and pilot fish.

IT is afferted, that the shark has brought under his command, a very small sist of the species of the gadus, that this latter sist proceeds its master in its progress, that it points out those parts of the sea which most abound in sist, and the method of obtaining the prey of which he is most greedy; and that the shark, notwithstanding its extreme voracity, in acknowledgment for these signal services, lives on good terms with this useful companion. Naturalists, who are always guarded against the exaggerations of travellers, and not being able to account for such an association, have doubtfully rejected these facts. It will be seen that they are wrong in so doing: the observations which I have made are accompanied by circumstances, which perhaps were never presented to any one but myself in so detailed a manner.

The 6th Prairial in the year 6, I was on board the frigate Alceste between Cape Bon and the Island of Malta. The sea was calm: the passengers were tired with its long duration, when their attention was arrested by a shark which they perceived approaching the vessel. It was preceded by pilot sish, who were nearly at the same distance from one another as they

were

were from the shark: the pilot fish approached the poop of Interesting narrative vessel, they examined it twice from one end to the other, and pilot fish, and being satisfied that there was nothing there to answer their purpose, they continued the course they had formerly held. During these different movements, the shark never lost sight of them, or rather he followed them so exactly, that it seemed as if he had been drawn along.

As foon as they were discovered, a failor immediately prepared a large fish hook, which he baited with pork; but the fhark and his companions, by the time the fisher had got ready were at the distance of 20 or 25 metres; he nevertheless threw his bait into the fea. The noise occasioned by the fall was heard at a distance. The fish were assonished and stopped their course; the pilots immediately turned, and prepared again to examine the poop of the vessel. The shark, during their absence, amused himself in a thousand different ways at the furface of the water: he turned on his back, then on his belly, dived into the sea, but always appeared again in the fame place. The pilots having arrived at the poop of the Alcefte, passed near the pork, and as soon as they perceived it, they returned to the shark with greater swiftness than when they left him. When they had reached him, he began his course: the pilots swimming one on his right and the other on his left, exerted themselves to keep before him; they with difficulty did fo, and returned a fecond time to the poop of the veffel; the shark followed them, and owing to the fagacity of his companions, perceived the prey that was destined for him.

It has been faid that the shark has a very powerful smell: I paid much attention to what passed when he was near the pork: it seemed to me that he did not perceive it till the very moment that his guides, as it were, pointed it out to him; he did not till then swim with greater swiftness, or rather make a spring to seize it. He first seized a portion without being hooked; but at the second attempt the hook pierced the left lip: he was hoisted on board.

At the expiration of two hours, during which I was occupied in the anatomy of this fqualus, I began to regret that I had not more nearly examined this species, which so willingly confecrates itself to the service of the shark: I was affured

that


that I might easily do it, as it was certain that they had not quitted the neighbourhood of the vessel, and some moments after they pointed one out to me, which I perceived was of the species of pilot sish of the sailors, and of the gasterosteus ductor of naturalists.

It would doubtless be curious to examine what interests can induce animals so different in their organization, their fize and habits to form such an intercourse. Does the pilot sish feed on the dung of the shark as Cit. Bosc has supposed; or does it impose on itself the painful duties of servitude, in order to find protection and safety in the company of so voracious a species.

Notice of a New Metal, by Professor Proust.


New metal Silemes. Professor Proust, in a letter to Dr. Delametherie, promises to send an account of a new metal he has lately discovered in an ore from Hungary. He is not yet acquainted with its metallic aspect, he is apprehensive it will not be easily reduced, from its retaining oxigen with considerable avidity, and like many other metals, it is susceptible of two degrees of oxidation: the solutions of its oxide at maximum is yellow, green at the minimum, when in these two states it colours glass; lastly, it may be ranked in the class of those metals which retain oxigen against sulphurated hidrogen. He has purified it by the same means he employed with nickel, cobalt, iron, manganese.


J. de Phys. Vendemaire, an 11.






# Barometers. by the Rev. James Witson, A.M.











### . U." Read's pneumatic Apparatus.





Fig. 2.




1


Mydrometer for directly ochibiting Specific Gravities.

Fig. 1.










Muchow Sallingbott it.





Muttow Sc. Bufsell Co.









Fig. 2. S. Bed of Sandstone deposited on the broken bed of Basaltes.



Distinguishers of Galvanism & Electricity.









Muster Se. Rafaell O.






Muslow Sc. Rugsell Co!






Mullow / Raybell at





Markow . Rujavill Co.







# D. Young's Apparatus for exhibiting the Colours of thin plates.

Fig. 1.





Fig. 2.





## Prevost on the Phenomena of Dew






Fig. 2.





Fig. 3. Fig. 4. Fig. 5. Fig. 6. Fig. 7. Fig. 8.





## INDEX.

Air, experiments on condensed, 163, 164, Air, Dalton on its mechanical condenfation and rarefaction, 160 Alderson, Dr. 60 Aldini, Professor, abstract of his experiments on galvanism, 298 Amontons on the expansion of common air, 210 Apparatus for driving bolts into ships. By Mr. R. Philips, 36 Archimedes, 86 Atkins, description of his hydrometer, B. Barites, analysis of hyperoxigenized muriate of. 230 Barometer, a method of increasing its fenfibility, by J. Wilson, 21 Barthold, 98 Baumé, 51 Bells, 40 Berard on palpable mathematics, 189, 190, 191 .- Description of his new fecret lock, 216, 276 Bergman, 118, 234, 239, 289 Bernoulli, 85 Berthollet, 172, 175, 176, 177, 236, 266, 245 Befant, his water wheel, 49 Bettancourt, 207 Biftres, for enamel, obtained from mixed oxides, 151

YOL. III.

ACCUM, Mr. 123.

Acids, their action on hyperoxygenized

muriate of potash, 178, 179

Black colours for enamel, difficult to be obtained, 152 Blow pipe, description of a cheap and simple, 1-Of Haas, 119 Blues for porcelain, obtained from the oxide of cobalt, 150 Books, account of, 62 Boffut, 81 Bostock, Dr. on the theory of galvanism, 69 .- On the unexpected production of fulphate of magnefia, 288 Bournon, Count de, 90, 98 Bramah, 217 British Museum, the present state of, Brongniart, on the colours obtained from metallic oxides, 101, 148 Buat, 29, 30, 32 Von Buch, 17

C. Caloric, Henry on its materiality, 197

Carlifle, 78
Cavendish, 206
Charcoal, effect of its galvanic ignition on muriatic gas, 138
Charles, 258
Chenevix, on oxigenized and hyperoxigenized muriatic acid, 171, 229,—On the magnetic property of nickel, 286
Chladni, 98
Clement, 301
Colours vitrifiable, obtained from metallic oxides, 103.—The changes they undergo on porcelain, 106, 107

Crawford, Dr. 133, 134, 201, 202, 204 Cruickshank, 4, 11, 54, 76, 176, 183 Cuthbertson on a method of distinguishing the two kinds of electricity, 188

D.

D.

Dalton on the expansion of elastic fluids by heat, 130.—On heat and cold produced by condensation and rarefaction of air, 160.—His theory of mixed gases, 267 Davy, 4, 5, 10, 11, 12, 71, 76, 130, 198, 199, 202.—On galvanic electricity, 135
Deluc, 212, 262
Desaguliers, 187
Desormes, 301
Dew, Prevost on, 290
Duvernois, 215, 264

#### E.

Ekcherg, A. G. on the properties of the earth Yttria and on Tantalium, 251 Electricity, on the two kinds, W. N. 121 Englefield, Sir H, on the separation of light and heat by refraction, 125, 126, 127 Eytelwein, 28, 30, 32, 80, 81, 86

F,

Falconet, 99
Fluxes for porcelain colours, 104
Fontana, 299
Fortis, 17, 18
Fourcroy, 135, 236, 237
Franklin, 78, 122

#### G.

Oalvanic apparatus, theory of its action, 9.

—Remarks on, 74, 75, 76, 77

Galvanism, outline of its history, 3.—

Bostock on the theory of, 69.—Experiments on, 72, 73.—Abstract of Aldini's experiments on, 298

Ganges, 33

Garnerin, 57

Garnerin, account of his sixth descent by a parachute, 143, 144

Garnet, Dr. some account of his life, 63

Gases, dilatation of the different, by heat, 214.—Gay Lussac on the dilatations of, 134, 257.—Mr. Dalton's theory of mixed gases, 267.—Cannot be held together by chemical affinity, 270 Gay Lussac on the dilatations of gases and

vapors, 207, 257

Geoffroy, 239.—On some peculiar habits of the shark and pilot fish, 302, 303, 304

Geyer, 253

Girard, 226, 228

Glass painted, best method of baking,

Glass, Wedgwood on a method of copying paintings on, 167, 168

Gough, Mr. John, his reply to Dr. Young's theory of compound founds, 39, 146

Greens, are obtained from copper, 151 Gren, 184

Guyton, 215, 264. — On the affinities of prussic acid, 193

#### H.

Haas, description of his blow pipe, 119, 120

Haldane, Col. 3, 5

Hall, Sir James, 118

Hamilton, Dr. 226

Hand Mill, description of the Indian, 186

Hauy, 189, 190

Heat, Henry on its materiality, 197

Henry, Mr. W. on the materiality of heat, 197

Hermstadt, 89

Howard, on certain stony substances sup-

Howard, on certain stony substances supposed to have fallen on the earth, 88, 153, 243

Hoyle, 172

Hull, proposal for a commercial college at, 60, 61

Hunter, Dr. 129

Hutton, 19, 20, 113, 115

Hydraulics, Dr. Young's remarks on, 25,

Hydrometer, description of Mr. Atkins's,

I.

ī.

Invifible Girl, account of, 56

Iron, Siberian, examination of, 158.—
From South America, 158.—From Bohemia, 159. From Senegal, ib.

Jameson, on the existence of petrifactions in primitive mountains, 13.—On rocks of the trapp-formation, 111

Jordan, 284

#### K.

Keir, Mr. description of his moveable crane, 124 Kirwan, 18 Klaproth, 157, 254, 255

#### L

Lachenaic on the claying of fugars, 41
Lagrange, 84
Lambert, 166
Langfdorff, 81
Lemery, 239
Light, cheap method of producing, 222
Light, method of increasing the quantity afforded by candles. By Mr, Ezek, Walker, 272, 273, 274
Lime, analysis of hyperoxigenized muriate of, 231
Lock, description of Berard's new secret, 218, 276
Locker, 59
De Luc, 22, 23, 132

#### M

Magnefia, analyfis of the hyperoxigenized muriate of, 232 Magnefia, Bostock, Dr. on the unexpected production of sulphate of, 288 Van Marum, 7, 8, 12, 71, 74, 143 Meraud, 108, 150 Mercury, analysis of muriate of, 237 Michell, 206 Moyes, Dr. 57 Muriates, characters of the hyperoxigenized, 180

Muriatic acid, remarks on oxigenized and hyperoxigenized. By R. Chenevix, Eq. 171

Muriatic acid, analysis of hyperoxigenized, 173

Muriatic acid, Chenevix on oxigenized and hyperoxigenized, 229

Murray, Mr. J. his improvement of Woulfe's apparatus, 225

#### N.

Newton, 284 Nickel, Chenevix on its magnetic property, 286 Nuguet, 211

0.

Oil, made to boil by galvanism, 136

## P. Pearson, Rev. Mr. 234.—On some wells

near Brighthelmstone, 65, 66, 67, 68

Pendulums, their vibrations in vacuo, 35 Petrifactions, remarks on their existence in primitive mountains. By R. Jamefon, 13 Philips, Mr. Richard, his apparatus for driving bolts into ships, 36 .- Description of it, 37 Pictet, 2 .- On the reflection of obscure heat, 223 Pilkington, Mr. 66 Pinel, Dr. 300 Laplace, 209 Playfair, 14, 15, 16, 111, 114, 117 Pneumatic apparatus, description of Read's, Potash, elements of the hyperoxigenized muriate of, 174 Prevoft on the dew, 290 Priestley, 8, 213, 245, 264, 275 Prizes proposed by the National Institute of France, 139, 140, 141, 142 Prouft,

#### INDEX.

Frout, 100, 154, 236
Pruffiates, their action on the acid folutions of barites, 196
Pruffic acid, Guyton on the affinities of, 193

R

Read, his pneumatic apparatus, 55 Regnier, 217, 218, 277,1278 Rennel, 33 Richardson, Dr. 17 Richter, 184 Robison, 31, 32 Roy, Col. 131, 132, 213 Rumford, Count, 198, 199 Rutherford, Dr. 6

\$.

Saunderson, 189, 192 Sauffure, 132, 213, 266 Scheele, 240, 241 Schober, 83 Silver, formation of the hyperoxigenized muriate of, 242 Smeaton, 82 Smith, Dr. 39, 145 Soda, method of preparing the hyperoxigenized muriate of, 185 Sounds, Mr. Gough's reply to Young's theory of compound, 39 Sound, transmitted by means of a string, Sowden, Captain, 58 Spallanzani, 112 Stancari, 212 Stone from Benares, analysis of, 93, 94. -From Sienna, 95, 96. From Yorkfhire, 96 .- From Bohemia, 96, 97 Sugars, new process for claying, by Cit. Lachenaic, 41

T.

Tantalium, Ekeberg on the newly discovered metal, 251
Thermometer, depression of, in vacuo, 161
Tremery on electric sluids, 223, 234

Trevithick, 85 Trommsdorff on a new gas, 301

Vandier, 182

IV.

Vauquelin, 135, 233, 255
Venturi, 27
Volta, 6, 72
Volcanoes lunar, on the probability of their eruptions reaching the earth. By a correspondent, 255.—Observations by W. N. 256

w.

Walker, Mr. Ezekiel, on the vibrations of pendulums in vacuo, 35.—On a method of increasing the light of candles, 272, 273, 274

Water, law of its velocity, iffuing from refervoirs, 26

Water wheel, description of a, by Mr. Besant, 49

Watts, 1, 32, 268

Wedgwood, 169. On a method of painting on glass, 167

Wilson, Rev. J. on a method of increasing the fensibility of the barometer, 21 Wilson, Mr. W. on the galvanic effects of particles of zins and copper in water, 147

Wollaston, Dr. 7, 70, 123

Woltmann, 83, 87

Woulfe's apparatus, Murray's improvement of, 225

Υ.

Yellows, are formed by the oxides of lead and antimony, 149

Young, Dr. on the most useful parts of hydraulics, from Eytelwein, 25, 79.—
His reply to Mr. Gough, 145.—His apparatus for exhibiting the colours of thin plates, 283

Yttria, Ekeberg on the properties of the earth, 251

Z.

52 1184 5

Ziegler, 208

END OF THE THIRD VOLUME.

Printed by W. Stratford, Grown-Court, Temple-Bar-



